arXiv:1211.3152v2 [hep-ph] 22 Nov 2012

MAN/HEP/2012/04
November 2012

Perturbative Non-Equilibrium Thermal Field Theory

Peter Millington^{a,b} **and Apostolos Pilaftsis**^{b}

[a] *Consortium for Fundamental Physics, School of Mathematics and Statistics, University of Sheffield, Sheffield S3 7RH, United Kingdom*

[b] *Consortium for Fundamental Physics, School of Physics and Astronomy, University of Manchester, Manchester M13 9PL, United Kingdom*

E-mail: p.w.millington@sheffield.ac.uk, apostolos.pilaftsis@manchester.ac.uk

Abstract: We present a new perturbative formulation of non-equilibrium thermal field theory, based upon non-homogeneous free propagators and time-dependent vertices. Our approach to non-equilibrium dynamics yields time-dependent diagrammatic perturbation series that are free of pinch singularities, without the need to resort to quasi-particle approximation or effective resummations of finite widths. In our formalism, the avoidance of pinch singularities is a consequence of the consistent inclusion of finite-time effects and the proper consideration of the time of observation. After introducing a physically meaningful definition of particle number densities, we derive master time evolution equations for statistical distribution functions, which are valid to all orders in perturbation theory. The resulting equations do not rely upon a gradient expansion of Wigner transforms or involve any separation of time scales. To illustrate the key features of our formalism, we study out-of-equilibrium decay dynamics of unstable particles in a simple scalar model. In particular, we show how finite-time effects remove the pinch singularities and lead to violation of energy conservation at early times, giving rise to otherwise kinematically forbidden processes. The non-Markovian nature of the memory effects as predicted in our formalism is explicitly demonstrated.

Keywords: non-equilibrium thermal field theory, non-homogeneous backgrounds

ArXiv ePrint: 1211.3152

Contents

1 Introduction **2**

2 Canonical Quantization **5**

3 The CTP Formalism **9**
- 3.1 The CTP Contour 10
- 3.2 The Free CTP Propagator 16
- 3.3 The Resummed CTP Propagator 18

4 Non-Homogeneous Backgrounds **20**
- 4.1 The Schwinger–Dyson Equation in the CTP Formalism 21
- 4.2 Applicability of the Gradient Expansion 27
- 4.3 Non-Homogeneous Free Propagators 29

5 The Thermodynamic Equilibrium Limit **34**

6 Perturbative One-Loop Expansion without Pinch Singularities **38**

7 The Particle Number Density **43**

8 Master Time Evolution Equations for Particle Number Densities **46**

9 Thermalization in a Scalar Model **50**
- 9.1 Time-Dependent Width 52
- 9.2 Generalized Two-Body Decay Kinematics 54
- 9.3 Non-Markovian Oscillations 64
- 9.4 Perturbative Time Evolution Equations 66
- 9.5 Inclusion of Thermal Masses 71

10 Conclusions **73**

A Propagator Properties and Identities **75**

B Correspondence between Imaginary and Real Time Formalisms **76**

C The Complex Scalar Field **79**

D Non-Homogeneous Density Operator **83**

E Kadanoff–Baym Equations **85**

F Non-Homogeneous Loop Integrals **86**

1 Introduction

As modern particle physics continues to advance at both the energy and intensity frontiers, we are increasingly concerned with transport phenomena in dense systems of ultra-relativistic particles, the so-called *density frontier*. One such system is the deconfined phase of Quantum Chromodynamics, known as the quark-gluon plasma [1], whose existence has been inferred from the observation of jet quenching in Pb-Pb collisions by the ATLAS [2], CMS [3] and ALICE [4] experiments at the CERN Large Hadron Collider (LHC). In addition to laboratory experiments, an understanding of such ultra-relativistic many-body dynamics is of interest in theoretical astro-particle physics and cosmology. Predictions about the evolution of the early Universe rely upon our understanding of the dynamics of states at the end and shortly after the phase of cosmological inflation.

The Wilkinson Microwave Anisotropy Probe (WMAP) [5, 6] measured a baryon-to-photon ratio at the present epoch of $\eta = n_B/n_\gamma = 6.116^{+0.197}_{-0.249} \times 10^{-10}$, where n_γ is the number density of photons and $n_B = n_b - n_{\bar{b}}$ is the difference in number densities of baryons and anti-baryons. This observed Baryon Asymmetry of the Universe (BAU) is also consistent with the predictions of Big Bang Nucleosynthesis (BBN) [7]. The generation of the BAU requires the presence of out-of-equilibrium processes and the violation of baryon number (B), charge (C) and charge-parity (CP), according to the Sakharov conditions [8]. One such set of processes is prescribed by the scenarios of baryogenesis via leptogenesis [9–11], in which an initial excess in lepton number (L), provided by the decay of heavy right-handed Majorana neutrinos, is converted to a baryon number excess through the $B + L$-violating sphaleron interactions. The description of such phenomena require a consistent approach to the non-equilibrium dynamics of particle number densities. Further examples to which non-equilibrium approaches are relevant include, for instance, the phenomena of reheating and preheating [12–15] and the generation of dark matter relic densities [16].

The classical evolution of particle number densities in the early Universe is described by Boltzmann transport equations; see for instance [17–24]. A semi-classical improvement to these equations may be achieved by substituting the classical Boltzmann distributions with quantum-statistical Bose–Einstein and Fermi–Dirac distribution functions for bosons and fermions, respectively. However, such improved approaches take into account finite-width and off-shell effects by means of effective field-theoretic methods. Hence, a complete and systematic field-theoretic description of quantum transport phenomena would be desirable.

The first framework for calculating Ensemble Expectation Values (EEVs) of field operators was provided by Matsubara [25]. This so-called Imaginary Time Formalism (ITF) of thermal field theory is derived by interpreting the canonical density operator as an evolution operator in negative imaginary time. Real-time Green's functions may then be obtained by subtle analytic continuation. Nevertheless, the applicability of the ITF remains limited to the description of processes occurring in thermodynamic equilibrium.

The calculation of EEVs of operators in non-static systems is performed using the so-called Real Time Formalism (RTF); see for example [26, 27]. In particular, for non-equilibrium systems, one uses the Closed Time Path (CTP) formalism, or the *in-in* formalism, due to Schwinger and Keldysh [28, 29]. The correspondence of these results with those obtained by the ITF in the equilibrium limit are discussed extensively in the literature [30–36]. A non-perturbative loopwise expansion of the CTP generating functional is then provided by the Cornwall–Jackiw–Tomboulis (CJT) effective action [37, 38], which was subsequently applied to the CTP formalism by Calzetta and Hu [39, 40].

Recently, the computation of the out-of-equilibrium evolution of particle number densities has received much attention, where several authors put forward quantum-corrected or quantum Boltzmann equations [41–50]. These approaches rely upon the Wigner transformation and gradient expansion [51] of a system of Kadanoff–Baym (KB) [52, 53] equations, originally applied in the non-relativistic regime [54, 55]. Often the truncation of the gradient expansion is accompanied by quasi-particle ansaetze for the forms of the propagators. Similar approaches have recently been applied to the quark-gluon plasma [56]. Dynamical equations may also be derived using functional renormalization techniques [57].

In this article, we present a new approach to non-equilibrium thermal field theory. Our approach is based upon a diagrammatic perturbation series, constructed from non-homogeneous free propagators and time-dependent vertices, which explicitly break space-time translational invariance. In particular, we show how the systematic inclusion of finite-time effects and the proper consideration of the time of observation render our perturbative expansion free of the pinch singularities, thereby enabling a consistent treatment of non-equilibrium dynamics. Unlike other methods, our approach does not require the use of quasi-particle approximation or other effective resummations of finite-width effects.

A key element of our formalism has been to define physically meaningful particle number densities in terms of off-shell Green's functions. This definition is unambiguous, as it can be closely linked to Noether's charge, associated with a partially conserved current. Subsequently, we derive master time evolution equations for the statistical distribution functions related to particle number densities. These time evolution equations do not rely on the truncation of a gradient expansion of the so-called Wigner transforms, neither do they involve separation of various time scales. Instead, they are built of non-homogeneous free propagators, with modified time-dependent Feynman rules, which enable us to analyze the pertinent kinematics fully. Our analysis shows that the systematic inclusion of finite-time effects leads to the microscopic violation of energy conservation at early times. Aside from preventing the appearance of pinch singularities, the effect of a finite time interval of evolution leads to contributions from processes that would otherwise be kinematically disallowed on grounds of energy conservation. Applying this formalism to a simple scalar model with unstable particles, we show that these evanescent processes contribute significantly to the rapidly-oscillating transient evolution of these systems, inducing late-time memory effects. We find that the spectral evolution of two-point correlation functions exhibits an oscillating pattern with time-varying frequencies. Such an oscillating pattern signifies a non-Markovian evolution of memory effects, which is a distinctive feature governing truly out-of-thermal-equilibrium dynamical systems.

The outline of the paper is as follows. In Section 2, we review the canonical quantization of a scalar field theory, placing particular emphasis on the inclusion of a finite time of coincidence for the three equivalent pictures of quantum mechanics, namely the Schrödinger, Heisenberg and Dirac (interaction) pictures. In Section 3, we introduce the

T	thermodynamic temperature
t	macroscopic time
$\tilde{t}$	microscopic real time
τ	microscopic negative imaginary time
$\mathfrak{t}$	complex time
$\mathrm{T}(\bar{\mathrm{T}})$	time-(anti-time-)ordering operator
$\mathrm{T}_{\mathcal{C}}$	path-ordering operator
Z	wavefunction renormalization
$\mathcal{Z}$	partition function/ generating functional
f	statistical distribution function
f_β	Boltzmann distribution
f_{B}	Bose–Einstein distribution
$\tilde{f}$	ensemble function
ρ	density operator/ density matrix
n	number density
N	total particle number

Table 1: Glossary for clarifying polysemous notation.

CTP formalism, limiting ourselves initially to consider its application to quantum field theory at zero temperature and density. This is followed by a discussion of the constraints upon the form of the CTP propagator. With these prerequisites reviewed, we proceed in Section 4 to discuss the generalization of the CTP formalism to finite temperature and density in the presence of both spatially and temporally inhomogeneous backgrounds. In the same section, we derive the most general form of the non-homogeneous free propagators for a scalar field theory. The thermodynamic equilibrium limit is outlined in Section 5. In Section 6, we demonstrate the absence of pinch singularities in the perturbation series arising in our approach. Subsequently, in Section 7, we define the concept of particle number density and relate this to a perturbative loopwise expansion of the resummed CTP propagators. In Section 8, we derive *new* master time evolution equations for statistical distribution functions, which go over to classical Boltzmann equations in the appropriate limits. Section 9 studies the thermalization of unstable particles in a simple scalar model, where particular emphasis is laid on the early-time behaviour and the impact of the finite-time effects. Finally, our conclusions and possible future directions are presented in Section 10.

For clarity, a glossary that might be useful to the reader to clarify polysemous notation is given in Table 1. Appendix A provides a summary of all important propagator definitions, along with their basic relations and properties. In Appendix B, we describe the correspondence between the RTF and ITF in the equilibrium limit at the one-loop level for a real scalar field theory with a cubic self-interaction. The generalization of our approach to complex scalar fields is outlined in Appendix C. Appendix D contains a series expansion of the most general non-homogeneous Gaussian-like density operator. In Appendix E, we summarize the derivation of the so-called Kadanoff–Baym equations. Lastly, in Appendix F, we describe key technical details involved in the calculation of loop integrals with non-homogeneous free propagators.

2 Canonical Quantization

In this section, we review the basic results obtained within the canonical quantization formalism for a massive scalar field theory. This discussion will serve as a precursor for our generalization to finite temperature and density, which follows in subsequent sections.

As a starting point, we consider a simple self-interacting theory of a real scalar field $\Phi(x)$ with mass M, which is described by the Lagrangian

$$\mathcal{L}(x) \;=\; \tfrac{1}{2}\partial_\mu\Phi(x)\partial^\mu\Phi(x) \;-\; \tfrac{1}{2}M^2\Phi^2(x) \;-\; \tfrac{1}{3!}g\Phi^3(x) \;-\; \tfrac{1}{4!}\lambda\Phi^4(x)\;, \tag{2.1}$$

where g and λ are dimensionful and dimensionless couplings, respectively. Throughout this article, we use the short-hand notation: $x \equiv x^\mu = (x^0, \mathbf{x})$, for the four-dimensional space-time arguments of field operators, and adopt the signature $(+,-,-,-)$ for the Minkowski space-time metric $\eta_{\mu\nu}$.

It proves convenient to start our canonical quantization approach in the Schrödinger picture, where the state vectors are *time-dependent*, whilst basis vectors and operators are, in the absence of external sources, *time-independent*. Hence, the *time-independent* Schrödinger-picture field operator, denoted by a subscript S, may be written in the familiar plane-wave decomposition

$$\Phi_{\mathrm{S}}(\mathbf{x};\tilde{t}_i) \;=\; \int \mathrm{d}\Pi_{\mathbf{p}}\;\Big(a_{\mathrm{S}}(\mathbf{p};\tilde{t}_i)e^{i\mathbf{p}\cdot\mathbf{x}} \;+\; a_{\mathrm{S}}^\dagger(\mathbf{p};\tilde{t}_i)e^{-i\mathbf{p}\cdot\mathbf{x}}\Big)\;, \tag{2.2}$$

where we have introduced the short-hand notation:

$$\int \mathrm{d}\Pi_{\mathbf{p}} \;\equiv\; \int\frac{\mathrm{d}^3\mathbf{p}}{(2\pi)^3}\,\frac{1}{2E(\mathbf{p})} \;=\; \int\frac{\mathrm{d}^4 p}{(2\pi)^4}\;2\pi\theta(p_0)\delta(p^2-M^2)\;, \tag{2.3}$$

for the Lorentz-invariant phase-space (LIPS). In (2.3), $E(\mathbf{p}) \;=\; \sqrt{\mathbf{p}^2+M^2}$ is the energy of the single-particle mode with three-momentum $\mathbf{p}$ and $\theta(p_0)$ is the generalized unit step function, defined by the Fourier representation

$$\theta(p_0) \;\equiv\; i\int_{-\infty}^{+\infty}\frac{\mathrm{d}\xi}{2\pi}\;\frac{e^{-ip_0\xi}}{\xi\;+\;i\epsilon} \;=\; \begin{cases} 1\;, & p_0 \;>\; 0\;; \\ \frac{1}{2}\;, & p_0 \;=\; 0\;; \\ 0\;, & p_0 \;<\; 0\;, \end{cases} \tag{2.4}$$

where $\epsilon = 0^+$. It is essential to stress here that we define the Schrödinger-, Heisenberg- and Interaction (Dirac)-pictures to be coincident at the finite *microscopic* boundary time $\tilde{t}_i$, such that

$$\Phi_{\mathrm{S}}(\mathbf{x};\tilde{t}_i) \;=\; \Phi_{\mathrm{H}}(\tilde{t}_i,\mathbf{x};\tilde{t}_i) \;=\; \Phi_{\mathrm{I}}(\tilde{t}_i,\mathbf{x};\tilde{t}_i)\;, \tag{2.5}$$

where implicit dependence upon the boundary time $\tilde{t}_i$ is marked by separation from explicit arguments by a semi-colon.

The *time-independent* Schrödinger-picture operators $a_{\mathrm{S}}^\dagger(\mathbf{p};\tilde{t}_i)$ and $a_{\mathrm{S}}(\mathbf{p};\tilde{t}_i)$ are the usual creation and annihilation operators, which act on the stationary vacuum $|0\rangle$, respectively creating and destroying *time-independent* single-particle momentum eigenstates.

Their defining properties are:

$$\begin{aligned} a^\dagger_{\rm S}(\mathbf{p};\tilde{t}_i)|0\rangle \ &= \ |\mathbf{p};\tilde{t}_i\rangle\ , & (2.6a)\\ a_{\rm S}(\mathbf{p};\tilde{t}_i)|\mathbf{p}';\tilde{t}_i\rangle \ &= \ (2\pi)^3 2E(\mathbf{p})\delta^{(3)}(\mathbf{p}\ -\ \mathbf{p}')|0\rangle\ , & (2.6b)\\ a_{\rm S}(\mathbf{p};\tilde{t}_i)|0\rangle \ &= \ 0\ . & (2.6c) \end{aligned}$$

Note that the momentum eigenstates $|\mathbf{p};\tilde{t}_i\rangle$ respect the orthonormality condition

$$\langle\mathbf{p}';\tilde{t}_i|\mathbf{p};\tilde{t}_i\rangle \ = \ (2\pi)^3 2E(\mathbf{p})\delta^{(3)}(\mathbf{p}\ -\ \mathbf{p}')\ . \tag{2.7}$$

We then define the *time-dependent* interaction-picture field operator $\Phi_{\rm I}(x;\tilde{t}_i)$ via

$$\Phi_{\rm I}(x;\tilde{t}_i) \ = \ e^{iH^0_{\rm S}(x_0\,-\,\tilde{t}_i)}\Phi_{\rm S}(\mathbf{x};\tilde{t}_i)e^{-iH^0_{\rm S}(x_0\,-\,\tilde{t}_i)}\ , \tag{2.8}$$

where $H^0_{\rm S}$ is the free-part of the Hamiltonian in the Schrödinger picture. Using the commutators

$$\begin{aligned} \big[\,H^0_{\rm S},\,a_{\rm S}(\mathbf{p};\tilde{t}_i)\,\big] \ &= \ -E(\mathbf{p})a_{\rm S}(\mathbf{p};\tilde{t}_i)\ , & (2.9a)\\ \big[\,H^0_{\rm S},\,a^\dagger_{\rm S}(\mathbf{p};\tilde{t}_i)\,\big] \ &= \ +E(\mathbf{p})a^\dagger_{\rm S}(\mathbf{p};\tilde{t}_i)\ , & (2.9b) \end{aligned}$$

the interaction-picture field operator may be written

$$\Phi_{\rm I}(x;\tilde{t}_i) \ = \ \int\!{\rm d}\Pi_{\mathbf{p}}\,\Big(a_{\rm S}(\mathbf{p};\tilde{t}_i)e^{-iE(\mathbf{p})(x_0\,-\,\tilde{t}_i)}e^{i\mathbf{p}\cdot\mathbf{x}}\ +\ a^\dagger_{\rm S}(\mathbf{p};\tilde{t}_i)e^{iE(\mathbf{p})(x_0\,-\,\tilde{t}_i)}e^{-i\mathbf{p}\cdot\mathbf{x}}\Big)\ , \tag{2.10}$$

or equivalently, in terms of interaction-picture operators only,

$$\Phi_{\rm I}(x;\tilde{t}_i) \ = \ \int\!{\rm d}\Pi_{\mathbf{p}}\,\Big(a_{\rm I}(\mathbf{p},0;\tilde{t}_i)e^{-iE(\mathbf{p})x_0}e^{i\mathbf{p}\cdot\mathbf{x}}\ +\ a^\dagger_{\rm I}(\mathbf{p},0;\tilde{t}_i)e^{iE(\mathbf{p})x_0}e^{-i\mathbf{p}\cdot\mathbf{x}}\Big)\ . \tag{2.11}$$

Notice that in (2.11) the *time-dependent* interaction-picture creation and annihilation operators, $a^\dagger_{\rm I}(\mathbf{p},\tilde{t};\tilde{t}_i)$ and $a_{\rm I}(\mathbf{p},\tilde{t};\tilde{t}_i)$, are evaluated at the microscopic time $\tilde{t}\ =\ 0$, after employing a relation analogous to (2.8). We may write the four-dimensional Fourier transform of the interaction-picture field operator as

$$\Phi_{\rm I}(p;\tilde{t}_i) \ = \ \int\!{\rm d}^4x\ e^{ip\cdot x}\,\Phi_{\rm I}(x;\tilde{t}_i) \ = \ 2\pi\delta(p^2-M^2)\Big(\theta(p_0)a_{\rm I}(\mathbf{p},0;\tilde{t}_i)+\theta(-p_0)a^\dagger_{\rm I}(-\mathbf{p},0;\tilde{t}_i)\Big)\ . \tag{2.12}$$

In the limit where the interactions are switched off adiabatically as $\tilde{t}_i\ \to\ -\infty$, one may define the asymptotic *in* creation and annihilation operators via

$$a^{(\dagger)}_{\rm in}(\mathbf{p}) \ \equiv \ Z^{-1/2}\lim_{\tilde{t}_i\,\to\,-\infty}a^{(\dagger)}_{\rm I}(\mathbf{p},0;\tilde{t}_i) \ = \ Z^{-1/2}\lim_{\tilde{t}_i\,\to\,-\infty}a^{(\dagger)}_{\rm S}(\mathbf{p};\tilde{t}_i)e^{+(-)iE(\mathbf{p})\tilde{t}_i}\ , \tag{2.13}$$

where $Z\ =\ 1\ +\ \mathcal{O}(\hbar)$ is the wavefunction renormalization. Evidently, keeping track of the finite boundary time $\tilde{t}_i$ plays an important role in ensuring that our forthcoming generalization to perturbative thermal field theory remains consistent with asymptotic field theory in the limit $\tilde{t}_i\ \to\ -\infty$. Hereafter, we will omit the subscript I on interaction-picture operators and suppress the implicit dependence on the boundary time $\tilde{t}_i$, except where it is necessary to do otherwise for clarity.

We start our quantization procedure by defining the commutator of interaction-picture fields

$$\big[\,\Phi(x),\ \Phi(y)\,\big]\ \equiv\ i\Delta^0(x,y;M^2)\ , \tag{2.14}$$

where $i\Delta^0(x,y;M^2)$ is the *free* Pauli-Jordan propagator. Herein, we denote free propagators by a superscript 0. The condition of micro-causality requires that the interaction-picture fields commute for space-like separations $(x-y)^2 < 0$. This restricts $i\Delta^0(x,y;M^2)$ to be invariant under spatial translations, having the Poincaré-invariant form

$$i\Delta^0(x,y;M^2)\ =\ \int \mathrm{d}\Pi_{\mathbf{p}}\ \Big(e^{-iE(\mathbf{p})(x^0\,-\,y^0)}e^{i\mathbf{p}\cdot(\mathbf{x}\,-\,\mathbf{y})}\ -\ \big(x\ \longleftrightarrow\ y\big)\Big)\ . \tag{2.15}$$

Observe that $i\Delta^0(x,y;M^2)$ represents the difference of two counter-propagating packets of plane waves and vanishes for $(x\ -\ y)^2\ <\ 0$.

It proves useful for our forthcoming analysis to introduce the *double* Fourier transform

$$i\Delta^0(p,p';M^2)\ =\ \iint \mathrm{d}^4x\,\mathrm{d}^4y\ e^{ip\cdot x}e^{-ip'\cdot y}\,i\Delta^0(x,y;M^2) \tag{2.16a}$$

$$=\ 2\pi\varepsilon(p_0)\delta(p^2-M^2)(2\pi)^4\delta^{(4)}(p\ -\ p')\ , \tag{2.16b}$$

where $\varepsilon(p_0)\ \equiv\ \theta(p_0)\ -\ \theta(-p_0)$ is the generalized signum function. Note that we have defined the Fourier transforms such that four-momentum p flows *away from* the point x and four-momentum p' flows *towards* the point y.

From (2.14) and (2.15), we may derive the equal-time commutation relations

$$i\Delta^0(x,y;M^2)\big|_{x^0\,=\,y^0\,=\,\tilde{t}}\ =\ \big[\,\Phi(\tilde{t},\mathbf{x}),\ \Phi(\tilde{t},\mathbf{y})\,\big]\ =\ 0\ , \tag{2.17a}$$

$$\partial_{x_0}i\Delta^0(x,y;M^2)\big|_{x^0\,=\,y^0\,=\,\tilde{t}}\ =\ \big[\,\pi(\tilde{t},\mathbf{x}),\ \Phi(\tilde{t},\mathbf{y})\,\big]\ =\ -i\delta^{(3)}(\mathbf{x}\ -\ \mathbf{y})\ , \tag{2.17b}$$

$$\partial_{x_0}\partial_{y_0}i\Delta^0(x,y;M^2)\big|_{x^0\,=\,y^0\,=\,\tilde{t}}\ =\ \big[\,\pi(\tilde{t},\mathbf{x}),\ \pi(\tilde{t},\mathbf{y})\,\big]\ =\ 0\ , \tag{2.17c}$$

where $\pi(x)\ =\ \partial_{x_0}\Phi(x)$ is the conjugate-momentum operator. In order to satisfy the canonical quantization relations (2.17), the creation and annihilation operators must respect the commutation relation

$$\big[\,a(\mathbf{p},\tilde{t}\,),\ a^\dagger(\mathbf{p}',\tilde{t}'\,)\,\big]\ =\ (2\pi)^3 2E(\mathbf{p})\delta^{(3)}(\mathbf{p}\ -\ \mathbf{p}')e^{-iE(\mathbf{p})(\tilde{t}\,-\,\tilde{t}')}\ , \tag{2.18}$$

with all other commutators vanishing. Here, we emphasize the appearance of an overall phase $e^{-iE(\mathbf{p})(\tilde{t}\,-\,\tilde{t}')}$ in (2.18) for $\tilde{t}\ \neq\ \tilde{t}'$ [cf. Section 4.1].

The vacuum expectation value of the commutator of Heisenberg-picture field operators may be expressed as a superposition of interaction-picture field commutators by means of the Källén–Lehmann spectral representation [58, 59]:

$$\langle 0|\big[\,\Phi_{\mathrm{H}}(x),\ \Phi_{\mathrm{H}}(y)\,\big]|0\rangle\ \equiv\ i\Delta(x,y)\ =\ \int_0^\infty \mathrm{d}s\ \sigma(s)\,i\Delta^0(x,y;s)\ , \tag{2.19}$$

where $i\Delta^0(x,y;s)$ is the free Pauli–Jordan propagator in (2.15) with M^2 replaced by s and $i\Delta(x,y)$ is the *dressed* or *resummed* propagator. The positive spectral density $\sigma(s)$

contains all information about the spectrum of single- and multi-particle states produced by the Heisenberg-picture field operators $\Phi_{\rm H}$. Note that for a homogeneous and stationary vacuum $|0\rangle$, $\sigma(s)$ is independent of the space-time coordinates and the resummed Pauli–Jordan propagator maintains its translational invariance. If $\sigma(s)$ is normalized, such that

$$\int_0^\infty {\rm d}s\, \sigma(s) \ =\ 1\,, \tag{2.20}$$

the equal-time commutation relations of Heisenberg-picture operators maintain exactly the form in (2.17). In this case, the spectral function cannot depend upon any fluctuations in the background. Clearly, for non-trivial 'vacua', or thermal backgrounds, the spectral density becomes a function also of the coordinates. The spectral representation of the resummed propagators may then depend non-trivially on space-time coordinates, i.e. $\sigma = \sigma(s;x,y)$; see for instance [41]. In this case, the convenient factorization of the Källén–Lehmann representation breaks down and the equal-time commutation relations of Heisenberg-picture field operators do not maintain their canonical form.

The retarded and advanced causal propagators are defined as

$$i\Delta_{\rm R}(x,y)\ \equiv\ \theta(x_0\ -\ y_0)i\Delta(x,y)\,,\qquad i\Delta_{\rm A}(x,y)\ \equiv\ -\theta(y_0\ -\ x_0)i\Delta(x,y)\,. \tag{2.21}$$

Using the Fourier representation of the unit step function in (2.4), we introduce a convenient representation of these causal propagators in terms of the convolution

$$i\Delta_{\rm R(A)}(p,p')\ =\ i\int\frac{{\rm d}k_0}{2\pi}\,\frac{i\Delta(p_0-k_0,p_0'-k_0;\mathbf{p},\mathbf{p}')}{k_0\ +(-)\,i\epsilon}\ . \tag{2.22}$$

The absolutely-ordered Wightman propagators are defined as

$$i\Delta_>(x,y)\ \equiv\ \langle\,\Phi(x)\Phi(y)\,\rangle\,,\qquad i\Delta_<(x,y)\ \equiv\ \langle\,\Phi(y)\Phi(x)\,\rangle\,. \tag{2.23}$$

We note that the two-point correlation functions $\Delta(x,y)$, $\Delta_{>,<}(x,y)$ and $\Delta_{\rm R,A}(x,y)$ satisfy the *causality relation*:

$$\Delta(x,y)\ =\ \Delta_>(x,y)\ -\ \Delta_<(x,y)\ =\ \Delta_{\rm R}(x,y)\ -\ \Delta_{\rm A}(x,y)\,. \tag{2.24}$$

Our next step is to define the non-causal Hadamard propagator, which is the vacuum expectation value of the field anti-commutator

$$i\Delta_1(x,y)\ \equiv\ \langle\,\{\,\Phi(x),\ \Phi(y)\,\}\,\rangle\,. \tag{2.25}$$

Correspondingly, the time-ordered Feynman and anti-time-ordered Dyson propagators are given by

$$i\Delta_{\rm F}(x,y)\ \equiv\ \langle\,{\rm T}\left[\,\Phi(x)\Phi(y)\,\right]\rangle\,,\qquad i\Delta_{\rm D}(x,y)\ \equiv\ \langle\,\bar{\rm T}\left[\,\Phi(x)\Phi(y)\,\right]\rangle\,, \tag{2.26}$$

where T and $\bar{\rm T}$ are the time- and anti-time-ordering operators, respectively. Explicitly, $\Delta_{\rm F}(x,y)$ and $\Delta_{\rm D}(x,y)$ may be written in terms of the absolutely-ordered Wightman propagators $\Delta_>(x,y)$ and $\Delta_<(x,y)$ as

$$\Delta_{\rm F}(x,y)\ =\ \theta(x_0\ -\ y_0)\Delta_>(x,y)\ +\ \theta(y_0\ -\ x_0)\Delta_<(x,y)\,, \tag{2.27a}$$

$$\Delta_{\rm D}(x,y)\ =\ \theta(x_0\ -\ y_0)\Delta_<(x,y)\ +\ \theta(y_0\ -\ x_0)\Delta_>(x,y)\,. \tag{2.27b}$$

The propagators $\Delta_1(x,y)$, $\Delta_{>,<}(x,y)$ and $\Delta_{\mathrm{F,D}}(x,y)$ obey the *unitarity relations*:

$$\Delta_1(x,y) \ = \ \Delta_{\mathrm{F}}(x,y) \ + \ \Delta_{\mathrm{D}}(x,y) \ = \ \Delta_{>}(x,y) \ + \ \Delta_{<}(x,y) \ = \ 2i\mathrm{Im}\,\Delta_{\mathrm{F}}(x,y)\ . \tag{2.28}$$

Finally, for completeness, we define the principal-part propagator

$$\Delta_{\mathcal{P}}(x,y) \ = \ \frac{1}{2}\Big(\Delta_{\mathrm{R}}(x,y) \ + \ \Delta_{\mathrm{A}}(x,y)\Big) \ = \ \mathrm{Re}\,\Delta_{\mathrm{F}}(x,y)\ . \tag{2.29}$$

Here, we should bear in mind that

$$\mathrm{Re}\big(\mathrm{Im}\big)\,\Delta_{\mathrm{F}}(x,y) \ \neq \ \iint\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4p'}{(2\pi)^4}\,e^{-ip\cdot x}e^{ip'\cdot y}\,\mathrm{Re}\big(\mathrm{Im}\big)\,\Delta_{\mathrm{F}}(p,p')\ , \tag{2.30}$$

implying that

$$\Delta_{\mathcal{P}}(p,p') \ \neq \ \mathrm{Re}\,\Delta_{\mathrm{F}}(p,p')\ , \tag{2.31}$$

unless $\Delta_{\mathrm{F}}(p,p') \ = \ \Delta_{\mathrm{F}}(-p,-p')$, which is not generally true in non-equilibrium thermal field theory [cf. (A.4f)].

The definitions and the relations discussed above are valid for both free and resummed propagators. In Appendix A, we list the properties of these propagators in coordinate, momentum and Wigner (see Section 4.2) representations, as well as a number of useful identities, which we will refer to throughout this article. More detailed discussion related to these propagators and their contour-integral representations may be found in [60]. In Appendix C, these considerations and the analysis of the following sections are generalized to the complex scalar field.

3 The CTP Formalism

In this section, we review the Closed Time Path (CTP) formalism, or the so-called *in-in* formalism, due to Schwinger and Keldysh [28, 29]. As an illuminating exercise, we consider the CTP formalism in the context of zero-temperature quantum field theory and derive the associated 2 × 2 matrix propagator, obeying basic field-theoretic constraints, such as CPT invariance, Hermiticity, causality and unitarity. Finally, we discuss the properties of the resummed propagator in the CTP formalism.

In the calculation of scattering-matrix elements, we are concerned with the transition amplitude between *in* and *out* asymptotic states, where single-particle states are defined in the infinitely-distant past and future. On the other hand, in quantum statistical mechanics, we are interested in the calculation of Ensemble Expectation Values (EEVs) of operators at a fixed given time t. Specifically, the evaluation of EEVs of operators requires a field-theoretic approach that allows us to determine the transition amplitude between states evolved to the *same time*. This approach is the Schwinger–Keldysh CTP formalism, which we describe in detail below.

For illustration, let us consider the following observable $\mathcal{O}$ in the Schrödinger picture:

$$\mathcal{O}_{(\mathrm{S})}(\tilde{t}_f;\tilde{t}_i) \ = \ \int[\mathrm{d}\Phi]\ {}_{\mathrm{S}}\langle\Phi(\mathbf{z}),\tilde{t}_f;\tilde{t}_i|\Phi_{\mathrm{S}}(\mathbf{x};\tilde{t}_i)\Phi_{\mathrm{S}}(\mathbf{y};\tilde{t}_i)|\Phi(\mathbf{z}),\tilde{t}_f;\tilde{t}_i\rangle_{\mathrm{S}}\ , \tag{3.1}$$

where $[\mathrm{d}\Phi]$ represents the functional integral over all field configurations $\Phi(\mathbf{z})$. In (3.1), $|\Phi(\mathbf{z}), \tilde{t}_f; \tilde{t}_i\rangle_{\mathrm{S}}$ is a *time-dependent* eigenstate of the *time-independent* Schrödinger-picture field operator $\Phi_{\mathrm{S}}(\mathbf{x}; \tilde{t}_i)$ with eigenvalue $\Phi(\mathbf{x})$, where the implicit dependence upon the boundary time $\tilde{t}_i$ has been restored.

We should remark here that there are seven independent space-time coordinates involved in the observable (3.1). These are the six spatial coordinates, $\mathbf{x}$ and $\mathbf{y}$, *plus* the microscopic time $\tilde{t}_f$. In addition, there is one implicit coordinate: the boundary time $\tilde{t}_i$. As we will see, exactly seven independent coordinates are required to construct physical observables that are compatible with Heisenberg's uncertainty principle. We choose the seven independent coordinates to be

$$t \;=\; \tilde{t}_f \,-\, \tilde{t}_i\;, \qquad \mathbf{X} \;=\; \frac{1}{2}\Big(\mathbf{x} \,+\, \mathbf{y}\Big)\;, \qquad \mathbf{p}\;, \tag{3.2}$$

where t and $\mathbf{X}$ are the macroscopic time and central space coordinates and $\mathbf{p}$ is the Fourier-conjugate variable to the relative spatial coordinate $\mathbf{R} \,=\, \mathbf{x} \,-\, \mathbf{y}$.

In the interaction picture, the same observable $\mathcal{O}$ in (3.1) is given by

$$\mathcal{O}_{(\mathrm{I})}(\tilde{t}_f; \tilde{t}_i) \;=\; \int [\mathrm{d}\Phi]\;\, {}_{\mathrm{I}}\langle \Phi(\mathbf{z}), \tilde{t}_f; \tilde{t}_i | \Phi_{\mathrm{I}}(\tilde{t}_f, \mathbf{x}; \tilde{t}_i) \Phi_{\mathrm{I}}(\tilde{t}_f, \mathbf{y}; \tilde{t}_i) | \Phi(\mathbf{z}), \tilde{t}_f; \tilde{t}_i\rangle_{\mathrm{I}}\;; \tag{3.3}$$

and, in the Heisenberg picture, by

$$\mathcal{O}_{(\mathrm{H})}(\tilde{t}_f; \tilde{t}_i) \;=\; \int [\mathrm{d}\Phi]\;\, {}_{\mathrm{H}}\langle \Phi(\mathbf{z}); \tilde{t}_i | \Phi_{\mathrm{H}}(\tilde{t}_f, \mathbf{x}; \tilde{t}_i) \Phi_{\mathrm{H}}(\tilde{t}_f, \mathbf{y}; \tilde{t}_i) | \Phi(\mathbf{z}); \tilde{t}_i\rangle_{\mathrm{H}}\;. \tag{3.4}$$

Notice that the prediction for the observable $\mathcal{O}$ does not depend on which picture we are using, i.e. $\mathcal{O}_{(\mathrm{S})}(\tilde{t}_f; \tilde{t}_i) \,=\, \mathcal{O}_{(\mathrm{I})}(\tilde{t}_f; \tilde{t}_i) \,=\, \mathcal{O}_{(\mathrm{H})}(\tilde{t}_f; \tilde{t}_i)$. This picture-independence of $\mathcal{O}$ is only possible because the time-dependent vectors and operators in $\mathcal{O}$ are evaluated individually at *equal times*. Otherwise, any potential observable, built out of time-dependent vectors and operators that are evaluated at *different times*, would be *picture-dependent* and therefore *unphysical*. Moreover, the prediction of the observable $\mathcal{O}$ should be invariant under time translations, i.e.

$$\mathcal{O}(\tilde{t}_f; \tilde{t}_i) \;=\; \mathcal{O}(t; 0) \;\equiv\; \mathcal{O}(t), \tag{3.5}$$

where $t \,=\, \tilde{t}_f - \tilde{t}_i$ is the macroscopic time, as we will see below. Herein and throughout the remainder of this article, the time agruments of time translationally-invariant quantities are written in terms of the macroscopic time t only.

3.1 The CTP Contour

In order to evaluate equal-time observables of the form in (3.4), we first introduce the *in* vacuum state $|0_{\mathrm{in}}; \tilde{t}_i\rangle$, which is at time $\tilde{t}_i$ a time-independent eigenstate of the Heisenberg field operator $\Phi_{\mathrm{H}}(x; \tilde{t}_i)$ with zero eigenvalue; see [39, 40]. We then need a means of driving the amplitude $\langle 0_{\mathrm{in}}; \tilde{t}_i | 0_{\mathrm{in}}; \tilde{t}_i\rangle$, which can be achieved by the appropriate introduction of external sources.

The procedure may be outlined with the aid of Figure 1 as follows. We imagine evolving our *in* state at time $\tilde{t}_i$ forwards in time under the influence of a source $J_+(x)$ to


$\mathrm{Im}\,\mathfrak{t}$
$\tilde z(0) \;=\; \tilde t_i$
$\mathcal{C}_+$
$\mathrm{Re}\,\mathfrak{t}$
$\tilde z(1) \;=\; \tilde t_i - i\epsilon$
$\mathcal{C}_-$
$\tilde z(1/2) \;=\; \tilde t_f - i\epsilon/2$
macroscopic time $t \;=\; \mathrm{Re}\,\tilde z(u) \;-\; \tilde t_i$
initial conditions:
macroscopic time $t \;=\; 0$
observation:
macroscopic time $t \;=\; \tilde t_f \;-\; \tilde t_i$

Figure 1: The closed-time path, $\mathcal{C} \;=\; \mathcal{C}_+ \cup \mathcal{C}_-$, running first along $\mathcal{C}_+$ from $\tilde t_i$ to $\tilde t_f - i\epsilon/2$ and then returning along $\mathcal{C}_-$ from $\tilde t_f - i\epsilon/2$ to $\tilde t_i - i\epsilon$. The relationship between *microscopic* and *macroscopic* times is indicated by a dashed black arrow.

some *out* state at time $\tilde t_f$ in the future, which will be a superposition over all possible future states. We then evolve this superposition of states backwards again under the influence of a second source $J_-(x)$, returning to the same initial time $\tilde t_i$ and the original *in* state. The sources $J_\pm(x)$ are assumed to vanish adiabatically at the boundaries of the interval $[\tilde t_i,\ \tilde t_f]$. We may interpret the path of this evolution as defining a closed contour $\mathcal{C} \;=\; \mathcal{C}_+ \cup \mathcal{C}_-$ in the complex-time plane ($\mathfrak{t}$-plane, $\mathfrak{t} \;\in\; \mathbb{C}$), which is the union of two anti-parallel branches: $\mathcal{C}_+$, running from $\tilde t_i$ to $\tilde t_f \;-\; i\epsilon/2$; and $\mathcal{C}_-$, running from $\tilde t_f \;-\; i\epsilon/2$ back to $\tilde t_i \;-\; i\epsilon$. We refer to $\mathcal{C}_+$ and $\mathcal{C}_-$ as the positive time-ordered and negative anti-time-ordered branches, respectively. As depicted in Figure 1, the small imaginary part $\epsilon \;=\; 0^+$ has been added to allow us to distinguish the two, essentially coincident, branches. We parametrize the distance along the contour, starting from $\tilde t_i$, by the real variable $u \;\in\; [0,\ 1]$, which increases monotonically along $\mathcal{C}$. We may then define the contour by a path $\tilde z(u) \;=\; \tilde t(u) - i\tilde\tau(u) \;\in\; \mathfrak{t}$, where $\tilde t(0) \;=\; \tilde t(1) \;=\; \tilde t_i$ and $\tilde t(1/2) \;=\; \tilde t_f$. Thus, the complex CTP contour $\tilde z(u)$ may be written down explicitly as

$$\tilde z(u) \;=\; \theta(\tfrac{1}{2} - u)\,\Big[\,\tilde t_i + 2u\,\big(\,\tilde t_f - \tilde t_i\,\big)\Big] + \theta(u - \tfrac{1}{2})\,\Big[\,\tilde t_i + 2\,\big(\,1 - u\,\big)\,\big(\,\tilde t_f - \tilde t_i\,\big)\Big] - i\epsilon u\ , \tag{3.6}$$

with $\theta(0) \;=\; 1/2$, as given in (2.4).

In the presence of external sources $J_\pm(x)$, the eigenstate $|\Phi(\mathbf{x}), \tilde t; \tilde t_i\rangle_{\mathrm{H},\,J}$ of the Heisenberg field operator $\Phi_{\mathrm{H}}(\tilde t, \mathbf{x}; \tilde t_i)$ becomes explicitly time-dependent, satisfying the eigenvalue equation

$$\Phi_{\mathrm{H}}(\tilde t, \mathbf{x}; \tilde t_i)|\Phi(\mathbf{x}), \tilde t; \tilde t_i\rangle_{\mathrm{H},\,J} \;=\; \Phi(\mathbf{x})|\Phi(\mathbf{x}), \tilde t; \tilde t_i\rangle_{\mathrm{H},J}\ . \tag{3.7}$$

Also, $|\Phi(\mathbf{x}), \tilde t; \tilde t_i\rangle_{\mathrm{H},\,J}$ forms a complete orthonormal basis, respecting the orthonormality condition

$$\int [\mathrm{d}\Phi]\ |\Phi(\mathbf{x}), \tilde{t}; \tilde{t}_i\rangle_{\mathrm{H},\,J}\ {}_{\mathrm{H},\,J}\langle\Phi(\mathbf{x}), \tilde{t}; \tilde{t}_i|\ =\ \mathbf{I}_\infty\ . \tag{3.8}$$

We may then write the CTP generating functional $\mathcal{Z}[J_\pm, t]$ as

$$\begin{aligned} \mathcal{Z}[J_\pm, t]\ &=\ \int [\mathrm{d}\Phi]\ {}_{J_-}\langle 0_{\mathrm{in}}, \tilde{t}_i; \tilde{t}_i|\Phi(\mathbf{x}), \tilde{t}_f; \tilde{t}_i\rangle_{\mathrm{H},\,J_-}\ \times\ {}_{\mathrm{H},\,J_+}\langle\Phi(\mathbf{x}), \tilde{t}_f; \tilde{t}_i|0_{\mathrm{in}}, \tilde{t}_i; \tilde{t}_i\rangle_{J_+} \\ &=\ \int [\mathrm{d}\Phi]\ {}_{J_-}\langle 0_{\mathrm{in}}, \tilde{t}_i; \tilde{t}_i|\ \bar{\mathrm{T}}\left(e^{-i\int_{\tilde{t}_i}^{\tilde{t}_f}\mathrm{d}^4x\ \Phi_{\mathrm{H}}(x)J_-(x)}\right)|\Phi(\mathbf{x}), \tilde{t}_f; \tilde{t}_i\rangle_{\mathrm{H},\,J_-} \\ &\qquad \times\ {}_{\mathrm{H},\,J_+}\langle\Phi(\mathbf{x}), \tilde{t}_f; \tilde{t}_i|\ \mathrm{T}\left(e^{i\int_{\tilde{t}_i}^{\tilde{t}_f}\mathrm{d}^4x\ \Phi_{\mathrm{H}}(x)J_+(x)}\right)|0_{\mathrm{in}}, \tilde{t}_i; \tilde{t}_i\rangle_{J_+}\ , \end{aligned} \tag{3.9}$$

where the x^0 integrations run from $\tilde{t}_i$ to $\tilde{t}_f$ and the 'latest' time (with $u = 1$) appears furthest to the left.

In order to preserve the correspondence with the ordinary S-matrix theory in the asymptotic limit $\tilde{t}_i \to -\infty$, we take $\tilde{t}_f = -\tilde{t}_i$. With this identification, the CTP generating functional $\mathcal{Z}[J_\pm, t]$ becomes *manifestly CPT* invariant. This enables one to easily verify that microscopic CPT invariance continues to hold, even when time translational invariance is broken by thermal backgrounds, as we will see in Section 4. Given that $\tilde{t}_i$ is the microscopic time at which the three pictures of quantum mechanics coincide and the interactions are switched on, it is also the point at which the boundary conditions may be specified fully and *instantaneously* in terms of on-shell free particle states. The microscopic time $\tilde{t}_i$ is therefore the natural origin for a *macroscopic* time $t = \tilde{t}_f - \tilde{t}_i = 2\tilde{t}_f$. This *macroscopic* time is also the total interval of microscopic time over which the system has evolved. This fact is illustrated graphically in Figure 1.

We denote by $\Phi_\pm(x) \equiv \Phi(x^0 \in \mathcal{C}_\pm, \mathbf{x})$ fields with the microscopic time variable x^0 confined to the positive and negative branches of the contour, respectively. Following [39, 40], we define the doublets

$$\Phi^a(x)\ =\ \Big(\Phi_+(x)\,,\ \Phi_-(x)\Big)\ , \qquad \Phi_a(x)\ =\ \eta_{ab}\Phi^b(x)\ =\ \Big(\Phi_+(x)\,,\ -\Phi_-(x)\Big)\ , \tag{3.10}$$

where the CTP indices $a,\ b = 1,\ 2$ and $\eta_{ab} = \mathrm{diag}\,(1,\ -1)$ is an $\mathbb{SO}\,(1,\ 1)$ 'metric.' Inserting into (3.9) complete sets of eigenstates of the Heisenberg field operator at intermediate times, we may derive a path-integral representation of the CTP generating functional:

$$\mathcal{Z}[J_a, t]\ =\ \mathcal{N}\int[\mathrm{d}\Phi^a]\ \exp\left[i\left(S[\Phi^a, t]\ +\ \int_{\Omega_t}\mathrm{d}^4x\ J_a(x)\Phi^a(x)\right)\right]\ , \tag{3.11}$$

where $\mathcal{N}$ is some normalization and $\Omega_t \simeq [-t/2,\ t/2] \times \mathbb{R}^3$ is the Minkowski space-time volume bounded by the hypersurfaces $x^0 = \pm t/2$. In (3.11), the action is

$$\begin{aligned} S[\Phi^a, t]\ =\ \int_{\Omega_t}\mathrm{d}^4x\ \Big[\ &\tfrac{1}{2}\eta_{ab}\partial_\mu\Phi^a(x)\partial^\mu\Phi^b(x)\ -\ \tfrac{1}{2}\big(M^2\eta_{ab}\ -\ i\epsilon\mathbb{I}_{ab}\big)\Phi^a(x)\Phi^b(x) \\ &-\ \tfrac{1}{3!}g\eta_{abc}\Phi^a(x)\Phi^b(x)\Phi^c(x)\ -\ \tfrac{1}{4!}\lambda\eta_{abcd}\Phi^a(x)\Phi^b(x)\Phi^c(x)\Phi^d(x)\Big]\ , \end{aligned} \tag{3.12}$$

where $\eta_{abc\cdots} = +1$ for $a = b = c = \cdots = 1$, $\eta_{abc\cdots} = -1$ for $a = b = c = \cdots = 2$, and $\eta_{abc\cdots} = 0$ otherwise. In (3.12), the $\epsilon = 0^+$ gives the usual Feynman prescription, ensuring convergence of the CTP path integral. We note that the damping term is proportional to the identity matrix $\mathbb{I}_{ab}$ and not to the 'metric' η_{ab}. This prescription requires the addition of a contour-dependent damping term, proportional to $\varepsilon(\frac{1}{2} - u)$, which has the same sign on both the positive and negative branches of the contour, $\mathcal{C}_+$ and $\mathcal{C}_-$, respectively.

In order to define properly a path-ordering operator $\mathrm{T}_{\mathcal{C}}$, we introduce the contour-dependent step function

$$\theta_{\mathcal{C}}(x^0 - y^0) \;\equiv\; \theta(u_x - u_y) \,, \tag{3.13}$$

where $x^0 = \tilde{z}(u_x)$ and $y^0 = \tilde{z}(u_y)$. By analogy, we introduce a contour-dependent delta function

$$\delta_{\mathcal{C}}(x^0 - y^0) \;=\; \frac{\delta(u_x - u_y)}{\left|\frac{\mathrm{d}\tilde{z}}{\mathrm{d}u}\right|} \;=\; \frac{\delta(u_x - u_y)}{2|\tilde{t}_f - \tilde{t}_i|} \;=\; \frac{\delta(u_x - u_y)}{2t} \,. \tag{3.14}$$

As a consequence, a path-ordered propagator $\Delta_{\mathcal{C}}(x,y)$ may be defined as follows:

$$i\Delta_{\mathcal{C}}(x,y) \;\equiv\; \langle\, \mathrm{T}_{\mathcal{C}} \left[\, \Phi(x)\Phi(y) \,\right] \rangle \,. \tag{3.15}$$

For x^0 and y^0 on the positive branch $\mathcal{C}_+$, the path-ordering $\mathrm{T}_{\mathcal{C}}$ is equivalent to the standard time-ordering T and we obtain the time-ordered Feynman propagator $i\Delta_{\mathrm{F}}(x,y)$. On the other hand, for x^0 and y^0 on the negative branch $\mathcal{C}_-$, the path-ordering $\mathrm{T}_{\mathcal{C}}$ is equivalent to anti-time-ordering $\bar{\mathrm{T}}$ and we obtain the anti-time-ordered Dyson propagator $i\Delta_{\mathrm{D}}(x,y)$. For x^0 on $\mathcal{C}_+$ and y^0 on $\mathcal{C}_-$, x^0 is always 'earlier' than y^0 and we obtain the absolutely-ordered negative-frequency Wightman propagator $i\Delta_<(x,y)$. Conversely, for y^0 on $\mathcal{C}_+$ and x^0 on $\mathcal{C}_-$, we obtain the positive-frequency Wightman propagator $i\Delta_>(x,y)$. In the $\mathbb{SO}\,(1,\ 1)$ notation, we write the CTP propagator as the 2×2 matrix

$$i\Delta^{ab}(x,y) \;\equiv\; \langle\, \mathrm{T}_{\mathcal{C}} \left[\, \Phi^a(x)\Phi^b(y) \,\right] \rangle \;=\; i \begin{bmatrix} \Delta_{\mathrm{F}}(x,y) & \Delta_<(x,y) \\ \Delta_>(x,y) & \Delta_{\mathrm{D}}(x,y) \end{bmatrix} \,. \tag{3.16}$$

In this notation, the CTP indices a, b are raised and lowered by contraction with the 'metric' η_{ab}, e.g.

$$i\Delta_{ab}(x,y) \;=\; \eta_{ac}\, i\Delta^{cd}(x,y)\, \eta_{db} \;=\; i \begin{bmatrix} \Delta_{\mathrm{F}}(x,y) & -\Delta_<(x,y) \\ -\Delta_>(x,y) & \Delta_{\mathrm{D}}(x,y) \end{bmatrix} \,. \tag{3.17}$$

Notice the difference in sign of the off-diagonal elements in (3.17) compared with (3.16). An alternative definition $i\widetilde{\Delta}^{ab}(x,y)$ of the CTP propagator uses the so-called Keldysh basis [35] and is obtained by means of an orthogonal transformation:

$$\widetilde{\Delta}^{ab}(x,y) \;\equiv\; O^a{}_c\, O^b{}_d\, \Delta^{cd}(x,y) \;=\; \begin{bmatrix} 0 & \Delta_{\mathrm{A}}(x,y) \\ \Delta_{\mathrm{R}}(x,y) & \Delta_1(x,y) \end{bmatrix} \,, \qquad O^{ab} \;=\; \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \,. \tag{3.18}$$

In Section 4, we will generalize these results to macroscopic ensembles by incorporating background effects in terms of physical sources. In this case, the surface integral $\oint_{\partial\Omega_t} \mathrm{d}s_\mu\ \Phi^a(x)\partial^\mu\Phi^b(x)$ may not in general vanish on the boundary hypersurface $\partial\Omega_t$ of the volume Ω_t. However, by requiring the '+'- and '−'-type fields to satisfy

$$\Phi_+(x)\partial^\mu\Phi_+(x)\big|_{x^\mu \,\in\, \partial\Omega_t} \;=\; \Phi_-(x)\partial^\mu\Phi_-(x)\big|_{x^\mu \,\in\, \partial\Omega_t} \,, \tag{3.19}$$

we can ensure that surface terms cancel between the positive and negative branches, $\mathcal{C}_+$ and $\mathcal{C}_-$, respectively. In this case, the free part of the action may be rewritten as

$$S^0[\Phi^a, t] \;=\; \iint_{\Omega_t} \mathrm{d}^4x\, \mathrm{d}^4y\; \tfrac{1}{2}\Phi^a(x)\Delta^{0,\,-1}_{ab}(x,y)\Phi^b(y)\;, \tag{3.20}$$

where

$$\Delta^{0,\,-1}_{ab}(x,y) \;=\; \delta^{(4)}(x\,-\,y)\Big[-\big(\Box^2_x\,+\,M^2\big)\eta_{ab}\,+\,i\epsilon\mathbb{I}_{ab}\Big] \tag{3.21}$$

is the free inverse CTP propagator and $\Box^2_x \;\equiv\; \frac{\partial}{\partial x^\mu}\frac{\partial}{\partial x_\mu}$ is the d'Alembertian operator. Note that the variational principle remains well-defined irrespectively of (3.19), since we are always free to choose the variation of the field $\delta\Phi^a(x)$ to vanish for x^μ on $\partial\Omega_t$.

We may complete the square in the exponent of the CTP generating functional $\mathcal{Z}[J_a, t]$ in (3.11) by making the following shift in the field:

$$\Phi^a(x) \;\equiv\; \Phi'^a(x) \;-\; \int_{\Omega_t} \mathrm{d}^4y\; \Delta^{0,\,ab}(x,y)J_b(y)\;, \tag{3.22}$$

where $i\Delta^{0,\,ab}(x,y)$ is the free CTP propagator. We may then rewrite $\mathcal{Z}[J_a,t]$ in the form

$$\begin{aligned}\mathcal{Z}[J_a,t] \;&=\; \mathcal{Z}^0[0,t]\exp\bigg[i\int_{\Omega_t}\mathrm{d}^4x\;\mathcal{L}^{\mathrm{int}}\bigg(\frac{1}{i}\frac{\delta}{\delta J_a}\bigg)\bigg]\\ &\quad\times\;\exp\bigg[-\frac{i}{2}\iint_{\Omega_t}\mathrm{d}^4x\,\mathrm{d}^4y\;J_a(x)\Delta^{0,\,ab}(x,y)J_b(y)\bigg]\;,\end{aligned} \tag{3.23}$$

where $\mathcal{L}^{\mathrm{int}}$ is the interaction part of the Lagrangian and $\mathcal{Z}^0[0,t]$ is the free part of the generating functional

$$\mathcal{Z}^0[0,t] \;=\; \mathcal{N}\int[\mathrm{d}\Phi^a]\; e^{iS^0[\Phi^a,\,t]}\;, \tag{3.24}$$

with the free action $S^0[\Phi^a,t]$ given by (3.20). We may then express the resummed CTP propagator $i\Delta^{ab}(x,y,\tilde{t}_f;\tilde{t}_i)$ as follows:

$$i\Delta^{ab}(x,y,\tilde{t}_f;\tilde{t}_i) \;=\; \frac{1}{\mathcal{Z}[0,t]}\,\frac{1}{i}\,\frac{\delta}{\delta J_a(x)}\,\frac{1}{i}\,\frac{\delta}{\delta J_b(y)}\,\mathcal{Z}[J_a,t]\bigg|_{J_a\,=\,0}\;, \tag{3.25}$$

where $\mathcal{Z}[0,t]$ is the generating functional with the external sources J_a set to zero. The functional derivatives satisfy

$$\frac{\delta}{\delta J_a(x)}\int_{\Omega_t}\mathrm{d}^4y\;J^b(y) \;=\; \eta^{ab}\delta^{(4)}(x\,-\,y)\;, \tag{3.26}$$

with $x^\mu,\ y^\mu \;\in\; \Omega_t$. We will see in Section 4.3 that the resummed CTP propagator $\Delta^{ab}(x,y,\tilde{t}_f;\tilde{t}_i)$ is not in general time translationally invariant.

In the absence of interactions, eigenstates of the free Hamiltonian will propagate uninterrupted from times infinitely distant in the past to times infinitely far in the future. As such, we may extend the limits of integration in the free part of the action to positive and negative infinity, since

$$\big(\Box^2_x\,+\,M^2\big)\Phi^a(x)\big|_{x^0\,\notin\,[-t/2,\;t/2]} \;=\; 0\;, \tag{3.27}$$

i.e. the sources $J_a(x)$ vanish for $x^0 \notin [-t/2,\ t/2]$. The free CTP propagator $\Delta^{0,\,ab}(x,y)$ is then obtained by inverting (3.21) subject to the inverse relation

$$\int \mathrm{d}^4 z\ \Delta^{0,\,-1}_{ab}(x,z)\Delta^{0,\,bc}(z,y)\ =\ \eta_a{}^c\delta^{(4)}(x\,-\,y)\ , \tag{3.28}$$

where the domain of integration over z^0 is extended to infinity. We expect to recover the familiar propagators of the *in-out* formalism of asymptotic field theory, which occur in S-matrix elements and in the reduction formalism due to Lehmann, Symanzik and Zimmermann [61]. The propagators will also satisfy unitarity cutting rules [62–65], thereby maintaining perturbative unitarity of the theory. Specifically, the free Feynman (Dyson) propagators $i\Delta^0_{\mathrm{F(D)}}(x,y)$ satisfy the inhomogeneous Klein–Gordon equation

$$-\,\big(\Box^2_x\,+\,M^2\big)i\Delta^0_{\mathrm{F(D)}}(x,y)\ =\ (-)i\delta^{(4)}(x\,-\,y)\ ; \tag{3.29}$$

and the free Wightman propagators $i\Delta^0_{\gtrless}(x,y)$ satisfy the homogeneous equation

$$-\,\big(\Box^2_x\,+\,M^2\big)i\Delta^0_{>(<)}(x,y)\ =\ 0\ . \tag{3.30}$$

In the double momentum representation, the free part of the action (3.20) may be written as

$$S^0[\Phi^a]\ =\ \iint\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4p'}{(2\pi)^4}\,\frac{1}{2}\Phi^a(p)\Delta^{0,\,-1}_{ab}(p,p')\Phi^b(-p')\ , \tag{3.31}$$

where

$$\Delta^{0,\,-1}_{ab}(p,p')\ =\ \begin{bmatrix} p^2\,-\,m^2\,+\,i\epsilon & 0 \\ 0 & -\big(p^2\,-\,m^2\,-\,i\epsilon\big)\end{bmatrix}(2\pi)^4\delta^{(4)}(p\,-\,p') \tag{3.32}$$

is the double momentum representation of the free inverse CTP propagator, satisfying the inverse relation

$$\int\frac{\mathrm{d}^4q}{(2\pi)^4}\ \Delta^{0,\,-1}_{ab}(p,q)\Delta^{0,\,bc}(q,p')\ =\ \eta_a{}^c(2\pi)^4\delta^{(4)}(p\,-\,p')\ . \tag{3.33}$$

Given that the free inverse CTP propagator is proportional to a four-dimensional delta function of the two momenta, it may be written more conveniently in the *single* Fourier representation as

$$\Delta^{0,\,-1}_{ab}(p)\ =\ \begin{bmatrix} p^2\,-\,m^2\,+\,i\epsilon & 0 \\ 0 & -\big(p^2\,-\,m^2\,-\,i\epsilon\big)\end{bmatrix}\ . \tag{3.34}$$

Hence, for *translationally invariant* backgrounds, we may recast (3.32) in the form

$$\Delta^{0,\,-1}_{ab}(p)\Delta^{0,\,bc}(p)\ =\ \eta_a{}^c\ , \tag{3.35}$$

where

$$\Delta^{0,\,ab}(p)\ \equiv\ \int\frac{\mathrm{d}^4p'}{(2\pi)^4}\ \Delta^{0,\,ab}(p,p') \tag{3.36}$$

is the single-momentum representation of the free CTP propagator.

3.2 The Free CTP Propagator

We proceed now to make the following ansatz for the most general *translationally invariant* form of the free CTP propagator, without evaluating the correlation functions directly:

$$\Delta^{0,\,ab}(p) = \begin{bmatrix} \big(p^2 - M^2 + i\epsilon\big)^{-1} + \tilde{c}_1(p)\delta(p^2 - M^2) & \tilde{c}_3(p)\delta(p^2 - M^2) \\ \tilde{c}_2(p)\delta(p^2 - M^2) & -\big(p^2 - M^2 - i\epsilon\big)^{-1} + \tilde{c}_4(p)\delta(p^2 - M^2) \end{bmatrix} . \tag{3.37}$$

The $\tilde{c}_i(p) \equiv \theta(p_0)c_i(p) + \theta(-p_0)c_i'(p)$ are as yet undetermined analytic functions of the four-momentum p, which may in general be complex. The diagonal elements are the Fourier transforms of the most general translationally invariant solutions to the inhomogeneous Klein–Gordon equation (3.29), whilst the off-diagonal elements are the most general translationally invariant solutions to the homogeneous Klein–Gordon equation (3.30).

The remaining freedom in the matrix elements of $\Delta^{0,\,ab}(p)$ is determined by the following field-theoretic requirements:

(i) ***CPT* Invariance.** Since the action should be invariant under CPT, the real scalar field Φ should be even under CPT. From the definitions of the propagators in (A.1), we obtain the CPT relations in (A.3). Consequently, the momentum representation of these relations in (A.4) imply that

$$\tilde{c}_{1(4)}(p) \;=\; \tilde{c}_{1(4)}(-p)\,, \qquad \tilde{c}_2(p) \;=\; \tilde{c}_3(-p)\,. \tag{3.38}$$

(ii) **Hermiticity.** The Hermiticity properties of the correlation functions defined in (A.1) give rise to the Hermiticity relations outlined in (A.4). These imply that

$$\tilde{c}_4(p) \;=\; -\tilde{c}_1^*(p)\,, \qquad \tilde{c}_2(p) \;=\; -\tilde{c}_3^*(-p)\,. \tag{3.39}$$

In conjunction with (3.38), we conclude that $\tilde{c}_2(p)$ and $\tilde{c}_3(p)$ must be purely imaginary-valued functions of the four-momentum p.

(iii) **Causality.** The free Pauli–Jordan propagator $\Delta^0(x,y)$ is proportional only to the real part of the free Feynman propagator $\mathrm{Re}\,\Delta^0_{\mathrm{F}}(x,y)$ [cf. (A.6b)]. The addition of an even-parity on-shell dispersive part to the Fourier transform of the free Feynman propagator $\Delta^0_{\mathrm{F}}(p)$ will contribute to the free Pauli–Jordan propagator terms that are non-vanishing for space-like separations $(x - y)^2 < 0$, thus violating the micro-causality condition outlined in Section 2. It follows then that $\tilde{c}_1(p)$ and $\tilde{c}_4(p)$ are also purely imaginary-valued functions. We shall therefore replace the $\tilde{c}_i(p)$ by the real-valued functions $\tilde{f}_i(p)$ through $\tilde{c}_i(p) \equiv -2\pi i \tilde{f}_i(p)$, where the minus sign and factor of 2π have been included for later convenience. The explicit form of the free Pauli–Jordan propagator in (2.16b), along with the causality relation (2.24), give rise to the constraint

$$\tilde{f}_2(p) \;-\; \tilde{f}_3(p) \;=\; \varepsilon(p_0)\,. \tag{3.40}$$

(iv) **Unitarity.** Finally, the unitarity relations in (2.28) require that

$$\tilde{f}_2(p) \,+\, \tilde{f}_3(p) \;=\; 1 \,+\, \tilde{f}_1(p) \,+\, \tilde{f}_4(p)\,. \tag{3.41}$$

Solving the system of the above four constraints for $\tilde{f}_{1,2,3,4}(p)$, we arrive at the following expression for the most general translationally invariant free CTP propagator:

$$\begin{aligned}\Delta^{0,\,ab}(p) \;&=\; \begin{bmatrix} \big(p^2 \,-\, M^2 \,+\, i\epsilon\big)^{-1} & -2\pi i\,\theta(-p_0)\delta(p^2 \,-\, M^2) \\ -2\pi i\,\theta(p_0)\delta(p^2 \,-\, M^2) & -\big(p^2 \,-\, M^2 \,-\, i\epsilon\big)^{-1} \end{bmatrix} \\ &\quad -\, 2\pi i\tilde{f}(p)\delta(p^2 \,-\, M^2)\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}\,. \end{aligned} \tag{3.42}$$

All elements of $\Delta^{0,\,ab}(p)$ contain terms dependent upon the same function

$$\tilde{f}(p) \;\equiv\; \tilde{f}_1(p) \;=\; \theta(p_0)f(p) \,+\, \theta(-p_0)f(-p)\,. \tag{3.43}$$

These terms correspond to the vacuum expectation of the normal-ordered product of fields $\langle{:}\Phi(x)\Phi(y){:}\rangle$, which vanishes for the trivial vacuum $|0\rangle$. Therefore, we must conclude that $\tilde{f}(p)$ also vanishes in this case. We then obtain the free *vacuum* CTP propagator $\widehat{\Delta}^{0,\,ab}(p)$, which contains the set of propagators familiar from the unitarity cutting rules of absorptive part theory [62, 65]:

$$\widehat{\Delta}^{0,\,ab}(p) \;=\; \begin{bmatrix} \big(p^2 \,-\, M^2 \,+\, i\epsilon\big)^{-1} & -2\pi i\,\theta(-p_0)\delta(p^2 \,-\, M^2) \\ -2\pi i\,\theta(p_0)\delta(p^2 \,-\, M^2) & -\big(p^2 \,-\, M^2 \,-\, i\epsilon\big)^{-1} \end{bmatrix}\,. \tag{3.44}$$

We may similarly arrive at (3.42) by considering the free CTP propagator in the Keldysh representation $\widetilde{\Delta}^{0,\,ab}(p)$ from (3.18). The constraints outlined above allow us to add to the free Hadamard propagator $\Delta^0_1(p)$ any purely-imaginary even function of p proportional to $\delta(p^2 \,-\, M^2)$, that is

$$\begin{aligned}\widetilde{\Delta}^{0,\,ab}(p) \;&=\; \begin{bmatrix} 0 & \big[(p_0 \,-\, i\epsilon)^2 \,-\, \mathbf{p}^2 \,-\, M^2\big]^{-1} \\ \big[(p_0 \,+\, i\epsilon)^2 \,-\, \mathbf{p}^2 \,-\, M^2\big]^{-1} & -2\pi i\,\delta(p^2 \,-\, M^2) \end{bmatrix} \\ &\quad -\, 2\pi i\,2\tilde{f}(p)\delta(p^2 \,-\, M^2)\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}\,. \end{aligned} \tag{3.45}$$

We note that there is no such freedom to add terms to the free retarded and advanced propagators, $\Delta^0_{\mathrm{R}}(p)$ and $\Delta^0_{\mathrm{A}}(p)$, which is a consequence of the micro-causality constraints on the form of the free Pauli–Jordan propagator $\Delta^0(p)$. Employing the fact that

$$\mathbf{O}^{\mathsf{T}}\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}\mathbf{O} \;=\; \frac{1}{2}\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}\,, \tag{3.46}$$

we recover (3.42), which serves as a self-consistency check for the correctness of our ansatz for the free CTP propagator.

3.3 The Resummed CTP Propagator

In order to obtain the *resummed* CTP propagator, we must invert the inverse resummed CTP propagator on the restricted domain $[-t/2,\ t/2]$ subject to the inverse relation

$$\int_{\Omega_t} \mathrm{d}^4 z\ \Delta^{-1}_{ab}(x, z, \tilde{t}_f; \tilde{t}_i) \Delta^{bc}(z, y, \tilde{t}_f; \tilde{t}_i)\ =\ \eta_a{}^c \delta^{(4)}(x\ -\ y)\ , \tag{3.47}$$

for $x^\mu,\ y^\mu \in \Omega_t$. We shall see in Section 4.1 that this restriction of the time domain implies that a closed analytic form for the resummed CTP propagator $\Delta^{ab}(x, y, \tilde{t}_f; \tilde{t}_i)$ is in general not possible, for systems out of thermal equilibrium (see also our discussion in Section 5).

The double momentum representation of the inverse relation (3.47) takes on the form

$$\iint \frac{\mathrm{d}^4 q}{(2\pi)^4} \frac{\mathrm{d}^4 q'}{(2\pi)^4}\ \Delta^{-1}_{ab}(p, q, \tilde{t}_f; \tilde{t}_i)(2\pi)^4 \delta^{(4)}_t(q\ -\ q')\Delta^{bc}(q', p', \tilde{t}_f; \tilde{t}_i)\ =\ \eta_a{}^c (2\pi)^4 \delta^{(4)}_t(p\ -\ p')\ , \tag{3.48}$$

where we have defined

$$\delta^{(4)}_t(p - p')\ \equiv\ \delta_t(p_0 - p'_0)\delta^{(3)}(\mathbf{p} - \mathbf{p}')\ =\ \frac{1}{(2\pi)^4} \iint_{\Omega_t} \mathrm{d}^4 x\, \mathrm{d}^4 y\ e^{ip\cdot x} e^{-ip'\cdot y}\ \delta^{(4)}(x - y)\ . \tag{3.49}$$

The restriction of the domain of time integration has led to the introduction of the analytic weight function

$$\delta_t(p_0\ -\ p'_0)\ =\ \frac{t}{2\pi}\, \mathrm{sinc}\Big[\Big(\frac{p_0\ -\ p'_0}{2}\Big)t\Big]\ , \tag{3.50}$$

which has replaced the ordinary energy-conserving delta function. As expected, we have

$$\lim_{t\,\to\,\infty} \delta_t(p_0\ -\ p'_0)\ =\ \delta(p_0\ -\ p'_0)\ , \tag{3.51}$$

so that the standard description of asymptotic quantum field theory is recovered in the limit $\tilde{t}_i\ =\ -t/2\ \to\ -\infty$. Moreover, the weight function δ_t satisfies the convolution

$$\int_{-\infty}^{+\infty} \mathrm{d}q_0\ \delta_t(p_0\ -\ q_0)\delta_t(q_0\ -\ p'_0)\ =\ \delta_t(p_0\ -\ p'_0)\ . \tag{3.52}$$

The emergence of the function δ_t is a consequence of our requirement that the time evolution and the mapping between quantum-mechanical pictures (see Section 2) are governed by the standard interaction-picture evolution operator

$$U(\tilde{t}_f, \tilde{t}_i)\ =\ \mathrm{T} \exp\left(i \int_{\tilde{t}_i}^{\tilde{t}_f} \mathrm{d}\tilde{t}\, H^{\mathrm{int}}(\tilde{t}; \tilde{t}_i\,)\right)\ . \tag{3.53}$$

This evolution is defined for times greater than the boundary time $\tilde{t}_i$, at which point the three pictures are coincident. We stress here that δ_t is neither a prescription, nor is it an *a priori* regularization of the Dirac delta function.

As we will see later, the oscillatory behaviour of the sinc function in δ_t is fundamentally important to the dynamical behaviour of the system. Let us therefore convince ourselves that these oscillations persist, even if we smear the switching on of the interactions, or



Figure 2: Comparison of $\delta_t(p_0 - p_0')$ (black dotted) and $\delta_A(p_0 - p_0')$ (blue dashed). The arbitrary mass M is included so that axes are dimensionless.

equivalently, if we impose an adiabatic switching off of the interaction Hamiltonian for microscopic times outside the interval $[-t/2,\ t/2]$. To this end, we introduce to the interaction Hamiltonian H^{int} in (3.53) the Gaussian function

$$A_t(\tilde{t}) \;=\; \exp\left(-\,\frac{\tilde{t}^2}{2t^2}\right)\,, \tag{3.54}$$

such that the evolution operator takes the form

$$U(\tilde{t}_f, \tilde{t}_i) \;=\; \mathrm{T}\exp\left(i\int_{\tilde{t}_i}^{\tilde{t}_f} \mathrm{d}\tilde{t}\; A_t(\tilde{t}) H^{\text{int}}(\tilde{t}; \tilde{t}_i)\right)\,. \tag{3.55}$$

Clearly, for $\tilde{t} \gg t$, $A_t(\tilde{t}) \to 0$, whereas for $\tilde{t} \ll t$, $A_t(\tilde{t}) \to 1$. To account for the effect of $A_t(\tilde{t})$ in the action, the following replacement needs to be made:

$$\begin{aligned}\delta_t(p_0 - p_0') \;\to\; \delta_A(p_0 \,-\, p_0') \;&\equiv\; \frac{1}{2\pi}\int_{-t/2}^{+t/2} \mathrm{d}\tilde{t}\; e^{-i(p_0\,-\,p_0')\tilde{t}}\, A_t(\tilde{t}) \\ =\; \frac{t}{2\sqrt{2\pi}}\, e^{-\frac{1}{2}(p_0\,-\,p_0')^2t^2}&\left[\mathrm{Erf}\left(\frac{1\,-\,2i(p_0\,-\,p_0')t}{2\sqrt{2}}\right) + \mathrm{Erf}\left(\frac{1\,+\,2i(p_0\,-\,p_0')t}{2\sqrt{2}}\right)\right]\,. \end{aligned} \tag{3.56}$$

Due to the error functions of complex arguments in (3.56), the oscillatory behaviour remains. The analytic behaviour of both the functions $\delta_t(p_0 - p_0')$ and $\delta_A(p_0 - p_0')$ is shown in Figure 2 in which we see that the smooth smearing by the Gaussian function $A_t(\tilde{t})$ has little effect on the central region of the sinc function, as one would expect.

4 Non-Homogeneous Backgrounds

Until now, we have considered the vacuum to be an 'empty' state with all quantum numbers zero. In this section, we replace that 'empty' vacuum state with some macroscopic background, which may in general be inhomogeneous and incoherent. This non-trivial 'vacuum' is described by the density operator ρ. Following a derivation of the CTP Schwinger–Dyson equation, we show that it is not possible to obtain a closed analytic form for the resummed CTP propagators in the presence of time-dependent backgrounds. Finally, we generalize the discussions in Section 3.2 to obtain non-homogeneous free propagators in which space-time translational invariance is explicitly broken.

The density operator ρ is necessarily Hermitian and, for an isolated system, evolves in the interaction picture according to the von Neumann or quantum Liouville equation

$$\frac{\mathrm{d}\rho(\tilde{t};\tilde{t}_i)}{\mathrm{d}(\tilde{t}-\tilde{t}_i)} \;=\; -i\left[H^{\mathrm{int}}(\tilde{t};\tilde{t}_i),\;\rho(\tilde{t};\tilde{t}_i)\right]\,, \tag{4.1}$$

where $H^{\mathrm{int}}(\tilde{t};\tilde{t}_i)$ is the interaction part of the Hamiltonian in the interaction picture, which is time-dependent. Notice that the time derivative appearing on the LHS of (4.1) is taken with respect to the *time translationally invariant* quantity $\tilde{t}-\tilde{t}_i$. Developing the usual Neumann series, we find that

$$\rho(\tilde{t};\tilde{t}_i) \;=\; U^{-1}(\tilde{t},\tilde{t}_i)\,\rho(\tilde{t}_i;\tilde{t}_i)\,U(\tilde{t},\tilde{t}_i)\,, \tag{4.2}$$

where U is the evolution operator in (3.53). Hence, in the absence of external sources and given the unitarity of the evolution operator, the partition function $\mathcal{Z} = \mathrm{Tr}\,\rho$ is time-independent. On the other hand, the partition function of an open or closed subsystem is in general time-dependent due to the presence of external sources.

We are interested in evaluating time-dependent Ensemble Expectation Values (EEVs) of field operators $\langle\bullet\rangle_t$ at the *macro*scopic time t, which corresponds to the *micro*scopic time $\tilde{t}_f = t/2$, where the *bra-ket* now denotes the weighted expectation

$$\langle\bullet\rangle_t \;=\; \frac{\mathrm{Tr}\left(\,\rho(\tilde{t}_f;\tilde{t}_i)\,\bullet\,\right)}{\mathrm{Tr}\,\rho(\tilde{t}_f;\tilde{t}_i)}\,. \tag{4.3}$$

In this case, EEVs of two-point products of field operators begin with a total of nine independent coordinates: the microscopic time of the density operator and the two four-dimensional space-time coordinates of the field operators. As discussed in the beginning of Section 3 [cf. (3.2)], this number is reduced to the required seven coordinates, i.e. one temporal and six spatial, after setting all microscopic times equal to $\tilde{t}_f = t/2$. Hence, physical observables in the interaction picture are, for instance, of the form

$$\langle\Phi(\tilde{t}_f,\mathbf{x};\tilde{t}_i)\Phi(\tilde{t}_f,\mathbf{y};\tilde{t}_i)\rangle_t \;=\; \frac{\mathrm{Tr}\left(\,\rho(\tilde{t}_f;\tilde{t}_i)\Phi(\tilde{t}_f,\mathbf{x};\tilde{t}_i)\Phi(\tilde{t}_f,\mathbf{y};\tilde{t}_i)\,\right)}{\mathrm{Tr}\,\rho(\tilde{t}_f;\tilde{t}_i)}\,. \tag{4.4}$$

In the presence of a non-trivial background, the *out* state of Section 3 is replaced by the density operator ρ at the time $\tilde{t}_f = t/2$. Consequently, the starting point of the corresponding CTP generating functional of EEVs is

$$\mathcal{Z}[\rho,J_\pm,t] \;=\; \mathrm{Tr}\left[\left(\bar{\mathrm{T}}e^{-i\int_{\Omega_t}\mathrm{d}^4x\,J_-(x)\Phi_{\mathrm{H}}(x)}\right)\rho_{\mathrm{H}}(\tilde{t}_f;\tilde{t}_i)\left(\mathrm{T}e^{i\int_{\Omega_t}\mathrm{d}^4x\,J_+(x)\Phi_{\mathrm{H}}(x)}\right)\right]. \tag{4.5}$$

Within the generating functional $\mathcal{Z}$ in (4.5), the Heisenberg-picture density operator ρ_{H} has explicit time dependence, as it is built out of state vectors that depend on time due to the presence of the external sources $J_{\pm}$. In the absence of such sources, however, the state vectors do not evolve in time, so ρ_{H} and the partition function $\mathcal{Z}[\rho, J_{\pm} = 0, t] = \mathrm{Tr}\,\rho$ in (4.5) become time-independent quantities.

In contrast to existing approaches, the explicit microscopic time $\tilde{t}_f = t/2$ of the density operator $\rho_{\mathrm{H}}(\tilde{t}_f; \tilde{t}_i)$ appearing in the CTP generating functional (4.5) is the *time of observation* and not the initial time $\tilde{t}_i = -t/2$, at which the boundary conditions are specified. As we shall see in Section 6, this subtle difference will contribute to the absence of pinch singularities in the resulting perturbation series.

4.1 The Schwinger–Dyson Equation in the CTP Formalism

In order to generate a perturbation series of correlation functions in the presence of non-homogeneous backgrounds, we must derive the Schwinger–Dyson equation in the CTP formalism. Of particular interest is the explicit form of the Feynman–Dyson series for the expansion of the resummed CTP propagator. We will show that, in the time-dependent case, this series does not collapse to the resummation known from zero-temperature field theory. In particular, we find that a closed analytic form for the resummed CTP propagator is not attainable in general.

We proceed by inserting into the generating functional $\mathcal{Z}$ in (4.5) complete sets of eigenstates of the Heisenberg field operator Φ_{H} at intermediate times. In this way, we obtain the path-integral representation

$$\begin{aligned}\mathcal{Z}[\rho, J_a, t] \;=\; &\int [\mathrm{d}\Phi^a]\ {}_{\mathrm{H}}\langle \Phi_-(\mathbf{x}), \tilde{t}_f; \tilde{t}_i |\, \rho_{\mathrm{H}}(\tilde{t}_f; \tilde{t}_i)\, | \Phi_+(\mathbf{x}), \tilde{t}_f; \tilde{t}_i \rangle_{\mathrm{H}} \\ &\times\ \exp\bigg[i\bigg(S[\Phi^a, t] \;+\; \int_{\Omega_t} \mathrm{d}^4x\ J_a(x)\Phi^a(x) \bigg)\bigg]\;. \end{aligned} \tag{4.6}$$

As before, we may extend the limits of integration to infinity in the free part of the action and the J-dependent term, due to the fact that the external sources vanish outside the time interval $[-t/2,\ t/2]$. *It is only in the interaction part of the action that the finite domain of integration must remain.*

Following [40], we write the kernel ${}_{\mathrm{H}}\langle \Phi_-(\mathbf{x}), \tilde{t}_f; \tilde{t}_i |\, \rho_{\mathrm{H}}(\tilde{t}_f; \tilde{t}_i)\, | \Phi_+(\mathbf{x}), \tilde{t}_f; \tilde{t}_i \rangle_{\mathrm{H}}$ as an infinite series of poly-local sources:

$${}_{\mathrm{H}}\langle \Phi_-(\mathbf{x}), \tilde{t}_f; \tilde{t}_i |\, \rho_{\mathrm{H}}(\tilde{t}_f; \tilde{t}_i)\, | \Phi_+(\mathbf{x}), \tilde{t}_f; \tilde{t}_i \rangle_{\mathrm{H}} \;=\; \exp\big(iK[\Phi^a, t\,]\big)\;, \tag{4.7}$$

where

$$K[\Phi^a, t\,] \;=\; K + \int_{\Omega_t} \mathrm{d}^4x\ K_a(x, \tilde{t}_f; \tilde{t}_i)\Phi^a(x) + \frac{1}{2}\iint_{\Omega_t} \mathrm{d}^4x\, \mathrm{d}^4x'\ K_{ab}(x, x', \tilde{t}_f; \tilde{t}_i)\Phi^a(x)\Phi^b(x') + \cdots \tag{4.8}$$

is a time translationally invariant quantity. The poly-local sources $K_{ab\cdots}$ encode the state of the system at the microscopic time of observation $\tilde{t}_f = t/2$, i.e. the time at which the EEV is evaluated, according to Figure 1. It follows that these sources must contribute only for $x_0 = x'_0 = \cdots = \tilde{t}_f$ and therefore be proportional to delta functions of the form $\delta(x_0 - \tilde{t}_f)\delta(x'_0 - \tilde{t}_f)\cdots$. For instance, the bi-local source K_{ab} in the double momentum representation must have the form

$$K_{ab}(x, x', \tilde{t}_f; \tilde{t}_i) \; = \; \iint \frac{\mathrm{d}^4 p}{(2\pi)^4} \, \frac{\mathrm{d}^4 p'}{(2\pi)^4} \; e^{-ip\cdot x} \, e^{ip'\cdot x'} \, e^{i(p_0 \, - \, p'_0)\tilde{t}_f} K_{ab}(\mathbf{p}, \mathbf{p}', t) \; , \tag{4.9}$$

so that the p_0 and p'_0 integrations yield the required delta functions. Here, it is understood that the bi-local K_{ab} sources occuring on the LHS and RHS of (4.9) are distinguished by the form of their arguments. We emphasise that $K_{ab}(x, x', \tilde{t}_f; \tilde{t}_i)$ is not a time translationally invariant quantity due to the explicit dependence upon $\tilde{t}_f$ on the RHS of (4.9). In contrast, $K_{ab}(\mathbf{p}, \mathbf{p}', t)$ *is* time translationally invariant.

Notice that we could extend the limits of integration to infinity for the time integrals in the expansion of the kernel given in (4.8) also. Nevertheless, for the following derivation, all space-time integrals are taken to run over the hypervolume Ω_t for consistency. We should reiterate here that the limits of time integration can be extended to an infinite domain Ω_∞ in all but the interaction part of the action. We will also suppress the time-dependencies of the action S and sources $K_{ab\cdots}$ for notational convenience.

We now absorb the constant K in (4.8) into the overall normalization of the CTP generating functional $\mathcal{Z}$ and the local source K_a into a redefinition of the external source J_a. Then, $\mathcal{Z}$ may be written down as

$$\begin{aligned} \mathcal{Z}[J_a, K_{ab}, \cdots] \; &= \; \int [\mathrm{d}\Phi^a] \; \exp\bigg[i \bigg(S[\Phi^a] \; + \int \mathrm{d}^4 x \; J_a(x) \Phi^a(x) \\ &\quad + \; \frac{1}{2} \iint \mathrm{d}^4 x \, \mathrm{d}^4 y \; K_{ab}(x, x') \Phi^a(x) \Phi^b(x') \\ &\quad + \; \frac{1}{6} \iiint \mathrm{d}^4 x \, \mathrm{d}^4 x' \, \mathrm{d}^4 x'' \; K_{abc}(x, x', x'') \Phi^a(x) \Phi^b(x') \Phi^c(x'') \; + \; \cdots \bigg) \bigg] \; . \end{aligned} \tag{4.10}$$

The Cornwall–Jackiw–Tomboulis (CJT) effective action [37] is given by the following Legendre transform:

$$\begin{aligned} \Gamma[\widehat{\Phi}^a, \mathcal{G}^{ab}, \mathcal{G}^{abc}, \cdots] \; &= \; \mathcal{W}[J_a, K_{ab}, K_{abc}] \; - \int \mathrm{d}^4 x \; J_a(x) \widehat{\Phi}^a(x) \\ &\quad - \; \frac{1}{2} \iint \mathrm{d}^4 x \, \mathrm{d}^4 x' \; K_{ab}(x, x') \left(\widehat{\Phi}^a(x) \widehat{\Phi}^b(x') \; + \; i\hbar \mathcal{G}^{ab}(x, x') \right) \\ &\quad - \; \frac{1}{6} \iiint \mathrm{d}^4 x \, \mathrm{d}^4 x' \, \mathrm{d}^4 x'' \; K_{abc}(x, x', x'') \left(\widehat{\Phi}^a(x) \widehat{\Phi}^b(x') \widehat{\Phi}^c(x'') \right. \\ &\quad \left. + \; 3i\hbar \mathcal{G}^{(ab}(x, x') \widehat{\Phi}^{c)}(x'') \; - \; \hbar^2 \mathcal{G}^{abc}(x, x', x'') \right) \; + \; \cdots \; , \end{aligned} \tag{4.11}$$

where $\mathcal{W}[J_a, K_{ab}, K_{abc}, \cdots] \; = \; -i\hbar \ln \mathcal{Z}[J_a, K_{ab}, K_{abc}, \cdots]$ is the generating functional of connected ensemble Green's functions. We obtain an infinite system of equations:

$$\widehat{\Phi}^a(x) \; = \; \frac{\delta \mathcal{W}}{\delta J_a(x)} \; = \; \langle \Phi^a(x) \rangle \; , \tag{4.12a}$$

$$\begin{aligned} i\hbar \mathcal{G}^{ab}(x, x') \; &= \; 2 \frac{\delta \mathcal{W}}{\delta K_{ab}(x, x')} \; - \; \widehat{\Phi}^a(x) \widehat{\Phi}^b(x') \\ &= \; -i\hbar \frac{\delta^2 \mathcal{W}}{\delta J_a(x) \delta J_b(x')} \; = \; \langle \, \mathrm{T}_{\mathcal{C}} \left[\Phi^a(x) \Phi^b(x') \right] \rangle \; - \; \langle \Phi^a(x) \rangle \langle \Phi^b(x') \rangle \; , \end{aligned} \tag{4.12b}$$

$$
\begin{aligned}
-\hbar^2 \mathcal{G}^{abc}(x,x',x'') \ &= \ 6\frac{\delta \mathcal{W}}{\delta K_{abc}(x,x',x'')} \ - \ 3i\hbar \mathcal{G}^{(ab}(x,x')\widehat{\Phi}^{c)}(x'') \ - \ \widehat{\Phi}^a(x)\widehat{\Phi}^b(x')\widehat{\Phi}^c(x'') \\
&= \ -\hbar^2 \frac{\delta^3 \mathcal{W}}{\delta J_a(x)\delta J_b(x')\delta J_c(x'')} \ , \qquad (4.12\mathrm{c})
\end{aligned}
$$

and

$$
\begin{aligned}
\frac{\delta \Gamma}{\delta \widehat{\Phi}^a(x)} \ &= \ -J_a(x) \ - \ \int \mathrm{d}^4x' \ K_{ab}(x,x')\widehat{\Phi}^b(x') \qquad (4.13\mathrm{a}) \\
& \quad - \ \frac{1}{2}\iint \mathrm{d}^4x' \, \mathrm{d}^4x'' \ K_{abc}(x,x',x'') \left(\widehat{\Phi}^b(x')\widehat{\Phi}^c(x'') \ + \ i\hbar \mathcal{G}^{bc}(x',x'') \right) \ - \ \cdots \ , \\
\frac{\delta \Gamma}{\delta \mathcal{G}^{ab}(x,x')} \ &= \ -\frac{i\hbar}{2} K_{ab}(x,x') \ - \ \frac{i\hbar}{2}\int \mathrm{d}^4x'' \ K_{abc}(x,x',x'')\widehat{\Phi}^c(x'') \ - \ \cdots \ , \qquad (4.13\mathrm{b}) \\
\frac{\delta \Gamma}{\delta \mathcal{G}^{abc}(x,x',x'')} \ &= \ -\frac{i\hbar}{6} K_{abc}(x,x',x'') \ - \ \cdots \ , \qquad (4.13\mathrm{c})
\end{aligned}
$$

where the parentheses (abc) on the RHS of (4.12c) denote cyclic permutation with respect to the indices a, b, c.

The above infinite system of equations (4.12) and (4.13) may be simplified by assuming that the density operator ρ is Gaussian, as we will do later in Section 4.3. In this case, the tri-local and higher kernels (K_{abc}, K_{abcd}, $\cdots$) can be set to zero in (4.13). We may then eliminate the three- and higher-point connected Green's functions ($\mathcal{G}^{abc}$, $\mathcal{G}^{abcd}$, $\cdots$) as dynamical variables by performing a second Legendre transform

$$
\Gamma[\widehat{\Phi}^a, \mathcal{G}^{ab}] \ \equiv \ \Gamma[\widehat{\Phi}^a, \mathcal{G}^{ab}, \widetilde{\mathcal{G}}^{abc}, \cdots] \ , \qquad (4.14)
$$

where the $\widetilde{\mathcal{G}}$'s are functionals of $\widehat{\Phi}^a$ and $\mathcal{G}^{ab}$ given by

$$
\frac{\delta \Gamma}{\delta \mathcal{G}^{abc\cdots}}[\widehat{\Phi}^a, \mathcal{G}^{ab}, \widetilde{\mathcal{G}}^{abc}, \cdots] \ = \ 0 \ . \qquad (4.15)
$$

The effective action is evaluated by expanding around the constant background field $\Phi^a_0(x) \ = \ \Phi^a(x) \ - \ \hbar^{1/2}\phi^a(x)$, defined at the saddle-point

$$
\frac{\delta S[\Phi^a]}{\delta \Phi^a(x)}\bigg|_{\Phi \, = \, \Phi_0} \ + \ J_a(x) \ + \ \int \mathrm{d}^4x' \ K_{ab}(x,x')\Phi^b_0(x') \ = \ 0 \ . \qquad (4.16)
$$

The result of this expansion is well known [37, 38] and, truncating to order $\hbar^2$, we obtain the two-particle-irreducible (2PI) CJT effective action

$$
\Gamma[\widehat{\Phi}^a, \mathcal{G}^{ab}] \ = \ S[\widehat{\Phi}^a] + \frac{i\hbar}{2}\,\mathrm{Tr}_x \Big[\mathrm{Ln}_x \, \mathrm{Det}_{ab} \, \mathcal{G}^{-1}_{ab} + \Big(G^{-1}_{ab} - K_{ab} \Big) * \mathcal{G}^{ab} - \eta_a^{\ a} \Big] + \hbar^2 \Gamma_2[\widehat{\Phi}^a, \mathcal{G}^{ab}] \ , \qquad (4.17)
$$

where a subscript x and the $*$'s indicate that the trace, logarithm and products should be understood as functional operations. The operator G^{-1}_{ab} is defined by

$$
\begin{aligned}
G^{-1}_{ab}(\widehat{\Phi}^a; x, x') \ &= \ \frac{\delta^2 S[\widehat{\Phi}^a]}{\delta \widehat{\Phi}^a(x)\delta \widehat{\Phi}^b(x')} \ + \ K_{ab}(x,x') \\
&= \ \Delta^{0,\,-1}_{ab}(x,x') \ + \ \frac{\delta^2 S^{\mathrm{int}}[\widehat{\Phi}^a]}{\delta \widehat{\Phi}^a(x)\delta \widehat{\Phi}^b(x')} \ + \ K_{ab}(x,x') \ , \qquad (4.18)
\end{aligned}
$$

where $\Delta^{0,-1}_{ab}(x,x')$ is the free inverse CTP propagator in (3.21) and $S^{\rm int}[\widehat{\Phi}^a]$ is the interaction part of the action. Obviously, all Green's functions depend upon the state of the system at the macroscopic time t through the bi-local source K_{ab}. For the Lagrangian in (2.1), we have

$$\begin{aligned} G^{-1}_{ab}(\widehat{\Phi}^a;x,x') \ &= \ \delta^{(4)}(x\ -\ x')\Big[-\ (\Box^2_x\ +\ M^2)\eta_{ab}\ +\ i\epsilon\mathbb{I}_{ab} \\ &\qquad -\ g\eta_{abc}\widehat{\Phi}^c(x)\ -\ \frac{1}{2}\lambda\eta_{abcd}\widehat{\Phi}^c(x)\widehat{\Phi}^d(x)\Big]\ +\ K_{ab}(x,x')\ , \end{aligned} \tag{4.19}$$

where $\eta_{abc\cdots}\ =\ +1$ for all indices $a\ =\ b\ =\ \cdots\ =\ 1$, $\eta_{abc\cdots}\ =\ -1$ for all indices $a\ =\ b\ =\ \cdots\ =\ 2$ and $\eta_{abc\cdots}\ =\ 0$ otherwise.

The overall normalization ($\eta_a^{\ a}$) of (4.17) has been chosen so that when $K_{ab}(x,x')\ =\ 0$, we may recover the conventional effective action [66] by making a further Legendre transform to eliminate $\mathcal{G}^{ab}$ as a dynamical variable:

$$\Gamma[\widehat{\Phi}^a]\ \equiv\ \Gamma[\widehat{\Phi}^a,\widetilde{\mathcal{G}}^{ab}]\ =\ S[\widehat{\Phi}^a]\ +\ \frac{i\hbar}{2}{\rm Tr}_x\,{\rm Ln}_x\,{\rm Det}_{ab}\,G^{-1}_{ab}\ +\ \mathcal{O}(\hbar^2) \tag{4.20}$$

Here, $\mathcal{G}^{-1}_{ab}$ has been replaced by G^{-1}_{ab} and $\widetilde{\mathcal{G}}^{ab}$ is a functional of $\widehat{\Phi}^a$.

In the CJT effective action (4.17), $\Gamma_2[\widehat{\Phi}^a,\mathcal{G}^{ab}]$ is the sum of all 2PI vacuum graphs:

$$\Gamma_2[\widehat{\Phi}^a,\mathcal{G}^{ab}]\ =\ -i\sum_{a,\,b\,=\,1,\,2}\Bigg[\frac{1}{8}$$



$$\delta_{ab}\ +\ \frac{1}{12}$$



$$\Bigg]\ , \tag{4.21}$$

where combinatorial factors have been written explicitly and we associate with each n-point vertex a factor of

$$iS^{(n)}_a(\widehat{\Phi}^a;x)\ =\ i\frac{\delta^n S[\widehat{\Phi}^a]}{\delta\big(\widehat{\Phi}^a(x)\big)^n}\ , \tag{4.22}$$

and each line a factor of $i\mathcal{G}^{ab}(\widehat{\Phi}^a;x,y)$. The three- and four-point vertices are

$$iS^{(3)}_a(\widehat{\Phi}^a;x)\ =\ -ig\eta_{aaa}\ -\ i\lambda\eta_{aaaa}\widehat{\Phi}^a(x)\ ,\qquad iS^{(4)}_a(\widehat{\Phi}^a;x)\ =\ -i\lambda\eta_{aaaa}\ . \tag{4.23}$$

Upon functional differentiation of the CJT effective action (4.17) with respect to $\mathcal{G}^{ab}(x,y)$, we obtain by virtue of (4.13b) the Schwinger–Dyson equation

$$\mathcal{G}^{-1}_{ab}(\widehat{\Phi}^a;x,y)\ =\ G^{-1}_{ab}(\widehat{\Phi}^a;x,y)\ +\ \Pi_{ab}(\widehat{\Phi}^a,\mathcal{G}^{ab};x,y)\ , \tag{4.24}$$

where

$$\Pi_{ab}(\widehat{\Phi}^a,\mathcal{G}^{ab};x,y)\ =\ -2i\hbar\frac{\delta\Gamma_2[\widehat{\Phi}^a,\mathcal{G}^{ab}]}{\delta\mathcal{G}^{ab}(x,y)}$$

$$=\ -i\hbar\Bigg[$$



$$\delta_{ab}\ +$$



$$\Bigg] \tag{4.25}$$

is the one-loop truncated CTP self-energy. A combinatorial factor of $\frac{1}{2}$ has been absorbed into the diagrammatics.

Suppressing the $\widehat{\Phi}^a$ and $\mathcal{G}^{ab}$ arguments for notational convenience, the CTP self-energy $\Pi_{ab}(x,y)$ may be written in matrix form as

$$\Pi_{ab}(x,y) \; = \; \begin{bmatrix} \Pi(x,y) & -\Pi_<(x,y) \\ -\Pi_>(x,y) & -\Pi^*(x,y) \end{bmatrix} , \tag{4.26}$$

where $\Pi(x,y)$ and $-\Pi^*(x,y)$ are the time- and anti-time-ordered self-energies; and $\Pi_>(x,y)$ and $\Pi_<(x,y)$ are the positive- and negative-frequency absolutely-ordered self-energies, respectively. In analogy to the propagator definitions discussed in Section 2 and Appendix A, we also define the self-energy functions

$$\Pi_1(x,y) \; = \; \Pi_>(x,y) \, + \, \Pi_<(x,y) \; = \; \Pi(x,y) \, - \, \Pi^*(x,y) \; = \; 2i\,\mathrm{Im}\,\Pi(x,y) \; , \tag{4.27a}$$

$$\Pi_{\mathcal{P}}(x,y) \; = \; \frac{1}{2}\Big(\Pi_{\mathrm{R}}(x,y) \, + \, \Pi_{\mathrm{A}}(x,y) \Big) \; = \; \mathrm{Re}\,\Pi(x,y) \; , \tag{4.27b}$$

$$2iM\Gamma(x,y) \; = \; \Pi_>(x,y) \, - \, \Pi_<(x,y) \; = \; \Pi_{\mathrm{R}}(x,y) \, - \, \Pi_{\mathrm{A}}(x,y) \; , \tag{4.27c}$$

which satisfy relations analogous to those described in Appendix A. $\Gamma(x,y)$ in (4.27c) is related to the usual Breit–Wigner width in the equilibrium and zero-temperature limits. The Keldysh representation [see (3.18)] $\widetilde{\Pi}_{ab}(x,y)$ of the CTP self-energy reads:

$$\widetilde{\Pi}_{ab}(x,y) \; = \; \begin{bmatrix} \Pi_1(x,y) & \Pi_{\mathrm{R}}(x,y) \\ \Pi_{\mathrm{A}}(x,y) & 0 \end{bmatrix} . \tag{4.28}$$

In the limit $\widehat{\Phi}^a(x) \; \to \; 0$, the Schwinger–Dyson equation (4.24) reduces to

$$\Delta^{-1}_{ab}(x,y,\tilde{t}_f;\tilde{t}_i) \; = \; \Delta^{0,\,-1}_{ab}(x,y) \, + \, K_{ab}(x,y,\tilde{t}_f;\tilde{t}_i) \, + \, \Pi_{ab}(x,y,\tilde{t}_f;\tilde{t}_i) \; , \tag{4.29}$$

in which $\Delta^{-1}_{ab}(x,y,\tilde{t}_f;\tilde{t}_i) \equiv \mathcal{G}^{-1}_{ab}(\widehat{\Phi}^a = 0; x,y,\tilde{t}_f;\tilde{t}_i)$ and $\Delta^{0,\,-1}_{ab}(x,y)$ is the free inverse CTP propagator defined in (3.21). We have re-introduced the dependence upon $\tilde{t}_f$ and $\tilde{t}_i$ for clarity. Notice that due to the explicit $\tilde{t}_f$-dependence of the bi-local source $K_{ab}(x,y,\tilde{t}_f;\tilde{t}_i)$ in (4.9), the inverse resummed CTP propagator $\Delta^{-1}_{ab}(x,y,\tilde{t}_f;\tilde{t}_i)$ and the CTP self-energy $\Pi_{ab}(x,y,\tilde{t}_f;\tilde{t}_i)$ are not time translationally invariant quantities.

In order to develop a self-consistent inversion of the Schwinger–Dyson equation in (4.29), the bi-local source $K_{ab}(x,y,\tilde{t}_f;\tilde{t}_i)$ is absorbed into an inverse non-homogeneous CTP propagator

$$D^{0,\,-1}_{ab}(x,y,\tilde{t}_f;\tilde{t}_i) \; = \; \Delta^{0,\,-1}_{ab}(x,y) \, + \, K_{ab}(x,y,\tilde{t}_f;\tilde{t}_i), \tag{4.30}$$

whose inverse, to leading order in K_{ab}, is the free CTP propagator $\Delta^{0,\,ab}(x,y,\tilde{t}_f;\tilde{t}_i)$, i.e.

$$D^{0,\,ab}(x,y,\tilde{t}_f;\tilde{t}_i) \; \equiv \; \Delta^{0,\,ab}(x,y,\tilde{t}_f;\tilde{t}_i) \, + \, \mathcal{O}(K^2) \; , \tag{4.31}$$

as we will illustrate in Sections 4.3 and 5. The contribution of the bi-local source K_{ab} is now absorbed into the free CTP propagator $\Delta^{0,\,ab}(x,y,\tilde{t}_f;\tilde{t}_i)$, whose time translational invariance is broken as a result. The Schwinger–Dyson equation (4.29) may then be written in the double momentum representation as

$$\Delta^{-1}_{ab}(p,p',\tilde{t}_f;\tilde{t}_i) \; = \; \Delta^{0,\,-1}_{ab}(p,p') \, + \, \Pi_{ab}(p,p',\tilde{t}_f;\tilde{t}_i) \; . \tag{4.32}$$

Since the stationary vacuum $|0\rangle$ has been replaced by the density operator ρ at the microscopic time $\tilde{t}_f = t/2$, we must use the following field-particle duality relation in the Wick contraction of interaction-picture fields:

$$\langle 0|\Phi(x;\tilde{t}_i)a^\dagger(\mathbf{k},\tilde{t}_f;\tilde{t}_i)|0\rangle \;=\; e^{-ik\cdot x}e^{iE(\mathbf{k})\tilde{t}_f}\ . \tag{4.33}$$

Here, the extra phase $e^{iE(\mathbf{k})\tilde{t}_f}$ arises from the fact that the creation and annihilation operators of the interaction-picture field $\Phi(x;\tilde{t}_i)$ are evaluated at the microscopic time $\tilde{t} = 0$ [cf. (2.12) and (2.18)], whereas the operator $a^\dagger(\mathbf{k},\tilde{t}_f;\tilde{t}_i)$, resulting from the expansion of the density operator ρ (see Section 4.3), is evaluated at the microscopic time $\tilde{t}_f = t/2$. Analytically continuing this extra phase to off-shell energies, we must therefore associate with each external vertex of the self-energy $\Pi_{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$, appearing in (4.32), a phase:

$$e^{ip_0\tilde{t}_f}\ , \tag{4.34}$$

where p_0 is the energy flowing *into* the vertex. This amounts to the absorption of an overall phase

$$e^{i(p_0\,-\,p_0')\tilde{t}_f} \tag{4.35}$$

into the definition of the self-energy $\Pi_{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$. We recall from (4.9) that the bi-local source $K_{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ already contains this *free phase* factor.

Convoluting from left and right on both sides of (4.32) first with the weight function $(2\pi)^4\delta_t^{(4)}(p\,-\,p')$ from (3.50) and then with $\Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ and $\Delta^{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$, respectively, we obtain the Feynman–Dyson series

$$\begin{aligned}\Delta^{ab}(p,p',\tilde{t}_f;\tilde{t}_i) \;&=\; \Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i) \;-\; \int\cdots\int\frac{\mathrm{d}^4q}{(2\pi)^4}\,\frac{\mathrm{d}^4q'}{(2\pi)^4}\,\frac{\mathrm{d}^4q''}{(2\pi)^4}\,\frac{\mathrm{d}^4q'''}{(2\pi)^4}\\ &\times\;\Delta^{0,\,ac}(p,q,\tilde{t}_f;\tilde{t}_i)(2\pi)^4\delta_t^{(4)}(q\,-\,q')\Pi_{cd}(q',q'',\tilde{t}_f;\tilde{t}_i)(2\pi)^4\delta_t^{(4)}(q''\,-\,q''')\Delta^{db}(q''',p',\tilde{t}_f;\tilde{t}_i)\ ,\end{aligned} \tag{4.36}$$

where $\delta_t^{(4)}(p\,-\,p')$ is defined in (3.49). Because of the form of $\delta_t(p_0\,-\,p_0')$ in (3.50), we see that this series does not collapse to an algebraic equation of resummation, as known from zero-temperature field theory. As we will see in Section 4.2, one cannot write down a closed analytic form for the resummed CTP propagator $\Delta^{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$.

Given that δ_t satisfies the convolution in (3.52), the weight functions may be absorbed into the external vertices of the self-energy $\Pi_{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$; see Section 9. The Feynman–Dyson series may then be written in the more concise form

$$\begin{aligned}\Delta^{ab}(p,p',\tilde{t}_f;\tilde{t}_i) \;&=\; \Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i)\\ &-\;\iint\frac{\mathrm{d}^4q}{(2\pi)^4}\,\frac{\mathrm{d}^4q'}{(2\pi)^4}\;\Delta^{0,\,ac}(p,q,\tilde{t}_f;\tilde{t}_i)\Pi_{cd}(q,q',\tilde{t}_f;\tilde{t}_i)\Delta^{db}(q',p',\tilde{t}_f;\tilde{t}_i)\ .\end{aligned} \tag{4.37}$$

Note that for finite t, $\delta_t(p_0\,-\,p_0')$ is analytic in the limit $p_0\,\to\,p_0'$. As we shall see in Section 6, the systematic incorporation of these finite-time effects ensures that the perturbation expansion is free of pinch singularities.

4.2 Applicability of the Gradient Expansion

Here, we will look more closely at the inverse relation (3.47) that determines the resummed CTP propagator. We will show that a full matrix inversion may only be performed in the thermodynamic equilibrium limit. Hence, the application of truncated gradient expansions and the use of partially resummed quasi-particle propagators, particularly for early times, become questionable in out-of-equilibrium systems.

We define the relative and central coordinates

$$R^{\mu}_{xy} \;=\; x^{\mu} \;-\; y^{\mu}\;, \qquad X^{\mu}_{xy} \;=\; \frac{x^{\mu} \;+\; y^{\mu}}{2}\;, \tag{4.38}$$

such that

$$x^{\mu} \;=\; X^{\mu}_{xy} \;+\; \frac{1}{2}R^{\mu}_{xy}\;, \qquad y^{\mu} \;=\; X^{\mu}_{xy} \;-\; \frac{1}{2}R^{\mu}_{xy}\;. \tag{4.39}$$

We then introduce the Wigner transform (see [51]), namely the Fourier transform with respect to the relative coordinate R^{μ}_{xy} only. Explicitly, the Wigner transform of a function $F(R, X)$ is

$$F(p, X) \;=\; \int \mathrm{d}^4R\; e^{ip\cdot R}\, F(R, X)\;. \tag{4.40}$$

The resummed CTP propagator $\Delta^{ab}(x, y)$ respects the inverse relation in (3.47). Here, we suppress the $\tilde{t}_f$- and $\tilde{t}_i$-dependence of the propagators for notational convenience. Inserting into (3.47), the Wigner transforms of the resummed and inverse resummed CTP propagators, the inverse relation takes the form

$$\begin{aligned} \int_{\Omega_t} \mathrm{d}^4z \iint \frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4p'}{(2\pi)^4}\, e^{-ip\cdot R_{xz}}\, e^{-ip'\cdot R_{zy}}\, \Delta^{-1}_{ab}(p, X_{xz})\Delta^{bc}(p', X_{zy}) \\ =\; \iint \frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4p'}{(2\pi)^4}\, e^{-ip\cdot x}\, e^{ip'\cdot y}\, \eta_a^{\ c}(2\pi)^4\delta^{(4)}_t(p \;-\; p')\;, \end{aligned} \tag{4.41}$$

where the z^0 domain of integration is restricted to be in the range $[-t/2,\ t/2]$.

In the case where deviations from homogeneity are small, i.e. when the characteristic scale of macroscopic variations in the background is large in comparison to that of the microscopic single-particle excitations, we may perform a gradient expansion of the inverse relation in terms of the soft derivative $\partial^{\mu}_{X_{xy}} \;\equiv\; \partial/\partial X_{xy,\,\mu}$. Writing $X_{xz} \;=\; X_{xy} \;+\; R_{zy}/2$ and $X_{zy} \;=\; X_{xy} \;-\; R_{xz}/2$ and after integrating by parts, we obtain

$$\begin{aligned} \iint \frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4p'}{(2\pi)^4}\, e^{-ip\cdot x}\, e^{ip'\cdot y}\,(2\pi)^4\delta^{(4)}_t(p - p') \\ \times\; \Delta^{-1}_{ab}(p, X)\exp\Big[-\frac{i}{2}\big(\overleftarrow{\partial_p}\cdot\overrightarrow{\partial_X} \;-\; \overleftarrow{\partial_X}\cdot\overrightarrow{\partial_{p'}}\big)\Big]\Delta^{bc}(p', X) \\ =\; \iint \frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4p'}{(2\pi)^4}\, e^{-ip\cdot x}\, e^{ip'\cdot y}\, \eta_a^{\ c}(2\pi)^4\delta^{(4)}_t(p \;-\; p')\;, \end{aligned} \tag{4.42}$$

where $X \;\equiv\; X_{xy}$.

We now define the central and relative momenta

$$q^{\mu} \;=\; \frac{p^{\mu} \;+\; p'^{\mu}}{2}\;, \qquad Q^{\mu} \;=\; p^{\mu} \;-\; p'^{\mu}\;, \tag{4.43}$$

which are the Fourier conjugates to the relative and central coordinates, R^μ and X^μ, respectively. It follows that

$$p^\mu \;=\; q^\mu \,+\, \frac{1}{2}Q^\mu\;, \qquad p'^\mu \;=\; q^\mu \,-\, \frac{1}{2}Q^\mu\;. \tag{4.44}$$

We may then Fourier transform (4.42) with respect to R_{xy} to obtain

$$\begin{aligned}&\int\!\frac{\mathrm{d}^4Q}{(2\pi)^4}\;e^{-iQ\cdot X}\;(2\pi)^4\delta_t^{(4)}(Q)\\ &\qquad\times\;\exp\Big[-i\,\Big(\diamondsuit^-_{q,X}\,+\,2\diamondsuit^+_{Q,X}\Big)\Big]\{\Delta^{-1}_{ab}(q+Q/2,X)\}\{\Delta^{bc}(q-Q/2,X)\}\\ &\qquad\qquad\qquad\qquad\;=\;\int\!\frac{\mathrm{d}^4Q}{(2\pi)^4}\;e^{-iQ\cdot X}\;\eta_a^{\ c}(2\pi)^4\delta_t^{(4)}(Q)\;,\end{aligned} \tag{4.45}$$

where, following [42, 43], we have introduced the diamond operator

$$\diamondsuit^\pm_{p,X}\{A\}\{B\}\;=\;\frac{1}{2}\{A,\;B\}^\pm_{p,X} \tag{4.46}$$

and $\{A,\;B\}^\pm_{p,X}$ denote the symmetric and anti-symmetric Poisson brackets

$$\{A,\;B\}^\pm_{p,X}\;\equiv\;\frac{\partial A}{\partial p^\mu}\,\frac{\partial B}{\partial X_\mu}\;\pm\;\frac{\partial A}{\partial X^\mu}\,\frac{\partial B}{\partial p_\mu}\;. \tag{4.47}$$

For $t > 0$, we may perform the integral on the RHS of (4.45), yielding

$$\begin{aligned}&\int\!\frac{\mathrm{d}^4Q}{(2\pi)^4}\;e^{-iQ\cdot X}\;(2\pi)^4\delta_t^{(4)}(Q)\\ &\times\exp\Big[-i\,\Big(\diamondsuit^-_{q,X}\,+\,2\diamondsuit^+_{Q,X}\Big)\Big]\{\Delta^{-1}_{ab}(q+Q/2,X)\}\{\Delta^{bc}(q-Q/2,X)\}\;=\;\eta_a^{\ c}\theta(t-2|X_0|)\;,\end{aligned} \tag{4.48}$$

In the above expressions, of particular concern is the $\diamondsuit^+_{Q,X}$ operator, where the relative momentum Q and the central coordinate X are conjugate to one another. Thus, if the derivatives with respect to X are assumed to be small, then the derivatives with respect to Q must be large. In this case, all orders of the gradient expansion may be significant, so it is inappropriate to truncate to a given order in the soft derivative ∂_X^μ.

As $t\,\to\,\infty$, we have the transition $\delta_t^{(4)}(Q)\;\to\;\delta^{(4)}(Q)$ and (4.48) reduces to

$$e^{-i\diamondsuit^-_{q,X}}\{\Delta^{-1}_{ab}(q,X)\}\{\Delta^{bc}(q,X)\}\;=\;\eta_a^{\ c}\;. \tag{4.49}$$

Even for these late times, we can perform the matrix inversion exactly only if we truncate the gradient expansion in (4.49) to *zeroth* order. However, such a truncation appears valid only for time-independent *and* spatially homogeneous systems. Employing a suitable quasi-particle approximation to the Wigner representation of the propagators, it can be shown [55] that this inversion may be performed at first order in the gradient expansion. However, off-shell contributions are not fully accounted for in such an approximation.

In conclusion, a closed analytic form for the resummed CTP propagator may only be obtained in the time-independent thermodynamic equilibrium limit. The truncation of the gradient expansion may be justifiable only to the late-time evolution of systems very close to equilibrium, even for spatially homogeneous thermal backgrounds. A similar conclusion is drawn from different arguments in [67].

4.3 Non-Homogeneous Free Propagators

Unlike the resummed CTP propagator, the free CTP propagator can be derived analytically, even in the presence of a time-dependent and spatially inhomogeneous background. The non-homogeneous free propagator will account explicitly for the violation of space-time translational invariance. Our derivation relies on the algebra of the canonical quantization commutators of creation and annihilation operators described in Section 2. Subsequently, we make connection of our results with the path-integral representation of the CTP generating functional in (4.10). Finally, we introduce a diagrammatic representation for the non-homogeneous free CTP propagator.

We note that the derived propagators are 'free' in the sense that their spectral structure is that of single-particle states, corresponding to the free part of the action (see Section 3). Their statistical structure, on the other hand, will turn out to contain a summation over contributions from all possible multi-particle states. The time-dependent statistical distribution function appearing in these propagators is therefore a dressed object. This subtle point is significant for the consistent definition of the number density in Section 7 and the derivation of the master time evolution equations in Section 8.

The starting point of our canonical derivation is the explicit form of the density operator ρ. We relax any assumptions about the form of the density operator and take it to be in general non-diagonal but Hermitian within the general Fock space. We may write the most general interaction-picture density operator at the microscopic time $\tilde{t}_f \;=\; t/2$ as

$$\begin{aligned}\rho(\tilde{t}_f;\tilde{t}_i) \;&=\; C\exp\bigg[-\int \mathrm{d}\Pi_{\mathbf{k}_1}\;W_{10}(\mathbf{k}_1:0)a^\dagger(\mathbf{k}_1,\tilde{t}_f)\;-\;\int \mathrm{d}\Pi_{\mathbf{k}'_1}\;W_{01}(0:\mathbf{k}'_1)a(\mathbf{k}'_1,\tilde{t}_f)\\ &-\iint \mathrm{d}\Pi_{\mathbf{k}_1}\,\mathrm{d}\Pi_{\mathbf{k}'_1}\;W_{11}(\mathbf{k}_1:\mathbf{k}'_1)a^\dagger(\mathbf{k}_1,\tilde{t}_f)a(\mathbf{k}'_1,\tilde{t}_f)\;-\;\cdots \\ &-\frac{1}{n!}\frac{1}{m!}\int\cdots\int\bigg(\prod_{i=1}^{n}\mathrm{d}\Pi_{\mathbf{k}_i}\bigg)\bigg(\prod_{j=1}^{m}\mathrm{d}\Pi_{\mathbf{k}'_j}\bigg)\;W_{nm}(\{\mathbf{k}_i\}:\{\mathbf{k}'_j\})\prod_{i=1}^{n}a^\dagger(\mathbf{k}_i,\tilde{t}_f)\prod_{j=1}^{m}a(\mathbf{k}'_j,\tilde{t}_f)\bigg]\,,\end{aligned} \tag{4.50}$$

where the constant C can be set to unity without loss of generality. The complex-valued weights $W_{nm}(\{\mathbf{k}\}_n;\{\mathbf{k}'\}_m)$ depend on the state of the system at time $\tilde{t}_f \;=\; t/2$ and satisfy the Hermiticity constraint:

$$W_{nm}(\{\mathbf{k}\}_n:\{\mathbf{k}'\}_m)\;\;=\;\;W^*_{mn}(\{\mathbf{k}'\}_m:\{\mathbf{k}\}_n)\;. \tag{4.51}$$

The density operator ρ may be written in the basis of momentum eigenstates by multiplying the exponential form in (4.50) by the completeness relation of the basis of Fock states at time $\tilde{t}_f \;=\; t/2$:

$$\mathbf{I}\;=\;|0\rangle\langle 0|\;+\;\sum_{\ell\,=\,1}^{\infty}\frac{1}{\ell!}\bigg(\prod_{k=1}^{\ell}\int\mathrm{d}\Pi_{\mathbf{p}_k}\bigg)|\{\mathbf{p}\}_\ell,\tilde{t}_f\rangle\langle\{\mathbf{p}\}_\ell,\tilde{t}_f|\;, \tag{4.52}$$

where $|\{\mathbf{p}\}_\ell,\tilde{t}_f\rangle$ is the multi-mode Fock state $|\mathbf{p}_1,\tilde{t}_f\rangle\;\otimes\;|\mathbf{p}_2,\tilde{t}_f\rangle\;\otimes\;\cdots\;\otimes\;|\mathbf{p}_\ell,\tilde{t}_f\rangle$. This usually gives an intractable infinite series of n-to-m-particle correlations. Taking all weights $W_{nm}(\{\mathbf{k}\}_n:\{\mathbf{k}\}_m)$ to be zero if $n\;+\;m\;>\;2$, i.e. taking a Gaussian-like density operator, it is still possible to generate all possible n-to-m-particle correlations. In Appendix D, we give the expansion of the general Gaussian-like density operator, where only sufficient terms are included to help us visualize its analytic form.

Exploiting the remaining freedom in the commutation relations in (2.18), we may account for our ignorance of the series expansion of the density operator by writing the EEVs of two-point products of interaction-picture creation and annihilation operators as

$$\begin{aligned}\langle a^\dagger(\mathbf{p}',\tilde{t}_f;\tilde{t}_i)a(\mathbf{p},\tilde{t}_f;\tilde{t}_i)\rangle_t &= 2\,\mathscr{E}(\mathbf{p},\mathbf{p}')f(\mathbf{p},\mathbf{p}',t)\ , &(4.53a)\\ \langle a(\mathbf{p}',\tilde{t}_f;\tilde{t}_i)a(\mathbf{p},\tilde{t}_f;\tilde{t}_i)\rangle_t &= 2\,\mathscr{E}(\mathbf{p},\mathbf{p}')g(\mathbf{p},\mathbf{p}',t)\ . &(4.53b)\end{aligned}$$

The energy factor $2\mathscr{E}(\mathbf{p},\mathbf{p}')$, having dimensions E^1, arises from the fact that the 'number operator' $a^\dagger(\mathbf{p},\tilde{t};\tilde{t}_i)a(\mathbf{p},\tilde{t};\tilde{t}_i)$ of quantum field theory has dimensions E^{-2}, i.e. it does *not* have the dimensions of a number. Bearing in mind that the density operator is constructed from on-shell Fock states, a natural ansatz for this energy factor is

$$\mathscr{E}(\mathbf{p},\mathbf{p}') = \sqrt{E(\mathbf{p})E(\mathbf{p}')}\ . \tag{4.54}$$

The complex-valued distributions f and g have dimensions E^{-3} and satisfy the identities:

$$\begin{aligned}f(\mathbf{p},\mathbf{p}',t) &= f^*(\mathbf{p}',\mathbf{p},t)\ , &(4.55a)\\ g(\mathbf{p},\mathbf{p}',t) &= g(\mathbf{p}',\mathbf{p},t)\ . &(4.55b)\end{aligned}$$

We refer to f and g as *statistical distribution functions.* In particular, we interpret the Wigner transform

$$n(\mathbf{q},\mathbf{X},t) = \int\frac{\mathrm{d}^3\mathbf{Q}}{(2\pi)^3}\,e^{i\mathbf{Q}\cdot\mathbf{X}}f(\mathbf{q}+\tfrac{\mathbf{Q}}{2},\mathbf{q}-\tfrac{\mathbf{Q}}{2},t)\ , \tag{4.56}$$

having dimensions E^3, as the number density of particles at macroscopic time t in the phase-space hypervolume between $\mathbf{q}$ and $\mathbf{q}+\mathrm{d}^3\mathbf{q}$ and $\mathbf{X}$ and $\mathbf{X}+\mathrm{d}^3\mathbf{X}$. Notice that $n(\mathbf{q},\mathbf{X},t)$ is real thanks to the Hermiticity constraint (4.55a).

The EEV of the two-point product $\langle a(\mathbf{p},\tilde{t}_f;\tilde{t}_i)a^\dagger(\mathbf{p}',\tilde{t}_f;\tilde{t}_i)\rangle_t$ follows from the definition (4.53a) and the canonical commutation relation in (2.18), giving

$$\langle a(\mathbf{p},\tilde{t}_f;\tilde{t}_i)a^\dagger(\mathbf{p}',\tilde{t}_f;\tilde{t}_i)\rangle_t = (2\pi)^3 2E(\mathbf{p})\delta^{(3)}(\mathbf{p}-\mathbf{p}') + 2\,\mathscr{E}(\mathbf{p},\mathbf{p}')f(\mathbf{p},\mathbf{p}',t)\ . \tag{4.57}$$

Hermitian conjugation of (4.53b) yields

$$\langle a^\dagger(\mathbf{p},\tilde{t}_f;\tilde{t}_i)a^\dagger(\mathbf{p}',\tilde{t}_f;\tilde{t}_i)\rangle_t = 2\,\mathscr{E}(\mathbf{p},\mathbf{p}')g^*(\mathbf{p},\mathbf{p}',t)\ . \tag{4.58}$$

Note that (4.53), (4.57) and (4.58) are consistent with the canonical quantization rules in (2.14) and (2.17).

When the linear terms in the exponent of the density operator ρ in (4.50) are non-zero, we may consider the EEVs of single creation or annihilation operators

$$\langle a(\mathbf{p},\tilde{t}_f;\tilde{t}_i)\rangle_t = 2\sqrt{E(\mathbf{p})}w(\mathbf{p},t)\ . \tag{4.59}$$

In this case, we may define the connected distribution functions

$$f_{\text{con}}(\mathbf{p},\mathbf{p}',t) \;\equiv\; f(\mathbf{p},\mathbf{p}',t) \,-\, w(\mathbf{p},t)w(\mathbf{p}',t)\;, \tag{4.60a}$$

$$g_{\text{con}}(\mathbf{p},\mathbf{p}',t) \;\equiv\; g(\mathbf{p},\mathbf{p}',t) \,-\, w(\mathbf{p},t)w(\mathbf{p}',t)\;, \tag{4.60b}$$

which obey the same symmetry properties given in (4.55).

We are now in a position to derive the most general form of the double momentum representation of the non-homogeneous free CTP propagator, satisfying the inverse relation (3.33). Proceeding as in Section 3.2, we make the following ansatz for the most general solution of the Klein–Gordon equation in the double momentum representation:

$$\begin{aligned}\Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i) \;&=\; \begin{bmatrix} \big(p^2 \,-\, M^2 \,+\, i\epsilon\big)^{-1} & -i2\pi\theta(-p_0)\delta(p^2 \,-\, M^2) \\ -i2\pi\theta(p_0)\delta(p^2 \,-\, M^2) & -\big(p^2 \,-\, M^2 \,-\, i\epsilon\big)^{-1}\end{bmatrix}(2\pi)^4\delta^{(4)}(p-p') \\ &\quad -\, i2\pi|2p_0|^{1/2}\delta(p^2 \,-\, M^2)\tilde{f}(p,p',t)e^{i(p_0\,-\,p_0')\tilde{t}_f}2\pi|2p_0'|^{1/2}\delta(p'^2 \,-\, M^2)\begin{bmatrix}1 & 1\\ 1 & 1\end{bmatrix}\,,\end{aligned} \tag{4.61}$$

which we confirm by evaluating the EEVs directly, using the algebra of (4.53).

In (4.61) the phase factor $e^{i(p_0\,-\,p_0')\tilde{t}_f}$ arises from the fact that the creation and annihilation operators appearing in the Fourier transform of the field operator given in (2.12) are evaluated at the time $\tilde{t} \,=\, 0$. The density operator, on the other hand, is evaluated at the time $\tilde{t}_f$. As a consequence, in the evaluation of the EEV, we have, for instance,

$$\langle a^\dagger(\mathbf{p}',0;\tilde{t}_i)a(\mathbf{p},0;\tilde{t}_i)\rangle_t \;=\; 2\,\mathscr{E}(\mathbf{p},\mathbf{p}')f(\mathbf{p},\mathbf{p}',t)e^{i[E(\mathbf{p})\,-\,E(\mathbf{p}')]\tilde{t}_f}\;, \tag{4.62}$$

which directly results from (4.53a).

The form of the function $\tilde{f}(p,p',t)$ is

$$\begin{aligned}\tilde{f}(p,p',t) \;&=\; \theta(p_0)\theta(p_0')f(\mathbf{p},\mathbf{p}',t) \,+\, \theta(-p_0)\theta(-p_0')f^*(-\mathbf{p},-\mathbf{p}',t) \\ &\qquad +\, \theta(p_0)\theta(-p_0')g(\mathbf{p},-\mathbf{p}',t) \,+\, \theta(-p_0)\theta(p_0')g^*(-\mathbf{p},\mathbf{p}',t)\;.\end{aligned} \tag{4.63}$$

The function $\tilde{f}$ satisfies the relations: $\tilde{f}(p,p',t) = \tilde{f}(-p',-p,t) = \tilde{f}^*(-p,-p',t)$, consistent with the properties in (A.4). It also contains all information about the state of the ensemble at the macroscopic time t. For this reason, we refer to $\tilde{f}$ as the *ensemble function*.

In the double momentum representation, the retarded and advanced propagators are

$$\Delta^0_{\mathrm{R(A)}}(p,p') \;=\; \frac{1}{(p_0\,+(-)\,i\epsilon)^2 \,-\, \mathbf{p}^2 \,-\, M^2}(2\pi)^4\delta^{(4)}(p\,-\,p')\;. \tag{4.64}$$

The Pauli-Jordan $\Delta^0(p,p')$, Hadamard $\Delta^0_1(p,p',\tilde{t}_f;\tilde{t}_i)$ and principal-part $\Delta^0_{\mathcal{P}}(p,p')$ propagators become

$$\Delta^0(p,p') \;=\; -\,i2\pi\varepsilon(p_0)\delta(p^2 \,-\, M^2)(2\pi)^4\delta^{(4)}(p\,-\,p')\;, \tag{4.65a}$$

$$\begin{aligned}\Delta^0_1(p,p',\tilde{t}_f;\tilde{t}_i) \;&=\; -\,i2\pi\delta(p^2 \,-\, M^2)(2\pi)^4\delta^{(4)}(p\,-\,p') \\ &\quad -\, i2\pi|2p_0|^{1/2}\delta(p^2 \,-\, M^2)2\tilde{f}(p,p',t)e^{i(p_0\,-\,p_0')\tilde{t}_f}2\pi|2p_0'|^{1/2}\delta(p'^2 \,-\, M^2)\,,\end{aligned} \tag{4.65b}$$

$$\Delta^0_{\mathcal{P}}(p,p') \;=\mathcal{P}\frac{1}{p^2 \,-\, M^2}(2\pi)^4\delta^{(4)}(p\,-\,p')\;. \tag{4.65c}$$

Propagator	**Double Momentum Representation**
Feynman (Dyson)	$i\Delta^0_{\mathrm{F(D)}}(p,p',\tilde{t}_f;\tilde{t}_i) \;=\; \frac{(-)i}{p^2 \,-\, M^2 \,+ (-)\, i\epsilon}\,(2\pi)^4\delta^{(4)}(p\,-\,p')$ $+\, 2\pi\lvert 2p_0\rvert^{1/2}\delta(p^2 \,-\, M^2)\tilde{f}(p,p',t)e^{i(p_0\,-\,p'_0)\tilde{t}_f}2\pi\lvert 2p'_0\rvert^{1/2}\delta(p'^2 \,-\, M^2)$
$+(-)$ve-freq. Wightman	$i\Delta^0_{>(<)}(p,p',\tilde{t}_f;\tilde{t}_i) \;=\; 2\pi\theta(+(-)p_0)\delta(p^2 \,-\, M^2)(2\pi)^4\delta^{(4)}(p\,-\,p')$ $+\, 2\pi\lvert 2p_0\rvert^{1/2}\delta(p^2 \,-\, M^2)\tilde{f}(p,p',t)e^{i(p_0\,-\,p'_0)\tilde{t}_f}2\pi\lvert 2p'_0\rvert^{1/2}\delta(p'^2 \,-\, M^2)$
Retarded (Advanced)	$i\Delta^0_{\mathrm{R(A)}}(p,p') \;=\; \frac{i}{(p_0 \,+ (-)\, i\epsilon)^2 \,-\, \mathbf{p}^2 \,-\, M^2}(2\pi)^4\delta^{(4)}(p\,-\,p')$
Pauli–Jordan	$i\Delta^0(p,p') \;=\; 2\pi\varepsilon(p_0)\delta(p^2 \,-\, M^2)(2\pi)^4\delta^{(4)}(p\,-\,p')$
Hadamard	$i\Delta^0_1(p,p',\tilde{t}_f;\tilde{t}_i) \;=\; 2\pi\delta(p^2 \,-\, M^2)(2\pi)^4\delta^{(4)}(p\,-\,p')$ $+\, 2\pi\lvert 2p_0\rvert^{1/2}\delta(p^2 \,-\, M^2)2\tilde{f}(p,p',t)e^{i(p_0\,-\,p'_0)\tilde{t}_f}2\pi\lvert 2p'_0\rvert^{1/2}\delta(p'^2 \,-\, M^2)$
Principal-part	$i\Delta^0_{\mathcal{P}}(p,p') \;=\; \mathcal{P}\frac{i}{p^2 \,-\, M^2}(2\pi)^4\delta^{(4)}(p\,-\,p')$

Table 2: The full complement of non-homogeneous free propagators, where $\tilde{f}(p,p',t)$ is the ensemble function defined in (4.63) and $\mathcal{P}$ denotes the Cauchy principal value.

Thus, at the tree level, only the Hadamard correlation function $\Delta^0_1(p,p',\tilde{t}_f;\tilde{t}_i)$ depends explicitly on the background and macroscopic time t, through the ensemble function $\tilde{f}(p,p',t)$ in (4.63). This is a consequence of the causality of the theory, as we would expect from the spectral decomposition (2.22) of the retarded and advanced propagators $\Delta^0_{\mathrm{R(A)}}(x,y)$ in terms of the canonical commutation relation (2.14). Notice that the complex phase factor $e^{i(p_0\,-\,p'_0)\tilde{t}_f}$ has only appeared in the Hadamard propagator (4.65b) and so it does not spoil causality. Beyond the tree level, the background contributions are expected to modify the structure of the Pauli–Jordan and causal propagators, according to our discussion of the Källén–Lehmann spectral representation in Section 2. The full complement of non-homogeneous free propagators is listed in Table 2.

It would be interesting to establish a connection between these canonically derived non-homogeneous free propagators and those derived by the path-integral representation of the CTP generating functional $\mathcal{Z}$. This will be achieved through the bi-local source K_{ab} via the tree-level Schwinger–Dyson equation in (4.30). The role of the bi-local source K_{ab} will be illustrated further, when discussing the thermodynamic equilibrium limit in Section 5.

We proceed by replacing the exponent of the CTP generating functional $\mathcal{Z}$ in (4.10) by its double momentum representation. Subsequently, we may complete the square in this exponent by making the following shift in the field:

$$\Phi^a(p) \;=\; \Phi'^a(p) \,-\, \widehat{\Delta}^{0,\,a}{}_b(p)J^b(-p)\;, \tag{4.66}$$



Figure 3: The Feynman–diagrammatic interpretation of the non-homogeneous free CTP propagator $i\Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ for the real scalar Φ, where the double line represents momentum-violating coupling to the thermal background through the bi-local source $K_{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$.

where $\widehat{\Delta}^{0,\,ab}(p)$ is the free vacuum CTP propagator in (3.44) in which the ensemble function $\tilde{f}$ of (4.61) is set to zero. Notice that the normal-ordered contribution does not appear in $\widehat{\Delta}^{0,\,ab}(p)$, as it is sourced from the bi-local term K_{ab}. Upon substitution of (4.66) into (4.10), the CTP generating functional $\mathcal{Z}$ takes on the form

$$\begin{aligned}\mathcal{Z}[J,K,t] \ &= \ \mathcal{Z}^0[0,K,t]\exp\bigg\{iS^{\rm int}\bigg[\frac{1}{i}\frac{\delta}{\delta J_a},t\bigg]\bigg\}\int[\mathrm{d}\Phi'^a]\ \exp\bigg\{\frac{i}{2}\iint\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4p'}{(2\pi)^4} \\ &\bigg[J^a(p)\widehat{\Delta}^{0,}{}_a{}^c(p)K_{cb}(p,p',t)\Phi'^b(-p') \ + \ \Phi'^a(p)K_{ac}(p,p',t)\widehat{\Delta}^{0,\,c}{}_b(p')J^b(-p') \\ &- \ J^a(p)\Big(\widehat{\Delta}^0_{ab}(p)(2\pi)^4\delta^{(4)}(p\,-\,p') \ - \ \widehat{\Delta}^{0,}{}_a{}^c(p)K_{cd}(p,p',t)\widehat{\Delta}^{0,\,d}{}_b(p')\Big)\,J^b(-p')\bigg]\bigg\}\,. \end{aligned} \tag{4.67}$$

For the Lagrangian in (2.1), the cubic self-interaction part $(-g\Phi^3)$ of the action may be written down explicitly as

$$\begin{aligned}&\frac{-ig}{3!}\iiint\frac{\mathrm{d}^4p_1}{(2\pi)^4}\,\frac{\mathrm{d}^4p_2}{(2\pi)^4}\,\frac{\mathrm{d}^4p_3}{(2\pi)^4}\,\eta_{abc}(2\pi)^4\delta_t^{(4)}(p_1\,+\,p_2\,+\,p_3) \\ &\qquad\times\ \frac{1}{i}\frac{\delta}{\delta J_a(p_1)}\,\frac{1}{i}\frac{\delta}{\delta J_b(p_2)}\,\frac{1}{i}\frac{\delta}{\delta J_c(p_3)}\ \subset\ iS^{\rm int}\bigg[\frac{1}{i}\frac{\delta}{\delta J_a},t\bigg]\ .\end{aligned} \tag{4.68}$$

Hence, in the three-point vertex, the usual energy-conserving delta function has been replaced by δ_t, defined in (3.50), as a result of the systematic inclusion of finite-time effects. This time-dependent modification of the Feynman rules is fundamental to our perturbative approach to non-equilibrium thermal field theory and will be discussed further in Section 9 in the context of a simple scalar model.

In (4.67), the remaining terms linear in the external source J yield contributions to the free propagator proportional to K^2 upon double functional differentiation with respect to J. As we shall see in Section 5, these contributions may be neglected. Employing (3.25), we find that the non-homogeneous free CTP propagator $\Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ may be expressed in terms of the free vacuum CTP propagator $\widehat{\Delta}^{0,\,ab}(p)$ and the bi-local source $K_{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ as follows:

$$i\Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i) \ = \ i\widehat{\Delta}^{0,\,ab}(p)(2\pi)^4\delta^{(4)}(p\,-\,p') \ + \ i\widehat{\Delta}^{0,\,ac}(p)iK_{cd}(p,p',\tilde{t}_f;\tilde{t}_i)i\widehat{\Delta}^{0,\,db}(p')\ , \tag{4.69}$$

where

$$K_{ab}(p,p',\tilde{t}_f;\tilde{t}_i) \ = \ e^{i(p_0\,-\,p'_0)\tilde{t}_f}\,K_{ab}(\mathbf{p},\mathbf{p}',t)\ . \tag{4.70}$$

The form of the free CTP propagator $\Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ in (4.69) is consistent with a perturbative inversion of (4.30) to leading order in the bi-local source K_{ab}. It is also consistent with the canonically derived form of the non-homogeneous free propagators in (4.61).

The result in (4.69) may be interpreted diagrammatically, where the non-homogeneous free CTP propagator $i\Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ for the real scalar Φ is associated with the Feynman diagram displayed in Figure 3. The bi-local source $K_{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ plays the role of a three-momentum-violating vertex that gives rise to the violation of translational invariance, thus encoding the spatial inhomogeneity of the background.

5 The Thermodynamic Equilibrium Limit

In this section, we derive the analytical forms of the free and resummed CTP propagators in the limit of thermal equilibrium. The results of this section are of particular importance for the discussion of pinch singularities in Section 6. We also show the connection between the equilibrium Bose–Einstein distribution function and the bi-local source K_{ab} introduced in Section 4.3.

In the limit of thermal equilibrium, the density operator ρ is diagonal in particle number, so all amplitudes except W_{11} vanish in (4.50). In this limit, the general density operator ρ, given explicitly in (D.1), reduces to the series

$$\begin{aligned}\rho \;=\; & |0\rangle\langle 0| \,+\, \iint \mathrm{d}\Pi_{\mathbf{k}_1}\,\mathrm{d}\Pi_{\mathbf{k}_1'}\,\Big((2\pi)^3 2E(\mathbf{k}_1)\delta^{(3)}(\mathbf{k}_1 - \mathbf{k}_1') \,-\, W_{11}(\mathbf{k}_1:\mathbf{k}_1') \\ & \quad +\, \frac{1}{2}\int \mathrm{d}\Pi_{\mathbf{q}_1}\,W_{11}(\mathbf{k}_1:\mathbf{q}_1)W_{11}(\mathbf{q}_1:\mathbf{k}_1') \,+\, \cdots\Big)|\mathbf{k}_1\rangle\langle\mathbf{k}_1'| \\ & +\, \frac{1}{2}\int\cdots\int \mathrm{d}\Pi_{\mathbf{k}_1}\,\mathrm{d}\Pi_{\mathbf{k}_2}\,\mathrm{d}\Pi_{\mathbf{k}_1'}\,\mathrm{d}\Pi_{\mathbf{k}_2'}\,\Big((2\pi)^3 2E(\mathbf{k}_1)\delta^{(3)}(\mathbf{k}_1 - \mathbf{k}_1') \,-\, W_{11}(\mathbf{k}_1:\mathbf{k}_1') \\ & \quad +\, \frac{1}{2}\int \mathrm{d}\Pi_{\mathbf{q}_1}\,W_{11}(\mathbf{k}_1:\mathbf{q}_1)W_{11}(\mathbf{q}_1:\mathbf{k}_1') \,+\, \cdots\Big)\Big((2\pi)^3 2E(\mathbf{k}_2)\delta^{(3)}(\mathbf{k}_2 - \mathbf{k}_2') \\ & \quad -\, W_{11}(\mathbf{k}_2:\mathbf{k}_2') \,+\, \frac{1}{2}\int \mathrm{d}\Pi_{\mathbf{q}_2}\,W_{11}(\mathbf{k}_2:\mathbf{q}_2)W_{11}(\mathbf{q}_2:\mathbf{k}_2') \,+\, \cdots\Big)|\mathbf{k}_1,\mathbf{k}_2\rangle\langle\mathbf{k}_1',\mathbf{k}_2'| \,+\, \cdots\,,\end{aligned} \tag{5.1}$$

where the time arguments in the multi-mode Fock states have been omitted for notational convenience. In the equilibrium limit, the statistical distribution function $g(\mathbf{p},\mathbf{p'},t)$ is trivially zero. Instead, $f(\mathbf{p},\mathbf{p'},t)$ is calculated from (4.53) and takes the form of the series

$$\begin{aligned}2\mathcal{E}(\mathbf{p},\mathbf{p}')f(\mathbf{p},\mathbf{p}',t) \;=\; & (2\pi)^3 2E(\mathbf{p})\delta^{(3)}(\mathbf{p} - \mathbf{p}') \,-\, W_{11}(\mathbf{p}:\mathbf{p}') \\ & +\, \frac{1}{2}\int \mathrm{d}\Pi_{\mathbf{q}}\,W_{11}(\mathbf{p}:\mathbf{q})W_{11}(\mathbf{q}:\mathbf{p}') \,+\, \cdots \\ & +\, \int \mathrm{d}\Pi_{\mathbf{q}}\,\Big((2\pi)^3 2E(\mathbf{p})\delta^{(3)}(\mathbf{p} - \mathbf{q}) \,-\, W_{11}(\mathbf{p}:\mathbf{q}) \,+\, \cdots\Big) \\ & \quad \times\,\Big((2\pi)^3 2E(\mathbf{q})\delta^{(3)}(\mathbf{q} - \mathbf{p}') \,-\, W_{11}(\mathbf{q}:\mathbf{p}') \,+\, \cdots\Big) \,+\, \cdots\,,\end{aligned} \tag{5.2}$$

where disconnected parts have been cancelled order-by-order in the expansion by the normalization $\mathrm{Tr}\,\rho$ in (4.3). The factor $\mathcal{E}(\mathbf{p},\mathbf{p'})$ is defined in (4.54).

The equilibrium density operator $\rho_{\rm eq}$ must also be diagonal in the momenta and is thus obtained by making the replacement:

$$W_{11}(\mathbf{k}:\mathbf{k}') \;\to\; \beta E(\mathbf{k})(2\pi)^3\delta^{(3)}(\mathbf{k}\,-\,\mathbf{k}') \tag{5.3}$$

in (5.1), where $\beta\,=\,1/T$ is the inverse thermodynamic temperature. In detail, we find

$$\rho_{\rm eq} \;=\; |0\rangle\langle 0| \,+\, \sum_{n=1}^{\infty}\frac{1}{n!}\int\cdots\int\prod_{i=1}^{n}\Big(\mathrm{d}\Pi_{\mathbf{k}_i}\,f_\beta(\mathbf{k}_i)\Big)\bigotimes_{i=1}^{n}|\mathbf{k}_i\rangle\bigotimes_{i=1}^{n}\langle\mathbf{k}_i|\ , \tag{5.4}$$

where the amplitudes are the Boltzmann distributions $f_\beta(\mathbf{k})\,=\,e^{-\beta E(\mathbf{k})}$. This last expression of $\rho_{\rm eq}$ can be shown to be fully equivalent to the Gaussian form

$$\rho_{\rm eq} \;=\; \exp\bigg(-\,\beta\int\mathrm{d}\Pi_{\mathbf{k}}\,E(\mathbf{k})a^\dagger(\mathbf{k})a(\mathbf{k})\bigg)\ , \tag{5.5}$$

which corresponds to the standard Boltzmann density operator

$$\rho_{\rm eq} \;=\; e^{-\beta H^0}\ , \tag{5.6}$$

where H^0 is the free part of the interaction-picture Hamiltonian. Note that our convention for the normalization of the density operator ρ, including $\rho_{\rm eq}$, is chosen so that the canonical partition function $\mathcal{Z}(\beta)\,=\,\mathrm{Tr}\,e^{-\beta H^0}$ appears explicitly in the definition of the EEV in (4.3).

We may now substitute the limit (5.3) into the series expansion of the statistical distribution function f in (5.2). Using the identities of summation

$$\sum_{n=0}^{\infty}\frac{(-x)^n}{n!}\;=\;e^{-x}\ ,\qquad \sum_{n=1}^{\infty}x^n\;=\;\frac{x}{1-x}\ , \tag{5.7}$$

we find the following correspondence in the equilibrium limit:

$$f(\mathbf{p},\mathbf{p}',t)\;\underset{\rm eq.}{\longrightarrow}\;f_{\rm eq}(\mathbf{p},\mathbf{p}')\;=\;(2\pi)^3\delta^{(3)}(\mathbf{p}\,-\,\mathbf{p}')f_{\rm B}\big(E(\mathbf{p})\big)\ , \tag{5.8a}$$

$$g(\mathbf{p},\mathbf{p}',t)\;\underset{\rm eq.}{\longrightarrow}\;g_{\rm eq}(\mathbf{p},\mathbf{p}')\;=\;0\ , \tag{5.8b}$$

where

$$f_{\rm B}(p_0)\;=\;\frac{1}{e^{\beta p_0}\,-\,1} \tag{5.9}$$

is the Bose–Einstein distribution function. The equilibrium statistical distribution functions in (5.8) depend only on the magnitude of the three-momentum $\mathbf{p}$ via the on-shell energy $E(\mathbf{p})$. This is a consequence of the homogeneity and isotropy implied by thermodynamic equilibrium. Moreover, the multiplying factor $\delta^{(3)}(\mathbf{p}\,-\,\mathbf{p}')$ on the RHS of (5.8) necessarily restores translational and rotational invariance.

It is well-known that the pinch singularities present in perturbative expansions cancel in the equilibrium limit (see Section 6) and we can safely take the limit $t\,\to\,\infty$ throughout the CTP generating functional, as we should expect for a system with static macroscopic properties. Working then in the single momentum representation, we obtain from (3.42) and (3.43) the free equilibrium CTP propagators

$$i\Delta^0_{\rm F}(p) \;=\; \Big(i\Delta^0_{\rm D}(p)\Big)^* \;=\; i\Big(p^2 \,-\, M^2 \,+\, i\varepsilon\Big)^{-1} \,+\, 2\pi f_{\rm B}(|p_0|)\delta(p^2 \,-\, M^2)\;, \tag{5.10a}$$

$$i\Delta^0_{>}(p) \;=\; 2\pi\Big(\theta(p_0) \,+\, f_{\rm B}(|p_0|)\Big)\delta(p^2 \,-\, M^2) \;\equiv\; 2\pi\varepsilon(p_0)\Big(1 \,+\, f_{\rm B}(p_0)\Big)\delta(p^2 \,-\, M^2)\;, \tag{5.10b}$$

$$i\Delta^0_{<}(p) \;=\; 2\pi\Big(\theta(-p_0) \,+\, f_{\rm B}(|p_0|)\Big)\delta(p^2 - M^2) \;\equiv\; 2\pi\varepsilon(p_0)f_{\rm B}(p_0)\delta(p^2 \,-\, M^2)\;. \tag{5.10c}$$

The form of the Wightman propagators written in terms of the signum function $\varepsilon(p_0)$ prove very useful in the calculation of loop diagrams, as detailed in Appendix B.

Returning to the free CTP generating functional $\mathcal{Z}$ in (4.67), it follows from the results above that in equilibrium the bi-local source $K_{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ must be proportional to a four-dimensional delta function of the momenta, i.e.

$$K_{ab}(p,p',\tilde{t}_f;\tilde{t}_i) \;\underset{\rm eq.}{\longrightarrow}\; K^{\rm eq}_{ab}(p,p') \;=\; (2\pi)^4\delta^{(4)}(p \,-\, p')K^{\rm eq}_{ab}(p)\;. \tag{5.11}$$

In addition, $K^{\rm eq}_{ab}(p)$ must satisfy

$$\widehat{\Delta}^{0,\,ac}(p)K^{\rm eq}_{cd}(p)\widehat{\Delta}^{0,\,db}(p) \;=\; 2\pi i\,\delta(p^2 \,-\, M^2)f_{\rm B}\big(E(\mathbf{p})\big)\begin{bmatrix}1 & 1\\ 1 & 1\end{bmatrix}\;. \tag{5.12}$$

Solving the resulting system of equations, keeping terms to leading order in ϵ, and noting that $K^{\rm eq}_{ab}(p)$ should be written in terms of the three-momentum only, we find

$$K^{\rm eq}_{ab}(p) \;=\; 2i\epsilon f_{\rm B}\big(E(\mathbf{p})\big)\begin{bmatrix}1 & 1\\ 1 & 1\end{bmatrix}\;. \tag{5.13}$$

By virtue of the limit representation of the delta function

$$\delta(x) \;=\; \lim_{\epsilon\,\to\,0}\,\frac{1}{\pi}\,\frac{\epsilon}{x^2 \,+\, \epsilon^2}\;, \tag{5.14}$$

we can verify that we do indeed recover (5.13) and the correct free CTP propagator by means of (4.69). We also confirm that the terms linear in J remaining in (4.67) may safely be ignored, since they yield contributions to the free propagator proportional to $K^2 \,\sim\, \epsilon^2$ upon double functional differentiation with respect to J.

Alternatively, interpreting the Boltzmann density operator in (5.6) as an evolution operator in negative imaginary time and using the cyclicity of the trace in the EEV, we derive the Kubo–Martin–Schwinger (KMS) relation (see for instance [68])

$$\Delta_>(x^0 - y^0, \mathbf{x} - \mathbf{y}) \;=\; \Delta_<(x^0 - y^0 + i\beta, \mathbf{x} - \mathbf{y})\;. \tag{5.15}$$

In the momentum representation, the KMS relation reads:

$$\Delta_>(p) \;=\; e^{\beta p_0}\Delta_<(p)\;, \tag{5.16}$$

which offers the final constraint on $\tilde{f}(p)$ in (3.43):

$$\tilde{f}(p) \;=\; \theta(p_0)f_{\rm B}(p_0) + \theta(-p_0)f_{\rm B}(-p_0) \;=\; f_{\rm B}(|p_0|)\;. \tag{5.17}$$

Furthermore, the KMS relation also leads to the fluctuation-dissipation theorem

$$\Delta_1(p) \;=\; \Big(1\;+\;2f_{\rm B}(p_0)\Big)\,\Delta(p)\;, \tag{5.18}$$

relating the causality and unitarity relations in (2.24) and (2.28). Subsequently, by means of the KMS relation, we may write all propagators in terms of the retarded propagator $\Delta_{\rm R}(p)$:

$$\begin{aligned}
{\rm Re}\,\Delta_{\rm F}(p) \;&=\; {\rm Re}\,\Delta_{\rm R}(p)\;, &\text{(5.19a)}\\
{\rm Im}\,\Delta_{\rm F}(p) \;&=\; \varepsilon(p_0)\Big(1\;+\;2f_{\rm B}(|p_0|)\Big){\rm Im}\,\Delta_{\rm R}(p)\;, &\text{(5.19b)}\\
\Delta_{>}(p) \;&=\; 2i\varepsilon(p_0)\Big(\theta(p_0)\;+\;f_{\rm B}(|p_0|)\Big){\rm Im}\,\Delta_{\rm R}(p)\;, &\text{(5.19c)}\\
\Delta_{<}(p) \;&=\; 2i\varepsilon(p_0)\Big(\theta(-p_0)\;+\;f_{\rm B}(|p_0|)\Big){\rm Im}\,\Delta_{\rm R}(p)\;. &\text{(5.19d)}
\end{aligned}$$

In the homogeneous equilibrium limit of the Schwinger–Dyson equation in (4.29), the inverse resummed CTP propagator is given by

$$\Delta^{-1}_{ab}(p) \;=\; \Delta^{0,\,-1}_{ab}(p)\;+\;\Pi_{ab}(p)\;. \tag{5.20}$$

In the absence of self-energy effects, the free equilibrium CTP propagator is obtained by inverting the equilbrium limit of (4.30):

$$D^{0,\,-1}_{ab}(p) \;=\; \Delta^{0,\,-1}_{ab}(p)\;+\;K^{\rm eq}_{ab}(p)\;. \tag{5.21}$$

Knowing that $K^{\rm eq}_{ab}\;\sim\;\epsilon$ from (5.13), the inversion of (5.21) can be done perturbatively to leading order in $K^{\rm eq}_{ab}(p)$, in which case the expression (4.69) gets reproduced. Beyond the tree-level, however, the contribution from the bi-local source $K^{\rm eq}_{ab}(p)$ may be neglected next to the self-energy term $\Pi_{ab}(p)$ and the inverse resummed CTP propagator is explicitly given by

$$\Delta^{-1}_{ab}(p) \;=\; \begin{bmatrix} p^2\;-\;M^2\;+\;\Pi(p) & -\Pi_{<}(p)\\ -\Pi_{>}(p) & -p^2\;+\;M^2\;-\;\Pi^*(p)\end{bmatrix}\,. \tag{5.22}$$

In this equilibrium limit, (5.22) may be inverted exactly, yielding the *equilibrium* resummed CTP propagator

$$\begin{aligned}
\Delta^{ab}(p) \;=\; &\bigg[\Big(p^2\;-\;M^2\;+\;{\rm Re}\,\Pi_{\rm R}(p)\Big)^2\;+\;\Big({\rm Im}\,\Pi_{\rm R}(p)\Big)^2\bigg]^{-1}\\
&\times\begin{bmatrix} p^2\;-\;M^2\;+\;\Pi^*(p) & -\Pi_{<}(p)\\ -\Pi_{>}(p) & -p^2\;+\;M^2\;-\;\Pi(p)\end{bmatrix}\,.
\end{aligned} \tag{5.23}$$

In this single momentum representation, the self-energies satisfy the unitarity and causality relations

$$\begin{aligned}
\Pi_1(p) \;&=\; \Pi_{>}(p)\;+\;\Pi_{<}(p) \;=\; \Pi(p)\;-\;\Pi^*(p) \;=\; 2i{\rm Im}\,\Pi(p)\;, &\text{(5.24a)}\\
2iM\Gamma(p) \;&=\; \Pi_{>}(p)\;-\;\Pi_{<}(p) \;=\; \Pi_{\rm R}(p)\;-\;\Pi_{\rm A}(p) \;=\; 2i{\rm Im}\,\Pi_{\rm R}(p)\;, &\text{(5.24b)}
\end{aligned}$$

where $\Gamma(p)$ is the Breit–Wigner width, relating the absorptive part of the retarded self-energy $\Pi_{\rm R}(p)$ to physical reaction rates [69, 70]. Notice that the KMS relation (5.15) leads also to the detailed balance condition

$$\Pi_{>}(p) \ = \ e^{\beta p_0}\, \Pi_{<}(p)\ . \tag{5.25}$$

Given the relations in (5.24), we find, in compliance with (5.19), an analogous set of relations for the elements of the CTP self-energy:

$$\begin{aligned}
{\rm Re}\,\Pi(p) \ &= \ {\rm Re}\,\Pi_{\rm R}(p)\ , && (5.26a)\\
{\rm Im}\,\Pi(p) \ &= \ \varepsilon(p_0)\Big(1\ +\ 2f_{\rm B}(|p_0|)\Big){\rm Im}\,\Pi_{\rm R}(p)\ , && (5.26b)\\
\Pi_{>}(p) \ &= \ 2i\varepsilon(p_0)\Big(\theta(p_0)\ +\ f_{\rm B}(|p_0|)\Big){\rm Im}\,\Pi_{\rm R}(p)\ , && (5.26c)\\
\Pi_{<}(p) \ &= \ 2i\varepsilon(p_0)\Big(\theta(-p_0)\ +\ f_{\rm B}(|p_0|)\Big){\rm Im}\,\Pi_{\rm R}(p)\ . && (5.26d)
\end{aligned}$$

Ignoring the dispersive parts of the self-energy, we expect to recover the free CTP propagators given in (5.10) in the limit ${\rm Im}\,\Pi(p)\ \to\ \epsilon\ =\ 0^+$. This limit is equivalent to

$${\rm Im}\,\Pi_{\rm R}(p)\ \to\ \epsilon_{\rm R}\ \equiv\ \varepsilon(p_0)\,\epsilon\ . \tag{5.27}$$

Expressing the equilibrium resummed CTP propagator in (5.23) in terms of the retarded absorptive self-energy ${\rm Im}\,\Pi_{\rm R}(p)$, we can convince ourselves that we do indeed reproduce the free equilibrium CTP propagators (5.10) in the limit (5.27).

In Appendix B, we discuss the correspondence of the results of this section with the Imaginary Time Formalsim (ITF), clarifying the analytic continuation of the imaginary-time propagator and self-energy.

6 Perturbative One-Loop Expansion without Pinch Singularities

Pinch singularities normally spoil the perturbation expansion of non-equilibrium Green's functions [71]. These pathologies arise from products of delta functions with identical arguments. In this section, we explicitly demonstrate, in contrast to [72], how pinch singularities do not occur in our perturbative approach.

The avoidance of pinch singularities is achieved by the systematic inclusion of finite-time effects, which results in finite upper and lower bounds on time domains, as shown in Figure 1. Thus, our approach is different from the semi-infinite time domains employed in the existing literature [73, 74]. In particular, in our approach the finite-time effects are not included *a priori* and the removal of the pinch singularities is achieved without any *ad hoc* regularization or obscure resummation. Most importantly, proper consideration is given to the dependence upon the time of observation t. As noted in Section 4.3, the statistical distribution functions appearing in the non-homogeneous free propagators are dressed objects, evaluated at the macroscopic time t. Hence, the role of the Feynman–Dyson series is to dress the *spectral* structure of the propagators only. The evolution of the statistical distribution functions is determined by master time evolution equations to be derived in Section 8.

For early times, the appearance of pinch singularities is prevented by microscopic violation of energy conservation. At late times, the thermalization of the system ensures that pinch singularities cancel by virtue of the KMS relation (5.15). In this section, we demonstrate this cancellation explicitly at the one-loop level and describe the absence of pinch singularities in our approach for all times.

In order to proceed perturbatively, we truncate the Feynman–Dyson series in (4.37) to leading order in the couplings and set the bi-local source K_{ab} to zero. This corresponds to keeping free CTP propagators $\Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ and one-loop CTP self-energies $\Pi^{(1)}_{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ on the RHS of (4.37). We then have the one-loop-inserted CTP propagator

$$\begin{aligned}\Delta^{(1),\,ab}(p,p',\tilde{t}_f;\tilde{t}_i) \;&=\; \Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i)\\ &-\iint\frac{\mathrm{d}^4q}{(2\pi)^4}\,\frac{\mathrm{d}^4q'}{(2\pi)^4}\;\Delta^{0,\,ac}(p,q,\tilde{t}_f;\tilde{t}_i)\Pi^{(1)}_{cd}(q,q',\tilde{t}_f;\tilde{t}_i)\Delta^{0,\,db}(q',p',\tilde{t}_f;\tilde{t}_i)\;. \qquad (6.1)\end{aligned}$$

It will prove convenient to work in a mixed CTP–Keldysh basis by inserting the transformation outlined in (3.18) between the external legs and self-energies on the RHS of (6.1). In this mixed CTP–Keldysh basis, we may rewrite (6.1) as

$$\begin{aligned}\Delta^{(1),\,ab}(p,p',\tilde{t}_f;\tilde{t}_i) \;&=\; \Delta^{0,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i)\\ &-\frac{1}{2}\iint\frac{\mathrm{d}^4q}{(2\pi)^4}\,\frac{\mathrm{d}^4q'}{(2\pi)^4}\;\Delta^{0,\,ac}_{\mathrm{ret}}(p,q,\tilde{t}_f;\tilde{t}_i)\widetilde{\Pi}^{(1)}_{cd}(q,q',\tilde{t}_f;\tilde{t}_i)\Delta^{0,\,db}_{\mathrm{adv}}(q',p',\tilde{t}_f;\tilde{t}_i)\;, \qquad (6.2)\end{aligned}$$

where, making use of the relations in (A.6),

$$\begin{aligned}\Delta^{ac}_{\mathrm{ret}}(p,q,\tilde{t}_f;\tilde{t}_i) \;&=\; \begin{bmatrix}\Delta_{\mathrm{F}}(p,q,\tilde{t}_f;\tilde{t}_i) & \Delta_{<}(p,q,\tilde{t}_f;\tilde{t}_i)\\ \Delta_{>}(p,q,\tilde{t}_f;\tilde{t}_i) & -\Delta^{*}_{\mathrm{F}}(p,q,\tilde{t}_f;\tilde{t}_i)\end{bmatrix}\begin{bmatrix}1 & 1\\ -1 & 1\end{bmatrix} \qquad (6.3)\\ &=\; \begin{bmatrix}\Delta_{\mathrm{R}}(p,q,\tilde{t}_f;\tilde{t}_i) & \Delta_{\mathrm{R}}(p,q,\tilde{t}_f;\tilde{t}_i)\;+\;2\Delta_{<}(p,q,\tilde{t}_f;\tilde{t}_i)\\ \Delta_{\mathrm{R}}(p,q,\tilde{t}_f;\tilde{t}_i) & -\Delta_{\mathrm{R}}(p,q,\tilde{t}_f;\tilde{t}_i)\;+\;2\Delta_{>}(p,q,\tilde{t}_f;\tilde{t}_i)\end{bmatrix}\,,\end{aligned}$$

and

$$\begin{aligned}\Delta^{ac}_{\mathrm{adv}}(p,q,\tilde{t}_f;\tilde{t}_i) \;&=\; \begin{bmatrix}1 & -1\\ 1 & 1\end{bmatrix}\begin{bmatrix}\Delta_{\mathrm{F}}(p,q,\tilde{t}_f;\tilde{t}_i) & \Delta_{<}(p,q,\tilde{t}_f;\tilde{t}_i)\\ \Delta_{>}(p,q,\tilde{t}_f;\tilde{t}_i) & -\Delta^{*}_{\mathrm{F}}(p,q,\tilde{t}_f;\tilde{t}_i)\end{bmatrix} \qquad (6.4)\\ &=\; \begin{bmatrix}\Delta_{\mathrm{A}}(p,q,\tilde{t}_f;\tilde{t}_i) & \Delta_{\mathrm{A}}(p,q,\tilde{t}_f;\tilde{t}_i)\\ \Delta_{\mathrm{A}}(p,q,\tilde{t}_f;\tilde{t}_i)\;+\;2\Delta_{>}(p,q,\tilde{t}_f;\tilde{t}_i) & -\Delta_{\mathrm{A}}(p,q,\tilde{t}_f;\tilde{t}_i)\;+\;2\Delta_{<}(p,q,\tilde{t}_f;\tilde{t}_i)\end{bmatrix}\,.\end{aligned}$$

In the same mixed CTP–Keldysh basis, the one-loop-inserted Feynman $\Delta^{(1)}_{\mathrm{F}}(p,p',\tilde{t}_f;\tilde{t}_i)$ and positive-frequency Wightman $\Delta^{(1)}_{>}(p,p',\tilde{t}_f;\tilde{t}_i)$ propagators may be written in the following forms:

$$\Delta^{(1)}_{\rm F}(p,p',\tilde{t}_f;\tilde{t}_i) \ = \ \Delta^{0}_{\rm F}(p,p',\tilde{t}_f;\tilde{t}_i) \ - \iint \frac{{\rm d}^4 q}{(2\pi)^4}\,\frac{{\rm d}^4 q'}{(2\pi)^4}\,\Big(\Delta^{0}_{\rm R}(p,q)\Pi^{(1)}(q,q',\tilde{t}_f;\tilde{t}_i)\Delta^{0}_{\rm A}(q',p')$$
$$+ \ \Delta^{0}_{\rm R}(p,q)\Pi^{(1)}_{\rm R}(q,q',\tilde{t}_f;\tilde{t}_i)\Delta^{0}_{>}(q',p',\tilde{t}_f;\tilde{t}_i) \ + \ \Delta^{0}_{<}(p,q,\tilde{t}_f;\tilde{t}_i)\Pi^{(1)}_{\rm A}(q,q',\tilde{t}_f;\tilde{t}_i)\Delta^{0}_{\rm A}(q',p')\Big)\ , \tag{6.5a}$$

$$\Delta^{(1)}_{>}(p,p',\tilde{t}_f;\tilde{t}_i) \ = \ \Delta^{0}_{\rm F}(p,p',\tilde{t}_f;\tilde{t}_i) \ - \iint \frac{{\rm d}^4 q}{(2\pi)^4}\,\frac{{\rm d}^4 q'}{(2\pi)^4}\,\Big(\Delta^{0}_{\rm R}(p,q)\Pi^{(1)}_{>}(q,q',\tilde{t}_f;\tilde{t}_i)\Delta^{0}_{\rm A}(q',p')$$
$$+ \ \Delta^{0}_{\rm R}(p,q)\Pi^{(1)}_{\rm R}(q,q',\tilde{t}_f;\tilde{t}_i)\Delta^{0}_{>}(q',p',\tilde{t}_f;\tilde{t}_i) \ + \ \Delta^{0}_{>}(p,q,\tilde{t}_f;\tilde{t}_i)\Pi^{(1)}_{\rm A}(q,q',\tilde{t}_f;\tilde{t}_i)\Delta^{0}_{\rm A}(q',p')\Big)\ , \tag{6.5b}$$

where we have used the identities

$$\Pi_1(p,p',\tilde{t}_f;\tilde{t}_i) \ + \ \Pi_{\rm R}(p,p',\tilde{t}_f;\tilde{t}_i) \ + \ \Pi_{\rm A}(p,p',\tilde{t}_f;\tilde{t}_i) \ = \ 2\Pi(p,p',\tilde{t}_f;\tilde{t}_i)\ , \tag{6.6a}$$
$$\Pi_1(p,p',\tilde{t}_f;\tilde{t}_i) \ + \ \Pi_{\rm R}(p,p',\tilde{t}_f;\tilde{t}_i) \ - \ \Pi_{\rm A}(p,p',\tilde{t}_f;\tilde{t}_i) \ = \ 2\Pi_{>}(p,p',\tilde{t}_f;\tilde{t}_i)\ , \tag{6.6b}$$

which can be derived from relations between the self-energies analogous to those in (A.6).

In the equilibrium limit, where translational invariance is restored, the expansions in (6.5a) and (6.5b) can be rewritten in the single momentum representation as follows:

$$\Delta^{(1)}_{\rm F}(p) \ = \ \Delta^{0}_{\rm F}(p) \tag{6.7a}$$
$$- \ \Big(\Delta^{0}_{\rm R}(p)\Pi^{(1)}(p)\Delta^{0}_{\rm A}(p) \ + \ \Delta^{0}_{\rm R}(p)\Pi^{(1)}_{\rm R}(p)\Delta^{0}_{>}(p) \ + \ \Delta^{0}_{<}(p)\Pi^{(1)}_{\rm A}(p)\Delta^{0}_{\rm A}(p)\Big)\ ,$$
$$\Delta^{(1)}_{>}(p) \ = \ \Delta^{0}_{>}(p) \tag{6.7b}$$
$$- \ \Big(\Delta^{0}_{\rm R}(p)\Pi^{(1)}_{>}(p)\Delta^{0}_{\rm A}(p) \ + \ \Delta^{0}_{\rm R}(p)\Pi^{(1)}_{\rm R}(p)\Delta^{0}_{>}(p) \ + \ \Delta^{0}_{>}(p)\Pi^{(1)}_{\rm A}(p)\Delta^{0}_{\rm A}(p)\Big)\ ,$$

where the single momentum representations of the equilibrium propagators and self-energies are those discussed in Section 5. Using the identities in (5.19) and (5.26), derived by virtue of the KMS relation (5.15) and detailed balance condition (5.25), we can rewrite (6.7a) and (6.7b) entirely in terms of the retarded functions $\Delta^{0}_{\rm R}(p)$ and $\Pi^{(1)}_{\rm R}(p)$. In this way, it can be shown that undesirable terms proportional to $\delta^2(p^2\ -\ M^2)$, which would signify pinch singularities, cancel exactly. We are then left with terms depending upon the following product:

$${\rm Re}\,\Delta^{0}_{\rm R}(p)\,{\rm Im}\,\Delta^{0}_{\rm R}(p)\ , \tag{6.8}$$

where

$${\rm Re}\,\Delta^{0}_{\rm R}(p) \ = \ \mathcal{P}\frac{1}{p^2\ -\ M^2}\ , \tag{6.9}$$

with $\mathcal{P}$ denoting the principal value integral, and

$${\rm Im}\,\Delta^{0}_{\rm R}(p) \ = \ -\pi\varepsilon(p_0)\delta(p^2\ -\ M^2)\ . \tag{6.10}$$

In order to show that (6.8) is also free of pinch singularities, we consider the limit representation of the Cauchy principle value:

$$\mathcal{P}\frac{1}{x} \ = \ \lim_{\epsilon\,\to\,0}\frac{x}{x^2\ +\ \epsilon^2} \tag{6.11}$$

and the limit representation of the delta function in (5.14). We may then show that

$$\mathrm{Re}\,\Delta^0_{\mathrm{R}}(p)\,\mathrm{Im}\,\Delta^0_{\mathrm{R}}(p) \;=\; \frac{\pi}{2}\varepsilon(p_0)\delta'(p^2 - M^2)\,, \tag{6.12}$$

where $\delta'(x)$ is the derivative of the delta function, satisfying

$$\int_{-\infty}^{+\infty} \mathrm{d}x\;\delta'(x)y(x) \;=\; -y'(0)\,, \tag{6.13}$$

provided the function $y(x)$ is analytically well-behaved at $x = 0$. Consequently, the expression (6.8) is free of pinch singularities.

Next, we use (6.12) and the free CTP propagators from (5.10) for the remaining terms in (6.7a) and (6.7b). This allows us to find a closed form for the one-loop-inserted equilibrium Feynman and Wightman propagators

$$\Delta^{(1)}_{\mathrm{F}}(p) \;=\; \frac{p^2 - M^2 + \Pi^{(1)}(p)}{\big(p^2 - M^2 + i\varepsilon\big)^2} \;+\; 2\pi i f_{\mathrm{B}}(|p_0|)\Big(\,\delta(p^2 - M^2) + \widehat{\Pi}^{*(1)}(p)\delta'(p^2 - M^2)\Big)\,, \tag{6.14a}$$

$$\begin{aligned}\Delta^{(1)}_{\gtrless}(p) \;=\;& \frac{\Pi^{(1)}_{\gtrless}(p)}{\big(p^2 - M^2 + i\varepsilon\big)^2} \\ &+ 2\pi i\Big(\,\theta(\pm p_0) + f_{\mathrm{B}}(|p_0|)\Big)\Big(\,\delta(p^2 - M^2) + \widehat{\Pi}^{*(1)}(p)\delta'(p^2 - M^2)\Big)\,,\end{aligned} \tag{6.14b}$$

where

$$\widehat{\Pi}(p) \;=\; \mathrm{Re}\,\Pi_{\mathrm{R}}(p) \;+\; i\varepsilon(p_0)\mathrm{Im}\,\Pi_{\mathrm{R}}(p) \tag{6.15}$$

is a self-energy-like function, bearing by itself no direct physical meaning. In (6.14b), we have included the negative-frequency Wightman propagator $\Delta^{(1)}_{<}(p)$ for completeness. The results in (6.14) are consistent with the properties and relations in Appendix A and reduce to the expected form in the zero-temperature limit. In conclusion, we have shown that the one-loop perturbative expansion of the equilibrium CTP propagator is well-defined and that potential pinch singularities cancel exactly in the thermodynamic equilibrium limit as a result of the KMS relation.

In non-equilibrium situations, the KMS relation does not hold. It would appear therefore that the application of perturbation theory in the regime of out of thermodynamic equilibrium would be plagued by pinch singularities. However, for finite t, the usual energy-conserving delta functions have been replaced by the time-dependent energy-non-conserving weight functions δ_t, defined in (3.50). Thus, for finite t, pinch singularities do not occur, even for general statistical distribution functions, since products of δ_t's with identical arguments remain analytic.

In the limit $t \to \infty$, one would naively expect pinch singularities to reappear, since $\delta_t(x) \to \delta(x)$ as given in (3.51). However, in the same limit, our boundary conditions are specified in the infinitely distant past. Hence, our system must have reached thermodynamic equilibrium. Thus, in the limit $t \to \infty$, the time-dependent statistical distribution functions appearing in the non-homogeneous free propagators are replaced with equilibrium Bose–Einstein distributions by the correspondence in (5.8). With this replacement, the pinch singularities cancel by virtue of the KMS relation, as described at the beginning of this section.

With the absence of pinch singularities established for early and infinitely-late times, we must now illustrate that the perturbation series remains free of these pathologies for all intermediate times. For this to be the case, we must demonstrate that the thermalization of the system takes place more rapidly than the growth of the pinch singularities discussed above.

Potential pinch singularities arise from terms like

$$g^2\delta_t^2(x)\ , \tag{6.16}$$

where the argument x and the coupling constant g have mass dimensions. For large t, this term is dominated by the contribution from $x \to 0$. By l'Hôpitals rule and (3.51), we may show that these pathologies grow like

$$g^2\delta_t^2(x) \underset{t\,\gg\,1/M}{\longrightarrow} \frac{gt}{4\pi}\, g\delta(x)\ . \tag{6.17}$$

As a consequence, any order of pinch singularity will grow like $(gt)^n$, where $n \in \mathbb{N}$ is some finite number. Therefore, in order for the perturbation series to be free of pinch singularities for all times, the deviation from thermodynamic equilibrium must decay more rapidly than a power law in the dimensionless parameter gt.

In the spatially homogeneous limit, we may approximate the late-time behaviour of the statistical distribution function f by the following master equation:

$$\frac{\mathrm{d}f(|\mathbf{p}|,t)}{\mathrm{d}t}\ =\ -\Gamma_>(p,t)f(|\mathbf{p}|,t)\ +\ \Gamma_<(p,t)\Big(1\ +\ f(|\mathbf{p}|,t)\Big)\ , \tag{6.18}$$

where the four-momentum $p^\mu = (E(\mathbf{p}),\mathbf{p})$ is on-shell. The master equation (6.18) is exactly the classical Boltzmann transport equation. The terms $\Gamma_>(p,t)\ =\ -i\Pi_>(p,t)/2M$ and $\Gamma_<(p,t) = -i\Pi_<(p,t)/2M$ relate to the on-shell decay and production rates, respectively. If the system is close to equilibrium, which is reasonable for late but nonetheless finite times, we may assume that $\Gamma_{>(<)}(p,t) \equiv \Gamma_{>(<)}(p)$ are approximately constant. Expanding around the equilibrium Bose–Einstein distribution f_{B}, we write

$$f(|\mathbf{p}|,t)\ =\ f_{\mathrm{B}}\big(E(\mathbf{p})\big)\ +\ \delta f(|\mathbf{p}|,t)\ . \tag{6.19}$$

Substituting this expansion into (6.18), the terms involving $f_{\mathrm{B}}\big(E(\mathbf{p})\big)$ cancel by virtue of the detailed balance condition (5.25) and the properties of the Bose–Einstein distribution. We then arrive at a first-order differential equation for the deviation $\delta f(|\mathbf{p}|,t)$:

$$\frac{\mathrm{d}\delta f(|\mathbf{p}|,t)}{\mathrm{d}t}\ =\ -\Big(\Gamma_>(p)\ -\ \Gamma_<(p)\Big)\ \delta f(|\mathbf{p}|,t)\ . \tag{6.20}$$

Notice that $\Gamma_>(p) - \Gamma_<(p)$ is the total Breit–Wigner width $\Gamma(p) \sim g^2$, defined in (5.24b). Thus,

$$\delta f(|\mathbf{p}|,t)\ =\ \exp\Big(-\Gamma(p)\,t\Big)\ \delta f(|\mathbf{p}|,0)\ , \tag{6.21}$$

which decays faster than any power law in gt, as required.

In summary, for early times, the analytic t-dependent vertices lead to microscopic violation of energy conservation. This energy non-conservation regularizes potential pinch singularities. For intermediate times, the t-dependent statistical distribution functions approach equilibrium faster than energy conservation is restored. In the limit $t \to \infty$, the time-dependent statistical distribution functions appearing in the non-homogeneous free propagators are replaced by their equilibrium forms via the correspondence in (5.8). We then obtain the well-known equilibrium thermal field theory in which energy conservation is fully restored and pinch singularities cancel exactly by virtue of the KMS relation. In conlusion, we have explicitly demonstrated how pinch singularities do not arise in our perturbative approach to non-equilibrium thermal field theory.

7 The Particle Number Density

It is important to establish a direct connection between off-shell Green's functions and physical observables. Such observables include the particle number density for which various interpretations have been reported in the literature [41–50]. In this section, we derive a physically meaningful definition of the particle number density in terms of the resummed CTP propagators.

In order to count off-shell contributions systematically, we suggest to 'measure' the particle number density in terms of charges, rather than by quanta of energy. The latter approach would necessitate the use of a quasi-particle approximation to identify 'single-particle' energies, which we do not follow here. Instead, we begin by introducing the Noether charge $\mathcal{Q}(x_0;\tilde{t}_i)$ of the Heisenberg-picture scalar field $\Phi_{\rm H}(x;\tilde{t}_i)$:

$$\mathcal{Q}(x_0;\tilde{t}_i) \ = \ i\int \mathrm{d}^3\mathbf{x}\, \Big(\Phi^\dagger_{\rm H}(x;\tilde{t}_i)\pi^\dagger_{\rm H}(x;\tilde{t}_i) \ - \ \pi_{\rm H}(x;\tilde{t}_i)\Phi_{\rm H}(x;\tilde{t}_i) \Big)\ . \tag{7.1}$$

Here, we include all time-dependencies explicitly for clarity. We recall that $\pi_{\rm H}(x;\tilde{t}_i) = \partial_{x_0}\Phi^\dagger_{\rm H}(x;\tilde{t}_i)$ is the conjugate momentum operator. Up to the usual infinite $T = 0$ vacuum contribution, the EEV of the charge operator $\mathcal{Q}(x_0;\tilde{t}_i)$ for a real scalar field Φ is zero, since particle and anti-particle contributions cancel. Therefore, we need to devise a method by which to separate the particle from the anti-particle degrees of freedom.

For the general case of a spatially and temporally inhomogeneous background, we need therefore to generalize the Noether charge operator $\mathcal{Q}(x_0;\tilde{t}_i)$ by writing it in terms of a charge density operator $\mathcal{Q}(\mathbf{p},\mathbf{X},X_0;\tilde{t}_i)$ as

$$Q(X_0;\tilde{t}_i) \ = \ \int \mathrm{d}^3\mathbf{X}\int\frac{\mathrm{d}^3\mathbf{p}}{(2\pi)^3}\ \mathcal{Q}(\mathbf{p},\mathbf{X},X_0;\tilde{t}_i)\ . \tag{7.2}$$

In the above, the three-momentum $\mathbf{p}$ is conjugate to the spatial part of the relative space-time coordinate $R^\mu = x^\mu - y^\mu$ and $X^\mu = (X^0,\mathbf{X}) = (x^\mu + y^\mu)/2$ is the central space-time coordinate [cf. (3.2)]. To this end, we proceed by inserting into (7.1) unity in the following form:

$$1 \ = \ \int \mathrm{d}^4y\ \delta^{(4)}(x\ -\ y) \ = \ \int\mathrm{d}^4y\int\frac{\mathrm{d}^3\mathbf{p}}{(2\pi)^3}\ e^{-i\mathbf{p}\cdot(\mathbf{x}\,-\,\mathbf{y})}\delta(x_0\ -\ y_0)\ . \tag{7.3}$$

Observe that in what follows, $x^0 = y^0 = X^0$ thanks to $\delta(x_0 - y_0)$. Subsequently, symmetrizing the integrand in x and y, we may write the charge operator as

$$\begin{aligned} \mathcal{Q}(X_0;\tilde{t}_i) \ &= \ \frac{i}{2}\int \mathrm{d}^3\mathbf{x} \int \mathrm{d}^4 y \int \frac{\mathrm{d}^3\mathbf{p}}{(2\pi)^3}\, e^{-i\mathbf{p}\cdot(\mathbf{x}-\mathbf{y})}\delta(x_0 - y_0) \\ &\quad \times \Big(\Phi^\dagger_{\mathrm{H}}(y;\tilde{t}_i)\pi^\dagger_{\mathrm{H}}(x;\tilde{t}_i) - \pi_{\mathrm{H}}(y;\tilde{t}_i)\Phi_{\mathrm{H}}(x;\tilde{t}_i) + (x \leftrightarrow y)\Big)\, . \end{aligned} \tag{7.4}$$

In terms of the central and relative coordinates, X^μ and R^μ, the charge density operator $\mathcal{Q}(\mathbf{p},\mathbf{X},X_0;\tilde{t}_i)$ may be appropriately identified from (7.4) as

$$\begin{aligned} \mathcal{Q}(\mathbf{p},\mathbf{X},X_0;\tilde{t}_i) \ &= \ \frac{i}{2}\int \mathrm{d}^4R\; e^{-i\mathbf{p}\cdot\mathbf{R}}\,\delta(R_0) \\ &\times \Big(\Phi^\dagger_{\mathrm{H}}(X - \tfrac{R}{2};\tilde{t}_i)\pi^\dagger_{\mathrm{H}}(X + \tfrac{R}{2};\tilde{t}_i) - \pi_{\mathrm{H}}(X - \tfrac{R}{2};\tilde{t}_i)\Phi_{\mathrm{H}}(X + \tfrac{R}{2};\tilde{t}_i) + (R \leftrightarrow -R)\Big)\, . \end{aligned} \tag{7.5}$$

Substituting for the definitions of the conjugate-momentum operators π_{H} and $\pi^\dagger_{\mathrm{H}}$, we may rewrite $\mathcal{Q}(\mathbf{p},\mathbf{X},X_0;\tilde{t}_i)$ in the following form:

$$\begin{aligned} \mathcal{Q}(\mathbf{p},\mathbf{X},X_0;\tilde{t}_i) \ &= \ i\int \mathrm{d}^4R\; e^{-i\mathbf{p}\cdot\mathbf{R}}\,\delta(R_0) \\ &\times\ \partial_{R_0}\Big(\Phi^\dagger_{\mathrm{H}}(X - \tfrac{R}{2};\tilde{t}_i)\Phi_{\mathrm{H}}(X + \tfrac{R}{2};\tilde{t}_i) - \Phi^\dagger_{\mathrm{H}}(X + \tfrac{R}{2};\tilde{t}_i)\Phi_{\mathrm{H}}(X - \tfrac{R}{2};\tilde{t}_i)\Big)\, . \end{aligned} \tag{7.6}$$

The EEV of $\mathcal{Q}(\mathbf{p},\mathbf{X},X_0;\tilde{t}_i)$ at the macroscopic time t is then obtained by taking the trace with the density operator ρ as given in (4.3) in the equal-time limit $X_0 = \tilde{t}_f$. We have seen in Section 3 that the equal-time limit is necessary to ensure that the observable charge density is picture-independent and that the number of independent coordinates is reduced to seven as required. Thus, we have

$$\begin{aligned} \langle \mathcal{Q}(\mathbf{p},\mathbf{X},\tilde{t}_f;\tilde{t}_i)\rangle_t \ &= \ \lim_{X_0 \to \tilde{t}_f}\ i\int \mathrm{d}^4R\; e^{-i\mathbf{p}\cdot\mathbf{R}}\,\delta(R_0) \\ &\qquad \times\ \partial_{R_0}\Big(i\Delta_<(R,X,\tilde{t}_f;\tilde{t}_i) - i\Delta_<(-R,X,\tilde{t}_f;\tilde{t}_i)\Big)\, , \end{aligned} \tag{7.7}$$

where we use the notation

$$i\Delta_<(R,X,\tilde{t}_f;\tilde{t}_i) \ = \ \langle \Phi^\dagger_{\mathrm{H}}(X - \tfrac{R}{2};\tilde{t}_i)\Phi_{\mathrm{H}}(X + \tfrac{R}{2};\tilde{t}_i)\rangle_t \tag{7.8}$$

for the resummed CTP Wightman propagator.

Let us comment on the two terms $i\Delta_<(R,X,\tilde{t}_f;\tilde{t}_i)$ and $i\Delta_<(-R,X,\tilde{t}_f;\tilde{t}_i)$ that occur on the RHS of (7.7). The first term $i\Delta_<(R,X,\tilde{t}_f;\tilde{t}_i)$ comprises *ensemble* positive-frequency particle modes and *ensemble plus vacuum* negative-frequency anti-particle modes. The second one $i\Delta_<(-R,X,\tilde{t}_f;\tilde{t}_i)$ comprises *ensemble plus vacuum* positive-frequency anti-particle modes and *ensemble* negative-frequency particle modes. Hence, we may extract the number density of particles by taking the sum of the positive-frequency contribution from $i\Delta_<(R,X,\tilde{t}_f;\tilde{t}_i)$ and the negative-frequency contribution from $i\Delta_<(-R,X,\tilde{t}_f;\tilde{t}_i)$.

We separate the positive- and negative-frequency parts of (7.7) by decomposing the equal-time delta function $\delta(R_0)$ via the limit representation

$$\delta(R_0) \ = \ \frac{i}{2\pi}\bigg(\frac{1}{R_0 + i\epsilon} - \frac{1}{R_0 - i\epsilon}\bigg)\, , \tag{7.9}$$

with $\epsilon = 0^+$. Thus, a physically meaningful definition of the number density of particles at the macroscopic time t is given by

$$
\begin{aligned}
n(\mathbf{p}, \mathbf{X}, \tilde{t}_f; \tilde{t}_i) \; &= \; - \lim_{X_0 \to \tilde{t}_f} \int \mathrm{d}^3\mathbf{R}\, e^{-i\mathbf{p}\cdot\mathbf{R}} \int \frac{\mathrm{d}R_0}{2\pi} \\
&\times \left(\frac{1}{R_0 + i\epsilon}\, \partial_{R_0} i\Delta_<(R, X, \tilde{t}_f; \tilde{t}_i) \; + \; \frac{1}{R_0 - i\epsilon}\, \partial_{R_0} i\Delta_<(-R, X, \tilde{t}_f; \tilde{t}_i) \right) . \quad (7.10)
\end{aligned}
$$

Using time translational invariance of the CTP contour, this observable may be recast in terms of the Wigner transform of the Wightman propagators as

$$
\begin{aligned}
n(\mathbf{p}, \mathbf{X}, t) \; &\equiv \; n(\mathbf{p}, \mathbf{X}, t; 0) \\
&= \; \lim_{X_0 \to t} \int \frac{\mathrm{d}p_0}{2\pi}\, p_0 \Big(\theta(p_0) i\Delta_<(p, X, t; 0) \; - \; \theta(-p_0) i\Delta_<(-p, X, t; 0) \Big) . \quad (7.11)
\end{aligned}
$$

Note that the number density of anti-particles $n^C(\mathbf{p}, \mathbf{X}, t)$ is obtained by C-conjugating the two negative-frequency Wightman propagators in (7.11). Useful relations between correlation functions and their C-conjugated counterparts are given in Appendices A and C.

We reiterate from (4.56) that the $n(\mathbf{p}, \mathbf{X}, t)\, \mathrm{d}^3\mathbf{p}\, \mathrm{d}^3\mathbf{X}$ is interpreted as the number of particles at macroscopic time t in the volume of phase-space between $\mathbf{p}$ and $\mathbf{p} + \mathrm{d}^3\mathbf{p}$ and $\mathbf{X}$ and $\mathbf{X} + \mathrm{d}^3\mathbf{X}$. The particle number per unit volume is obtained by integrating over all momentum modes, i.e.

$$
n(\mathbf{X}, t) \; = \; \int \frac{\mathrm{d}^3\mathbf{p}}{(2\pi)^3}\, n(\mathbf{p}, \mathbf{X}, t) \quad (7.12)
$$

and finally the total particle number, by integrating over all space, i.e.

$$
N(t) \; = \; \int \mathrm{d}^3\mathbf{X}\, n(\mathbf{X}, t) \, . \quad (7.13)
$$

By inserting the inverse Wigner transform

$$
i\Delta_<(p, X, t; 0) \; = \; \int \frac{\mathrm{d}^4P}{(2\pi)^4}\, e^{-iP\cdot X}\, i\Delta_<(p + \tfrac{P}{2}, p - \tfrac{P}{2}, t; 0) \, , \quad (7.14)
$$

into (7.11), the particle number per unit volume $n(\mathbf{X}, t)$ may be expressed in terms of the double momentum representation of the Wightman propagators via (7.12). After making the coordinate transformation $p \to -p$ in the negative-frequency contribution, we then obtain

$$
n(\mathbf{X}, t) \; = \; \lim_{X_0 \to t} 2 \iint \frac{\mathrm{d}^4p}{(2\pi)^4} \frac{\mathrm{d}^4P}{(2\pi)^4}\, e^{-iP\cdot X}\, \theta(p_0) p_0 i\Delta_<(p + \tfrac{P}{2}, p - \tfrac{P}{2}, t; 0) \, , \quad (7.15)
$$

The particle number per unit volume $n(\mathbf{X}, t)$ is also related to the statistical distribution function f, through

$$
n(\mathbf{X}, t) \; = \; \int \frac{\mathrm{d}^3\mathbf{p}}{(2\pi)^3} \int \frac{\mathrm{d}^3\mathbf{P}}{(2\pi)^3}\, e^{i\mathbf{P}\cdot\mathbf{X}}\, f(\mathbf{p} + \mathbf{P}/2, \mathbf{p} - \mathbf{P}/2, t) \, , \quad (7.16)
$$

using (4.56). Equation (7.16) relies on our earlier observation that the statistical distribution function f appearing in the non-homogeneous free propagators of Section 4.3 is a dressed object, comprising all possible multi-particle background contributions.

It is instructive to check that our definitions for the number density lead to the expected results for the free and quasi-particle equilibrium cases. Substituting the free equilibrium Wightman propagator (5.10c) of the real scalar field into (7.11), we obtain

$$n_{\mathrm{eq}}(\mathbf{p},\mathbf{X},t) \;=\; f_{\mathrm{B}}\big(E(\mathbf{p})\big)\;, \tag{7.17}$$

exactly as we would expect for the number density of free particles in thermodynamic equilibrium. Inserting instead the resummed equilibrium Wightman propagator given in (5.23) in the narrow width limit, we get

$$n_{\mathrm{qp}}(\mathbf{p},\mathbf{X},t) \;=\; f_{\mathrm{B}}\big(\mathcal{E}(\mathbf{p})\big)\;, \tag{7.18}$$

where $\mathcal{E}(\mathbf{p})$ is the solution to the gap equation,

$$\mathcal{E}^2(\mathbf{p}) \;=\; \mathbf{p}^2 \;+\; M^2 \;-\; \mathrm{Re}\,\Pi_{\mathrm{R}}(\mathcal{E}(\mathbf{p})+i\epsilon,\mathbf{p})\;, \tag{7.19}$$

and $n_{\mathrm{qp}}(\mathbf{p},\mathbf{X},t)$ then represents the number density of quasi-particles.

8 Master Time Evolution Equations for Particle Number Densities

Having established a direct relationship between the non-homogeneous CTP propagators and the particle number density in Section 7, we are now in a position to derive in this section master time evolution equations for the particle number density $n(\mathbf{X},t)$ and the statistical distribution function f. This is achieved in analogy to the derivation of the well-known Kadanoff–Baym equations [52, 53] by partially inverting the CTP Schwinger–Dyson equation obtained in Section 4.1. Our approach, however, differs significantly from other methods. As discussed in Section 4.2, we do not rely on a truncation of a gradient expansion of the resulting expressions. More details of the gradient expansion are given in Appendix E. In the next section, we will employ a loopwise truncation of the time evolution equations in terms of non-homogeneous free CTP propagators. As we will see, these dynamical equations are nonetheless resummed *to all orders* in a gradient expansion.

We begin our derivation of the time evolution equations with the double momentum representation of the Schwinger–Dyson equation in (4.32). We convolute (4.32) consecutively from the right with the weight function $(2\pi)^4\delta_t^{(4)}(q_1\,-\,q_2)$ defined in (3.49) and then with the resummed CTP propagator $\Delta^{ab}(q_2,p_2,\tilde{t}_f;\tilde{t}_i)$. By making use of the inverse relation (3.48), we obtain the following expression:

$$\begin{aligned}\iint\!\frac{\mathrm{d}^4q_1}{(2\pi)^4}\,\frac{\mathrm{d}^4q_2}{(2\pi)^4}\,\Delta^{0,\,-1}_{ac}(p_1,q_1)(2\pi)^4\delta_t^{(4)}(q_1\,-\,q_2)\Delta^{c}_{\ b}(q_2,p_2,\tilde{t}_f;\tilde{t}_i) \;&=\; \eta_{ab}(2\pi)^4\delta_t^{(4)}(p_1\,-\,p_2)\\ -\iint\!\frac{\mathrm{d}^4q_1}{(2\pi)^4}\,\frac{\mathrm{d}^4q_2}{(2\pi)^4}\,\Pi_{ac}(p_1,q_1,\tilde{t}_f;\tilde{t}_i)(2\pi)^4\delta_t^{(4)}&(q_1\,-\,q_2)\Delta^{c}_{\ b}(q_2,p_2,\tilde{t}_f;\tilde{t}_i)\;,\end{aligned} \tag{8.1}$$

where the contribution of the bi-local source K_{ab} is neglected next to the self-energies. It is essential to remark that the LHS of (8.1) has the following coordinate-space representation:

$$\int_{\Omega_t} \mathrm{d}^4 z\ \Delta^{0,\,-1}_{ab}(x,z)\Delta^{b}_{\ c}(z,y,\tilde{t}_f;\tilde{t}_i)\ . \tag{8.2}$$

Substituting for the free inverse CTP propagator $\Delta^{0,\,-1}_{ab}(x,y)$ given in (3.21), we may then confirm via (8.2) that evaluating the q_1 and q_2 integrals on the LHS of (8.1) yields

$$\begin{aligned}\iint \frac{\mathrm{d}^4 q_1}{(2\pi)^4}\,\frac{\mathrm{d}^4 q_2}{(2\pi)^4}\ \Delta^{0,\,-1}_{ac}(p_1,q_1)(2\pi)^4\delta_t^{(4)}(q_1\ -\ q_2)\Delta^{c}_{\ b}(q_2,p_2,\tilde{t}_f;\tilde{t}_i)&\\ =\ \Big(p_1^2\ -\ M^2\Big)\Delta_{ab}(p_1,p_2,\tilde{t}_f;\tilde{t}_i)\ .&\end{aligned} \tag{8.3}$$

Recalling that the self-energy $\Pi_{ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ contains $\delta_t^{(4)}(p-p')$ functions in the vertices, we may perform the q_1 integral on the RHS of (8.1) by making use of (3.52). Consequently, (8.1) may be written down in the following concise form:

$$\begin{aligned}\Big(p_1^2\ -\ M^2\Big)\Delta_{ab}(p_1,p_2,\tilde{t}_f;\tilde{t}_i)\ &=\ \eta_{ab}(2\pi)^4\delta_t^{(4)}(p_1\ -\ p_2)\\ &\quad -\int\frac{\mathrm{d}^4 q}{(2\pi)^4}\ \Pi_{ac}(p_1,q,\tilde{t}_f;\tilde{t}_i)\Delta^{c}_{\ b}(q,p_2,\tilde{t}_f;\tilde{t}_i)\ .\end{aligned} \tag{8.4}$$

With the definition of the particle number per unit volume $n(\mathbf{X},t)$ from (7.12) in mind, we equate the element $(a,b)\ =\ (1,2)$ of each side of (8.4) to extract the interacting Klein–Gordon equation of the negative-frequency resummed Wightman propagator:

$$\begin{aligned}&\Big(p_1^2\ -\ M^2\Big)\Delta_{<}(p_1,p_2,\tilde{t}_f;\tilde{t}_i)\\ &=\ -\int\frac{\mathrm{d}^4 q}{(2\pi)^4}\ \Big(\Pi(p_1,q,\tilde{t}_f;\tilde{t}_i)\Delta_{<}(q,p_2,\tilde{t}_f;\tilde{t}_i)\ -\ \Pi_{<}(p_1,q,\tilde{t}_f;\tilde{t}_i)\Delta_{\mathrm{D}}(q,p_2,\tilde{t}_f;\tilde{t}_i)\Big)\ .\end{aligned} \tag{8.5}$$

Using the decomposition of $\Delta_{\mathrm{D}}(p,p',\tilde{t}_f;\tilde{t}_i)$ from (A.6e) and an analogous identity for $\Pi(p,p',\tilde{t}_f;\tilde{t}_i)$ analagous, we may rewrite (8.5) as

$$\begin{aligned}\Big(p_1^2\ -\ M^2\Big)\Delta_{<}(p_1,p_2,\tilde{t}_f;\tilde{t}_i)\ +\int\frac{\mathrm{d}^4 q}{(2\pi)^4}\ \Pi_{\mathcal{P}}(p_1,q,\tilde{t}_f;\tilde{t}_i)\Delta_{<}(q,p_2,\tilde{t}_f;\tilde{t}_i)&\\ =\ -\frac{1}{2}\int\frac{\mathrm{d}^4 q}{(2\pi)^4}\ \Big[\Pi_{>}(p_1,q,\tilde{t}_f;\tilde{t}_i)\Delta_{<}(q,p_2,\tilde{t}_f;\tilde{t}_i)&\\ -\ \Pi_{<}(p_1,q,\tilde{t}_f;\tilde{t}_i)\Big(\Delta_{>}(q,p_2,\tilde{t}_f;\tilde{t}_i)\ -\ 2\Delta_{\mathcal{P}}(q,p_2,\tilde{t}_f;\tilde{t}_i)\Big)\Big]\ ,&\end{aligned} \tag{8.6}$$

where the subscript $\mathcal{P}$ denotes principal-part evaluation of the functions given in (2.29) and (4.27b).

Introducing the central and relative momenta, $p\ =\ (p_1\ +\ p_2)/2$ and $P\ =\ p_1\ -\ p_2$, respectively, we write (8.6) in the following form:

$$\begin{aligned}\Big[\big(p_0\ +\ \tfrac{P_0}{2}\big)^2\ -\ E^2\big(\mathbf{p}\ +\ \tfrac{\mathbf{P}}{2}\big)\Big]\Delta_{<}(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)&\\ +\ \mathscr{F}(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\ =\ \mathscr{C}(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\ ,&\end{aligned} \tag{8.7}$$

where we have defined the following:

$$\begin{aligned}
\mathscr{F}(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i) \;&\equiv\; -\!\int\!\frac{\mathrm{d}^4q}{(2\pi)^4}\; i\Pi_{\mathcal{P}}(p+\tfrac{P}{2},q,\tilde{t}_f;\tilde{t}_i)\; i\Delta_<(q,p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i), && \text{(8.8a)}\\
\mathscr{C}(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i) \;&\equiv\; \frac{1}{2}\!\int\!\frac{\mathrm{d}^4q}{(2\pi)^4}\;\Big[\,i\Pi_>(p+\tfrac{P}{2},q,\tilde{t}_f;\tilde{t}_i)\; i\Delta_<(q,p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\\
&\quad -\; i\Pi_<(p+\tfrac{P}{2},q,\tilde{t}_f;\tilde{t}_i)\,\Big(\,i\Delta_>(q,p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\;-\;2i\Delta_{\mathcal{P}}(q,p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\,\Big)\Big]\;. && \text{(8.8b)}
\end{aligned}$$

With the aim of finding a master time evolution equation for the particle number density $n(\mathbf{X},t)$ in (7.12), we integrate both sides of (8.7) with the measure

$$\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;e^{-iP\cdot X}\,\theta(p_0)\;. \tag{8.9}$$

Explicitly, this gives

$$\begin{aligned}
&\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;e^{-iP\cdot X}\,\theta(p_0)\Big[\,\big(p_0\;+\;\tfrac{P_0}{2}\big)^2\;-\;E^2(\mathbf{p}+\tfrac{\mathbf{P}}{2})\Big]\Delta_<(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\\
&\qquad+\;\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;e^{-iP\cdot X}\,\theta(p_0)\,\mathscr{F}(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\\
&\qquad\qquad=\;\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;e^{-iP\cdot X}\,\theta(p_0)\,\mathscr{C}(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\;.
\end{aligned}\tag{8.10}$$

Adding to (8.10) the complex conjugate of the same expression with $P\to -P$ and using the identities in (A.4), we may extract the terms proportional to $p\cdot P$ on the LHS of (8.10). In this way, we obtain

$$\begin{aligned}
&2\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;e^{-iP\cdot X}\,\theta(p_0)\,p\cdot P\,\Delta_<(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\\
&\quad+\;\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;e^{-iP\cdot X}\,\theta(p_0)\Big(\mathscr{F}(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\;+\;\mathscr{F}^*(p-\tfrac{P}{2},p+\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\Big)\\
&\quad=\;\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;e^{-iP\cdot X}\,\theta(p_0)\Big(\mathscr{C}(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\;+\;\mathscr{C}^*(p-\tfrac{P}{2},p+\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\Big)\;.
\end{aligned}\tag{8.11}$$

The first term on the LHS of (8.11) may be rewritten as

$$2\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;\theta(p_0)\Big(\,ip_0\partial_{X_0}\;-\;\mathbf{p}\cdot\mathbf{P}\,\Big)e^{-iP\cdot X}\,\Delta_<(p+\tfrac{P}{2},p-\tfrac{P}{2},\tilde{t}_f;\tilde{t}_i)\;. \tag{8.12}$$

Using time translational invariance of the CTP contour and taking the limit $X_0\;\to\;t$ in (8.12), we recognise that the derivative term with respect to t is precisely the time-derivative of the particle number density $n(\mathbf{X},t)$ in (7.12). Hence, from (8.11) and (8.12), we arrive at our master time evolution equation for $n(\mathbf{X},t)$:

$$\begin{aligned}
&\partial_t n(\mathbf{X},t)\;-\;\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;e^{-iP\cdot X}\,\mathbf{p}\cdot\mathbf{P}\,\theta(p_0)\Delta_<(p+\tfrac{P}{2},p-\tfrac{P}{2},t;0)\\
&\quad+\;\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;e^{-iP\cdot X}\,\theta(p_0)\Big(\mathscr{F}(p+\tfrac{P}{2},p-\tfrac{P}{2},t;0)\;+\;\mathscr{F}^*(p-\tfrac{P}{2},p+\tfrac{P}{2},t;0)\Big)\\
&\quad=\;\iint\!\frac{\mathrm{d}^4p}{(2\pi)^4}\,\frac{\mathrm{d}^4P}{(2\pi)^4}\;e^{-iP\cdot X}\theta(p_0)\Big(\mathscr{C}(p+\tfrac{P}{2},p-\tfrac{P}{2},t;0)\;+\;\mathscr{C}^*(p-\tfrac{P}{2},p+\tfrac{P}{2},t;0)\Big)\;,
\end{aligned}\tag{8.13}$$

with $X_0 = t$, where $\mathscr{F}$ and $\mathscr{C}$ are defined in (8.8a) and (8.8b). Comparing with the full non-truncated form of the Kadanoff–Baym kinetic equation in (E.4b), the series of nested Poisson brackets in (E.4b) has been replaced by a single convolution integral over the central momentum P in (8.13).

In addition to the master time evolution equation for the particle number density $n(\mathbf{X}, t)$ in (8.13), we may respectively find a time evolution equation for the statistical distribution function $f(\mathbf{p} + \frac{\mathbf{P}}{2}, \mathbf{p} - \frac{\mathbf{P}}{2}, t)$. Specifically, given the relation (7.16) and the differential equation (8.13), the following time evolution equation may be derived for f:

$$\begin{aligned}\partial_t f(\mathbf{p} + \tfrac{\mathbf{P}}{2}, \mathbf{p} - \tfrac{\mathbf{P}}{2}, t) \;-\; \iint \frac{\mathrm{d}p_0}{2\pi}\,\frac{\mathrm{d}P_0}{2\pi}\; e^{-iP_0 t}\,\mathbf{p}\cdot\mathbf{P}\,\theta(p_0)\Delta_<(p + \tfrac{P}{2}, p - \tfrac{P}{2}, t; 0) \\ +\; \iint \frac{\mathrm{d}p_0}{2\pi}\,\frac{\mathrm{d}P_0}{2\pi}\; e^{-iP_0 t}\,\theta(p_0)\Big(\mathscr{F}(p + \tfrac{P}{2}, p - \tfrac{P}{2}, t; 0) \;+\; \mathscr{F}^*(p - \tfrac{P}{2}, p + \tfrac{P}{2}, t; 0)\Big) \\ =\; \iint \frac{\mathrm{d}p_0}{2\pi}\,\frac{\mathrm{d}P_0}{2\pi}\; e^{-iP_0 t}\,\theta(p_0)\Big(\mathscr{C}(p + \tfrac{P}{2}, p - \tfrac{P}{2}, t; 0) \;+\; \mathscr{C}^*(p - \tfrac{P}{2}, p + \tfrac{P}{2}, t; 0)\Big)\,.\end{aligned} \tag{8.14}$$

It is important to stress here that (8.14) provides a self-consistent time evolution equation for f valid *to all orders* in perturbation theory and to *all orders* in gradient expansion. Nevertheless, it is a formidable task to provide an evaluation of (8.14) to all orders in perturbation theory. Therefore, in the next section we develop a systematic perturbative expansion to non-equilibrium dynamics based on the result (8.14) to all orders in gradients.

We note that a physical intepretation may be attributed to the different terms contributing to (8.14). Specifically, all terms on the LHS of (8.14) may be associated with the total derivative in the phase-space $(\mathbf{X},\ \mathbf{p})$, appearing in the classical Boltzmann transport equation [75]:

$$D_t \;=\; \partial_t \;+\; \mathbf{v}\cdot\nabla_{\mathbf{X}} \;+\; \mathbf{F}\cdot\nabla_{\mathbf{p}}\,, \tag{8.15}$$

where $\mathbf{v}$ is the average non-relativistic velocity of the particle distribution and $\mathbf{F}$ is the force acting on this distribution. In particular, the $\mathscr{F}$ terms on the LHS of (8.14) are the *force* terms, generated by the potential due to the dispersive part of the self-energy. On the RHS of (8.14), the $\mathscr{C}$ terms represent the *collision* terms.

It would be interesting to discuss the spatially homogeneous limit of (8.14) at late times. In this case, energy conservation holds to a good approximation, so the time-dependent weight functions δ_t may be replaced by the standard Dirac delta functions, even in the vertices of the self-energies contained in the force $\mathscr{F}$ and collision $\mathscr{C}$ terms of (8.14). Moreover, as a consequence of the assumed spatial homogeneity, all propagators and self-energies, which in general depend on two momenta p_1 and p_2, will now be proportional to the four-dimensional delta function $(2\pi)^4\delta^{(4)}(p_1 - p_2)$. Likewise, the statistical distribution function f takes on the form

$$f(\mathbf{p} + \tfrac{\mathbf{P}}{2}, \mathbf{p} - \tfrac{\mathbf{P}}{2}, t) \;=\; (2\pi)^3\delta^{(3)}(\mathbf{P})f(|\mathbf{p}|, t)\,. \tag{8.16}$$

Because of the above simplifications, one can then work in the single momentum representation, by integrating over the three-momentum $\mathbf{P}$. Thus, we find the following time evolution equation for $f(|\mathbf{p}|, t)$:

$$\partial_t f(|\mathbf{p}|, t) \;=\; \int \frac{\mathrm{d}p_0}{2\pi}\,\theta(p_0)\Big(i\Pi_>(p, t)i\Delta_<(p, t) \;-\; i\Pi_<(p, t)i\Delta_>(p, t)\Big)\,, \tag{8.17}$$

where the purely-imaginary force term $\mathscr{F}$ and off-shell effects from $\Delta_{\mathcal{P}}$ have vanished for $P^\mu = (0, \mathbf{0})$. This result corresponds to the semi-classical Boltzmann transport equation in (6.18) or, equivalently, to the zeroth-order truncation of the gradient-expanded Kadanoff–Baym kinetic equation in (E.5b) with $X_0 = t$.

9 Thermalization in a Scalar Model

We now apply the formalism developed in the proceeding sections to a simple scalar model. In particular, we introduce the modified Feynman rules that result from the systematic inclusion of finite-time effects and the violation of both energy conservation and space-time translational invariance. As a playground for studying the kinematics in the early-time energy-non-conserving regime, we calculate the time-dependent thermal width of a heavy scalar Φ. We show that processes, which would normally be kinematically disallowed, contribute significantly to the prompt *shock regime* of the initial evolution. We also show that the subsequent dynamics exhibit non-Markovian behaviour, acquiring oscillations with time-dependent frequencies. This evolution signifies the occurence of memory effects, as are expected in truly out-of-equilibrium systems. Finally, we look in more detail at the time evolution equations of this simple model and demonstrate the importance of the violation of energy conservation to the statistical dynamics.

We consider a simple scalar theory, which comprises one heavy real scalar field Φ and one light pair of complex scalar fields $(\chi^\dagger,\, \chi)$, described by the Lagrangian

$$\begin{aligned}\mathcal{L}(x) \;=\; & \tfrac{1}{2}\partial_\mu\Phi(x)\partial^\mu\Phi(x) \;-\; \tfrac{1}{2}M^2\Phi^2(x) \;+\; \partial_\mu\chi^\dagger(x)\partial^\mu\chi(x) \;-\; m^2\chi^\dagger(x)\chi(x)\\ & -\; g\Phi(x)\chi^\dagger(x)\chi(x) \;-\; \tfrac{1}{4}\lambda\big[\chi^\dagger(x)\chi(x)\big]^2\,, \end{aligned} \tag{9.1}$$

where $M \gg m$. Appendix C describes the generalization of our approach to the complex scalar field χ.

We formulate a perturbative approach based upon the following *modified* Feynman rules:

- sum over all topologically distinct diagrams at a given order in perturbation theory.

- assign to each Φ-propagator line a factor of



$$= \; i\Delta_{\Phi}^{0,\,ab}(p, p', \tilde{t}_f; \tilde{t}_i)\;.$$

- assign to each χ-propagator line a factor of



$$= \; i\Delta_{\chi}^{0,\,ab}(p, p', \tilde{t}_f; \tilde{t}_i)\;.$$

- assign to each three-point Φ vertex a factor of



$$= -ig\eta_{abc}\,(2\pi)^4\delta_t\big(\textstyle\sum_{i=1}^{3} p_{0,i}\big)\;\delta^{(3)}\big(\textstyle\sum_{i=1}^{3}\mathbf{p}_i\big)\ ,$$

  where δ_t is defined in (3.50) and the pre-factors $\eta_{abc\ldots}$ are given after (3.12).

- assign to each four-point χ vertex a factor of



$$= -i\lambda\eta_{abcd}\,(2\pi)^4\delta_t\big(\textstyle\sum_{i=1}^{4} p_{0,i}\big)\;\delta^{(3)}\big(\textstyle\sum_{i=1}^{4}\mathbf{p}_i\big)\ .$$

- associate with each external vertex a phase

$$e^{ip_0\tilde{t}_f}\ ,$$

  where p_0 is the energy flowing *into* the vertex.

- contract all internal CTP indices.

- integrate with the measure

$$\int\frac{\mathrm{d}^4p}{(2\pi)^4}$$

  over the four-momentum associated with each contracted pair of CTP indices.

- consider the combinatorial symmetry factors, where appropriate.

Notice that there are a number of modifications with respect to the standard Feynman rules. In particular, the familiar energy-conserving delta function has been replaced by δ_t in the vertices. This is indicated diagrammatically by the dotted line terminated in a cross, representing the violation of energy conservation. This loss of energy conservation is a consequence of Heisenberg's uncertainty principle, due to the finite macroscopic time of observation t, over which the interactions have been switched on. The time-dependent vertices vanish in the limit $t \to 0$, as we should expect. The loss of space-time translational invariance leads to a doubling of the number of integrations with respect to the zero-temperature and equilibrium cases.



Figure 4: One-loop local χ self-energy: $i\Pi^{\mathrm{loc}(1)}_{\chi,\,ab}(q,q',\tilde{t}_f;\tilde{t}_i) \;\propto\; \eta_{ab}$.

At the one-loop level, we have three diagrams. The local χ self-energy shown in Figure 4:

$$
\begin{aligned}
i\Pi^{\mathrm{loc}(1)}_{\chi,\,ab}(q,q',\tilde{t}_f;\tilde{t}_i) \;&=\; \frac{-i\lambda}{2!}(2\pi\mu)^{2\epsilon}e^{i(q_0\,-\,q'_0)\tilde{t}_f}\iint\frac{\mathrm{d}^dk}{(2\pi)^d}\,\frac{\mathrm{d}^4k'}{(2\pi)^4}\\
&\quad\times\;(2\pi)^4\delta^{(4)}_t(q\,-\,q'\,-\,k\,+\,k')\eta_{abcd}i\Delta^{0,\,cd}_{\chi}(k,k',\tilde{t}_f;\tilde{t}_i)\;, \qquad (9.2)
\end{aligned}
$$

and the two non-local diagrams for the Φ and χ self-energies shown in Figure 5:

$$
\begin{aligned}
i\Pi^{(1)}_{\Phi,\,ab}(q,q',\tilde{t}_f;\tilde{t}_i) \;&=\; \frac{(-ig)^2}{2!}(2\pi\mu)^{2\epsilon}e^{i(q_0\,-\,q'_0)\tilde{t}_f}\int\cdots\int\frac{\mathrm{d}^dk_1}{(2\pi)^d}\,\frac{\mathrm{d}^4k'_1}{(2\pi)^4}\,\frac{\mathrm{d}^4k_2}{(2\pi)^4}\,\frac{\mathrm{d}^4k'_2}{(2\pi)^4}\\
&\quad\times\;(2\pi)^4\delta^{(4)}_t(q\,-\,k_1\,-\,k_2)(2\pi)^4\delta^{(4)}_t(q'\,-\,k'_1\,-\,k'_2)\\
&\quad\times\;\eta_{acd}i\Delta^{0,\,ce}_{\chi}(k_1,k'_1,\tilde{t}_f;\tilde{t}_i)i\Delta^{C,\,0,\,df}_{\chi}(k_2,k'_2,\tilde{t}_f;\tilde{t}_i)\eta_{efb}\;, \qquad (9.3a)\\
i\Pi^{(1)}_{\chi,\,ab}(q,q',\tilde{t}_f;\tilde{t}_i) \;&=\; \frac{(-ig)^2}{2!}(2\pi\mu)^{2\epsilon}e^{i(q_0\,-\,q'_0)\tilde{t}_f}\int\cdots\int\frac{\mathrm{d}^dk_1}{(2\pi)^d}\,\frac{\mathrm{d}^4k'_1}{(2\pi)^4}\,\frac{\mathrm{d}^4k_2}{(2\pi)^4}\,\frac{\mathrm{d}^4k'_2}{(2\pi)^4}\\
&\quad\times\;(2\pi)^4\delta^{(4)}_t(q\,-\,k_1\,-\,k_2)(2\pi)^4\delta^{(4)}_t(q'\,-\,k'_1\,-\,k'_2)\\
&\quad\times\;\eta_{acd}i\Delta^{0,\,ce}_{\chi}(k_1,k'_1,\tilde{t}_f;\tilde{t}_i)i\Delta^{0,\,df}_{\Phi}(k_2,k'_2,\tilde{t}_f;\tilde{t}_i)\eta_{efb}\;. \qquad (9.3b)
\end{aligned}
$$

A detailed description of the techniques required to perform these loop integrals is provided in Appendix F.

9.1 Time-Dependent Width

In this section, we study the time-dependent width of the heavy scalar Φ. In particular, we investigate the spectral evolution that results from the restoration of energy conservation, without solving explicitly the system of evolution equations for the statistical dynamics.

We consider the following situation. We prepare two isolated but coincident subsystems $\mathscr{S}_\Phi$ and $\mathscr{S}_\chi$, both separately in thermodynamic equilibrium and at the same temperature T, with all interactions turned off. The subsystem $\mathscr{S}_\Phi$ contains only the real scalar field Φ, whilst $\mathscr{S}_\chi$ contains only the complex scalar χ. At macroscopic time $t\,=\,0$, we turn on the interactions and allow the system $\mathscr{S}\,=\,\mathscr{S}_\Phi\cup\mathscr{S}_\chi$ to re-thermalize. For our numerical analysis, we take for definiteness the thermodynamic temperature to be $T\,=\,10$ GeV, the mass of the heavy Φ-scalar $M\,=\,1$ GeV, the mass of the complex χ-scalar $m\,=\,0.01$ GeV, and their trilinear coupling $g\,=\,0.1$ GeV.



(a) $i\Pi^{(1)}_{\Phi,\,ab}(q,q',\tilde{t}_f;\tilde{t}_i)$ (b) $i\Pi^{(1)}_{\chi,\,ab}(q,q',\tilde{t}_f;\tilde{t}_i)$

Figure 5: Non-local one-loop Φ and χ self-energies.

The free propagators of the fields Φ and χ at time $t = 0$ are the equilibrium propagators in (5.10) and (C.25), containing the Bose–Einstein distributions at temperature T. We take the chemical potential of the complex scalar to be vanishingly small in comparison to its mass, i.e. $\mu \ll m$, such that $f_\chi(|\mathbf{p}|,0) = f^C_\chi(|\mathbf{p}|,0) = f_{\mathrm{B}}\big(E_\chi(\mathbf{p})\big)$. Without solving the system of time evolution equations, the form of the the statistical distribution functions of the Φ and χ scalars is unknown for $t \neq 0$. We assume that the heat bath of χ's is sufficiently large so as to remain unperturbed by the addition of the real scalar Φ. Specifically, we may consider the number density of χ's to remain unchanged and the free equilibrium χ propagators in (C.25) to persist for all times.

By the optical theorem, the width Γ_Φ of the scalar Φ is defined in terms of the absorptive part of the retarded Φ self-energy $\mathrm{Im}\,\Pi_{\Phi,\,\mathrm{R}}$ via

$$\Gamma_\Phi(q_1,q_2,\tilde{t}_f;\tilde{t}_i) \;=\; \frac{1}{M}\mathrm{Im}\,\Pi_{\Phi,\,\mathrm{R}}(q_1,q_2,\tilde{t}_f;\tilde{t}_i)\;, \tag{9.4}$$

where

$$\mathrm{Im}\,\Pi_{\Phi,\,\mathrm{R}}(q_1,q_2,\tilde{t}_f;\tilde{t}_i) \;=\; \frac{1}{2i}\Big(\Pi_{\Phi,\,>}(q_1,q_2,\tilde{t}_f;\tilde{t}_i) \;-\; \Pi_{\Phi,\,<}(q_1,q_2,\tilde{t}_f;\tilde{t}_i)\Big)\;. \tag{9.5}$$

At the one-loop level, the self-energies $\Pi^{(1)}_{\Phi,\,\gtrless}(q_1,q_2,\tilde{t}_f;\tilde{t}_i)$ are given by (9.3a).

Employing the relative and central momenta, $Q = q_1 - q_2$ and $q = (q_1 + q_2)/2$, and using the results of Appendix F, the Laplace transform with respect to the macroscopic time t of the one-loop Φ width is

$$\begin{aligned}\Gamma^{(1)}_\Phi(q+Q/2,q-Q/2,s) \;&=\; (2\pi)^4\frac{1}{\pi}\,\frac{s}{Q_0^2\,+\,4s^2}\,e^{iQ_0\tilde{t}_f}\,\delta^{(3)}(\mathbf{Q})\\ &\times\;\frac{g^2}{32\pi^2 M}\sum_{\{\alpha\}}\int\mathrm{d}^3\mathbf{k}\;\frac{1}{\pi}\,\frac{\alpha_1\alpha_2}{E_1E_2}\,\frac{1\,+\,f_{\mathrm{B}}\big(\alpha_1E_1\big)\,+\,f_{\mathrm{B}}\big(\alpha_2E_2\big)}{\big(q_0^2\,-\,\alpha_1E_1\,-\,\alpha_2E_2\big)^2\,+\,s^2}\;,\end{aligned} \tag{9.6}$$

where we use the short-hand notation $\{\alpha\}$ for the summation over $\alpha_1,\ \alpha_2\ =\ \pm 1$. For the sake of generality, we distinguish the χ^+ and χ^- decay products by assigning them different masses m_1 and m_2, respectively, so that

$$E_1 \ \equiv\ E_1(\mathbf{k}) \ =\ \sqrt{|\mathbf{k}|^2\ +\ m_1^2}\,, \tag{9.7a}$$

$$E_2 \ \equiv\ E_2(\mathbf{q}-\mathbf{k}) \ =\ \sqrt{|\mathbf{k}|^2\ -\ 2|\mathbf{k}||\mathbf{q}|\cos\theta\ +\ |\mathbf{q}|^2\ +\ m_2^2}\,. \tag{9.7b}$$

In our numerical analysis, however, we take $m_1\ =\ m_2\ =\ m$.

Performing the inverse Wigner transform with respect to Q of (9.6) in the equal-time limit $X_0\ =\ \tilde{t}_f$ and subsequently the inverse Laplace transform with respect to s, we obtain the time-dependent Φ width

$$\Gamma^{(1)}_{\Phi}(q,t)\ =\ \frac{g^2}{64\pi^2 M}\sum_{\{\alpha\}}\int \mathrm{d}^3\mathbf{k}\ \frac{\alpha_1\alpha_2}{E_1E_2}\ \frac{t}{\pi}\,\mathrm{sinc}\big[\big(q_0\ -\ \alpha_1E_1\ -\ \alpha_2E_2\big)t\big] \\ \times\ \Big(1\ +\ f_{\mathrm{B}}\big(\alpha_1E_1\big)\ +\ f_{\mathrm{B}}\big(\alpha_2E_2\big)\Big)\,. \tag{9.8}$$

In the limit $t\ \to\ \infty$, the sinc function on the RHS of (9.8) yields the standard energy-conserving delta function:

$$\lim_{t\,\to\,\infty}\ \frac{t}{\pi}\,\mathrm{sinc}\big[\big(q_0\ -\ \alpha_1E_1\ -\ \alpha_2E_2\big)t\big]\ =\ \delta\big(q_0\ -\ \alpha_1E_1\ -\ \alpha_2E_2\big)\ , \tag{9.9}$$

thereby recovering the known equilibrium result.

Figure 6 contains a series of contour plots of the differential one-loop Φ width $\mathrm{d}^2\Gamma^{(1)}_{\Phi}$ evaluated over the dominant region of the $|\mathbf{k}|$-θ phase space, where the four-momentum q^μ of the Φ scalar is on-shell, i.e. $q^2 = M^2$. For late times, the integrand is highly oscillatory and we expect the dominant peak of the sinc function to approach the region of phase space permitted in the limit $t\ \to\ \infty$ [cf. (9.9)], for which

$$|\mathbf{k}|\ =\ \frac{M^2|\mathbf{q}|\cos\theta\ +\ \sqrt{\big(|\mathbf{q}|^2\ +\ M^2\big)\big[M^4\ -\ 4m^2\big(M^2\ +\ |\mathbf{q}|^2\sin^2\theta\big)\big]}}{2\big(M^2\ +\ |\mathbf{q}|^2\sin^2\theta\big)}\,. \tag{9.10}$$

For early times, the admissible phase space is greatly expanded. For later times, the frequency of oscillation increases and the width of the central peak of the sinc function narrows, lying along a curve in the phase space. To proceed further, we need to develop a method for dealing with the kinematics in the absence of exact energy conservation.

9.2 Generalized Two-Body Decay Kinematics

At zero temperature and density, it is convenient to analyze the two-body decay kinematics by performing a Lorentz boost to the rest frame of the decaying particle. However, at finite temperature, the dependence on the thermodynamic temperature of the heat bath breaks the Lorentz covariance of the integral. As such, we cannot eliminate dependence upon the the three-momentum of the decaying particle by any such Lorentz boost; the dependence will reappear in the 'boosted' temperature:

$$T'\ =\ \gamma T\ , \tag{9.11}$$



Figure 6: Contour plots of $|\mathbf{k}|$ versus θ for $\mathrm{d}^2\Gamma_\Phi^{(1)}$ for discrete values of Mt, with $q^2 = M^2$ and $|\mathbf{q}| = 10\,\mathrm{GeV}$. The region of phase-space permitted in the $t \to \infty$ limit is shown by the blue dashed line, corresponding to the delta function in (9.9).

where

$$\gamma = \left(1 + \frac{|\mathbf{q}|^2}{M^2}\right)^{1/2} \tag{9.12}$$

is the usual Lorentz boost factor for the heavy scalar field Φ. As a result, we are compelled to analyze the kinematics of the two-body decay in the rest frame of the heat bath, which we define to be the frame in which the EEV of the three-momentum operator $\langle \widehat{\mathbf{P}} \rangle$ is minimized. For an isotropic heat bath, this is the frame in which $\langle \widehat{\mathbf{P}} \rangle = \mathbf{0}$, that is the *comoving* frame. In this section, we look more closely at the kinematics in the absence of energy conservation.

For this purpose, let us introduce the variable

$$u \equiv \big(q_0 - \alpha_1 E_1 - \alpha_2 E_2\big)t\ , \tag{9.13}$$

which may be interpreted in terms of energy *borrowed from* or *lent to* the heat bath. We shall hereafter refer to the variable u as the *evanescent action* of the process, since it has the correct dimensions and quantifies the extended kinematically allowed phase-space configurations. We also define the *evanescent energy*

$$q_u(t) \ \equiv \ q_0 \ - \ \frac{u}{t} \ , \tag{9.14}$$

which satisfies

$$\lim_{u/t \,\to\, 0} q_u(t) \ = \ q_0 \ . \tag{9.15}$$

With this substitution, we obtain the kinematic constraint

$$q_u(t) \ - \ \alpha_1 E_1 \ - \ \alpha_2 E_2 \ = \ 0 \ . \tag{9.16}$$

Since u can take large positive values, $q_u(t)$ is not necessarily restricted to positive values for early times, even when $q_0 \ = \ \sqrt{|\mathbf{q}|^2 \ + \ M^2} \ > \ 0$ is on-shell. Processes with $u/t \ \neq \ 0$ are referred to as *evanescent*.

In order to make the coordinate transformation $|\mathbf{k}| \ \to \ u$ in (9.8), we must solve (9.16) for the magnitude of the three-momentum $|\mathbf{k}(t)|$, which becomes implicitly time-dependent. Specifically, we find

$$\begin{aligned} |\mathbf{k}^{(b)}(t)| \ &= \ \frac{1}{2\big(q_u^2(t) \ - \ |\mathbf{q}|^2\cos^2\!\theta\big)}\Big[\,\big(q_u^2(t) \ - \ |\mathbf{q}|^2 \ + \ m_1^2 \ - \ m_2^2\big)|\mathbf{q}|\cos\theta \\ &\quad + \ b\,q_u(t)\sqrt{\lambda\big(q_u^2(t) \ - \ |\mathbf{q}|^2, m_1^2, m_2^2\big) \ - \ 4m_1^2|\mathbf{q}|^2\sin^2\!\theta}\ \Big] \ , \end{aligned} \tag{9.17}$$

with $b \ = \ \pm 1$ and

$$\lambda(x,y,z) \ = \ (x-y-z)^2 \ - \ 4yz \ . \tag{9.18}$$

After a little algebra, we obtain the energies

$$\begin{aligned} E_1^{(b)}(t) \ &= \ \alpha_1\frac{1}{2\big(q_u^2(t) \ - \ |\mathbf{q}|^2\cos^2\!\theta\big)}\Big[\,\big(q_u^2(t) \ - \ |\mathbf{q}|^2 \ + \ m_1^2 \ - \ m_2^2\big)q_u(t) \\ &\quad + \ b\,|\mathbf{q}|\cos\theta\sqrt{\lambda\big(q_u^2(t) \ - \ |\mathbf{q}|^2, m_1^2, m_2^2\big) \ - \ 4m_1^2|\mathbf{q}|^2\sin^2\!\theta}\ \Big] \ , \end{aligned} \tag{9.19}$$

$$\begin{aligned} E_2^{(b)}(t) \ &= \ \alpha_2\frac{1}{2\big(q_u^2(t) \ - \ |\mathbf{q}|^2\cos^2\!\theta\big)}\Big[\,\big(q_u^2(t) \ - \ |\mathbf{q}|^2\cos 2\theta \ - \ m_1^2 \ + \ m_2^2\big)q_u(t) \\ &\quad - \ b\,|\mathbf{q}|\cos\theta\sqrt{\lambda\big(q_u^2(t) \ - \ |\mathbf{q}|^2, m_1^2, m_2^2\big) \ - \ 4m_1^2|\mathbf{q}|^2\sin^2\!\theta}\ \Big] \ , \end{aligned} \tag{9.20}$$

where the overall factors of α_1 and α_2 are necessary to satisfy the initial constraint (9.16). It is clear that these results collapse to the kinematics of equilibrium field theory in the limit $u/t \ \to \ 0$.

	$\alpha_1 = \alpha_2 = +1$	$\alpha_1 = -\alpha_2$	$\alpha_1 = \alpha_2 = -1$
$u_+(t)$	$\omega_0(q,\alpha_1,\alpha_2)t$	$\omega_0(q,\alpha_1,\alpha_2)t$	$+\infty$
$u_-(t)$	$-\infty$	0	$\omega_0(q,\alpha_1,\alpha_2)t$

Table 3: Limits of integration of the evanescent action u for each of the four processes contributing to the non-equilibrium thermal Φ width for $t > 0$, where the angular frequency $\omega_0(q,\alpha_1,\alpha_2)$ is defined in (9.23).

Keeping m_1 and m_2 distinct, the Φ width for $t > 0$ is given by

$$\begin{aligned}\Gamma_\Phi^{(1)}(q,t) \;=\; & \frac{g^2}{64\pi^2 M}\sum_{\{\alpha\},\,b\,=\,\pm 1}\int_0^\pi \mathrm{d}\theta\int_{u_-(t)}^{u_+(t)}\mathrm{d}u\;\mathrm{sinc}(u)\,\frac{\sin\theta}{\big(q_u^2(t)\,-\,|\mathbf{q}|^2\cos^2\!\theta\big)^2}\\ & \times\,\Big[\lambda\big(q_u^2(t)\,-\,|\mathbf{q}|^2,m_1^2,m_2^2\big)\,-\,4m_1^2|\mathbf{q}|^2\sin^2\!\theta\Big]^{-1/2}\\ & \times\,\Big[\big(q_u^2(t)\,-\,|\mathbf{q}|^2\,+\,m_1^2\,-\,m_2^2\big)|\mathbf{q}|\cos\theta\\ & +\,b\,q_u(t)\sqrt{\lambda\big(q_u^2(t)\,-\,|\mathbf{q}|^2,m_1^2,m_2^2\big)\,-\,4m_1^2|\mathbf{q}|^2\sin^2\!\theta}\,\Big]^2\\ & \times\,\alpha_1\alpha_2\Big(1\,+\,f_{\mathrm{B}}\big(\alpha_1E_1^{(b)}(t)\big)\,+\,f_{\mathrm{B}}\big(\alpha_2E_2^{(b)}(t)\big)\Big)\,. \end{aligned} \tag{9.21}$$

The reality of the loop momentum $|\mathbf{k}(t)|$ in (9.17) requires that the discriminant

$$\lambda\big(q_u^2(t)\,-\,|\mathbf{q}|^2,m_1^2,m_2^2\big)\,-\,4m_1^2|\mathbf{q}|^2\sin^2\!\theta\;\geq\;0\,, \tag{9.22}$$

which is now a *time-dependent* kinematic constraint. Furthermore, we require $|\mathbf{k}^{(b)}(t)| \geq 0$, $E_1^{(b)}(t) \geq m_1$ and $E_2^{(b)}(t) \geq m_2$. For $t > 0$, the limits of integration $u_\pm(t)$ are given in Table 3, where we have defined

$$\omega_0(q,\alpha_1,\alpha_2)\;\equiv\;q_0\,-\,\big(\alpha_1 m_1\,+\,\alpha_2 m_2\big)\left(1\,+\,\frac{|\mathbf{q}|^2}{\big(m_1\,+\,m_2\big)^2}\right)^{1/2}\,, \tag{9.23}$$

which is the angular frequency of the sine-integral-like oscillations of the integral. For $t = 0$, $u_\pm(t) = 0$ and the domain of integration over u collapses to zero. Given the analytic behaviour of the integrand in the limit $t \to 0$, the integral vanishes, as we expect.

For on-shell decay modes with $q^2 = M^2$, the angular frequency $\omega_0(q,\alpha_1,\alpha_2)$ in (9.23) becomes

$$\omega_0(|\mathbf{q}|,\alpha_1,\alpha_2)\;\equiv\;\omega_0(q,\alpha_1,\alpha_2)\big|_{q^2\,=\,M^2}\;=\;\sqrt{|\mathbf{q}|^2\,+\,M^2}-\sqrt{|\mathbf{q}|^2\,+\,\big(m_1\,+\,m_2\big)^2}\,. \tag{9.24}$$

Thus, the evolution of the phase space for on-threshold decays with $M^2 = (m_1 + m_2)^2$ is *critically damped*. We note that in the large momentum limit $|\mathbf{q}| \gg M$,

$$\omega_0(|\mathbf{q}|,\alpha_1,\alpha_2)\big|_{|\mathbf{q}|\,\gg\,M}\;\simeq\;\frac{M^2\,-\,\big(m_1\,+\,m_2\big)^2}{2|\mathbf{q}|}\,, \tag{9.25}$$



Figure 7: The dependence of $\omega_0(|\mathbf{q}|, \alpha_1, \alpha_2)/M$ on $|\mathbf{q}|/M$ for processes related to 1 → 2 decay (solid black), Landau damping (blue dashed) and 3 → 0 total annihilation (red dotted), where $q^2 = M^2$ and $m_1 = m_2 = m = 0.01\,\text{GeV}$. The latter two of these processes become highly oscillatory for large $|\mathbf{q}|/M$.



(a) 1 → 2 decay

(b) 2 → 1 Landau damping

(c) 3 → 0 total annihilation

Figure 8: Processes contributing to the time-dependent Φ width: (a) the familiar 1 → 2 decay; (b) 2 → 1 Landau damping; and (c) 3 → 0 total annihilation.

such that the evolution of the phase space for high-momentum modes is similarly damped. The momentum dependence of $\omega_0(|\mathbf{q}|, \alpha_1, \alpha_2)$ is shown in Figure 7.

The summation over α_1 and α_2 yields four distinct contributions to the decay width. For $\alpha_1 = \alpha_2 = +1$, we obtain the contribution from the familiar 1 → 2 decay process



Figure 9: Contour plots of u versus θ for the $1 \to 2$ decay contribution to $\mathrm{d}^2\Gamma_\Phi^{(1)}$ for discrete values of Mt, with $q^2 = M^2$ and $|\mathbf{q}| = 10$ GeV. The solid excluded regions to the right of the red dotted line lie exterior to the limits of integration over u.

presented in Figure 8a. For $\alpha_1 = -\alpha_2 = \pm 1$, we obtain the two $2 \to 1$ *Landau–damping* contributions displayed in Figure 8b. For $\alpha_1 = \alpha_2 = -1$, we obtain the $3 \to 0$ *total annihilation* process shown in Figure 8c. In the latter, the decay 'products' appear in the *initial state* along with the decaying particle. For late times, the Landau damping and total annihilation processes are kinematically disallowed, as we would expect. They are only permitted in the *evanescent regime* at early times.

Figures 9–10 contain contour plots of u versus θ for the four contributions to the on-shell differential width $\mathrm{d}^2\Gamma_\Phi^{(1)}$. The equilibrium kinematics are obtained in the large-time limit $t \to \infty$ as follows. As can be seen from the (u, θ) contour plots in Figure 9, for the $1 \to 2$ decay process, the limits of integration grow to encompass the full range of the sinc function. At the same time, the u dependence of the phase-space pre-factors vanishes, since $q_u(t) \to q_0$ as $t \to \infty$. For the $3 \to 0$ total annihilation process, the domain of integration vanishes in the large-time limit $t \to \infty$. Given that the integrand is finite in the same limit, the contribution therefore vanishes as expected, which is confirmed by our (u, θ) contour plots in Figure 10. For the two Landau–damping contributions, the large-time behaviour becomes more subtle. As $t \to \infty$, the domain of integration covers approximately all positive u. However, the kinematically allowed phase space cannot be attained with any values of u and θ, so that the two Landau–damping contributions also vanish in the large-time limit. This behaviour is reflected in our (u, θ) contour plots of Figure 11. Thus, only the usual $1 \to 2$ energy-conserving decay remains for late times.

In order to reduce the statistical error in our `Monte Carlo` integration over the (u, θ)



Figure 10: Contour plots of u versus θ for the $3 \to 0$ total annihilation contribution to $\mathrm{d}^2\Gamma_\Phi^{(1)}$ for discrete values of Mt, where $q^2 = M^2$ and $|\mathbf{q}| = 10$ GeV. The regions to the left of the red dotted line are exterior to the domain of integration over u.

phase space, we use a Gaussian sampling bias to ensure that the majority of sampling points fall over the dominant region of the sinc function of u in (9.21). We define the weight function

$$\varpi(u) \;=\; \frac{\mathrm{d}u}{\mathrm{d}r} \;\equiv\; \exp\left(-\,\frac{(u\,-\,u_0)^2}{2\sigma_u^2}\right) \tag{9.26}$$

where

$$r(u) \;\equiv\; \frac{\mathrm{Erf}\left(\frac{1}{\sqrt{2}}\,\frac{u-u_0}{\sigma_u}\right)\;-\;\mathrm{Erf}\left(\frac{1}{\sqrt{2}}\,\frac{u_- - u_0}{\sigma_u}\right)}{\mathrm{Erf}\left(\frac{1}{\sqrt{2}}\,\frac{u_+ - u_0}{\sigma_u}\right)\;-\;\mathrm{Erf}\left(\frac{1}{\sqrt{2}}\,\frac{u_- - u_0}{\sigma_u}\right)} \;\in\; [0,\;1]\;. \tag{9.27}$$

After performing the change of the variable $u \to r$ in (9.21) for $u_0 = 0$, the limits of integration become time-independent and are identical for the decay, total-annihilation and Landau–damping contributions. The dependence upon the limits $u_+(t)$ and $u_-(t)$ appears instead within the transformed integrand of (9.21) in the new variables (r, θ).

The width of the dominant region of the u phase-space is taken to be the distance between the central maximum and the maximum at which the amplitude of the sinc function has fallen to 0.1% with respect to the central maximum. Hence, we require for the distant maximum to be at the location where $\mathrm{sinc}\, u_n \sim 0.001$. The extrema u_n of the sinc function satisfy the transcendental equation

$$u_n \;=\; \tan u_n\;, \tag{9.28}$$



(a) $\alpha_1 \;=\; -\alpha_2 \;=\; +1$



(b) $\alpha_1 \;=\; -\alpha_2 \;=\; -1$

Figure 11: Contour plots of u versus θ for the two Landau–damping contributions ($\alpha_1 = -\alpha_2 = \pm 1$) to $\mathrm{d}^2\Gamma_\Phi^{(1)}$ for discrete values of Mt, assuming $q^2 = M^2$ and $|\mathbf{q}| = 10$ GeV. The regions to the right of the red dotted line are exterior to the domain of integration over u. The contour plots differ between the two contributions due to the asymmetry of the integrands in the Φ three-momentum.



Figure 12: The four separate contributions to the ratio $\bar{\Gamma}_{\Phi}^{(1)}$ in (9.30) versus Mt, for on-shell decays with $|\mathbf{q}| = 1$ GeV, 10 GeV and 100 GeV. The two Landau–damping contributions are identical up to numerical errors.

whose solutions for $n \geq 1$ may be expressed as [76]

$$u_n = \pm n\pi \exp\left\{\frac{1}{\pi}\int_0^1 \mathrm{d}\xi\, \frac{1}{\xi} \arg\left[\left(1 + \frac{1}{2}\xi \ln\frac{1-\xi}{1+\xi} \pm \frac{1}{2}\pi i \xi\right)^2 + n^2\pi^2\xi^2\right]\right\}. \qquad (9.29)$$

The required extremum is then given by $n = 318$, the 159th maximum of the sinc function, with $u_{318} = 1000$. The Gaussian weight function $\varpi(u)$ in (9.26) is then taken to have a variance of $\sigma_u^2 = (u_{318}/2)^2$, such that 95% of the sampling points fall within this dominant region.



Figure 13: The ratio $\bar{\Gamma}_{\Phi}^{(1)}$ in (9.30) versus Mt, for on-shell decays with $|\mathbf{q}| = 1$ GeV (solid black), 10 GeV (blue dotted) and 100 GeV (red dashed).

Our interest is in the deviation of the one-loop Φ width $\Gamma_{\Phi}^{(1)}$ from the known equilibrium result. It is therefore convenient to define the ratio $\bar{\Gamma}_{\Phi}^{(1)}(|\mathbf{q}|,t)$ of the time-dependent width to its late-time equilibrium value for $q^2 = M^2$:

$$\bar{\Gamma}_{\Phi}^{(1)}(|\mathbf{q}|,t) \;=\; \frac{\Gamma_{\Phi}^{(1)}(|\mathbf{q}|,t)}{\Gamma_{\Phi}^{(1)}(|\mathbf{q}|,t\,\to\,\infty)} \;=\; \frac{\Gamma_{\Phi}^{(1)1\to 2}(|\mathbf{q}|,t)\;+\;2\Gamma_{\Phi}^{(1)\mathrm{Landau}}(|\mathbf{q}|,t)\;+\;\Gamma_{\Phi}^{(1)3\to 0}(|\mathbf{q}|,t)}{\Gamma_{\Phi}^{(1)}(|\mathbf{q}|,t\to\infty)}\,. \tag{9.30}$$

In Figure 12, we plot separately the four contributions from Figure 8 to this ratio as a function of Mt for a series of discrete momenta. The evanescent Landau–damping processes yield a prompt contribution, which can be as high as 10–20% at early times. The evanescent total annihilation process contributes similarly at the level of about 5%.

In Figure 13, we plot the total ratio $\bar{\Gamma}_{\Phi}^{(1)}(|\mathbf{q}|,t)$ as a function of Mt. The total Φ width $\Gamma_{\Phi}^{(1)}(|\mathbf{q}|,t)$ is vanishing for $Mt = 0$, as we would expect. This is followed promptly by a sharp rise, which is particularly pronounced in the infra-red modes, resulting from the sudden switching on of the interactions. This so-called *shock regime* is followed by the superposition transient oscillations of angular frequency ω_0 with short time scales and *non-Markovian* oscillations of longer time scales. The latter of these oscillations exhibit time-dependent frequencies, the origin of which will be discussed in the next section.



Figure 14: Non-Markovian oscillations of the $1 \to 2$ decay contribution to the ratio $\bar{\Gamma}_\Phi^{(1)}$ in (9.30) against Mt, by performing a moving average over bins of $150Mt$.

9.3 Non-Markovian Oscillations

As we have seen in Figures 12 and 13, the time-dependent Φ width contains a superposition of damped oscillatory contributions. The longer-lived of these oscillations exhibit time-dependent frequencies. These non-Markovian oscillations are illustrated more clearly in Figure 14 for the $|\mathbf{q}| = 1$ GeV mode in which a moving time average is carried out to eliminate the higher-frequency Markovian oscillations. In this section, we describe the origin of the time-dependent oscillations and show that they are not a numerical artefact, but are instead an intrinsic feature inherent to the dynamics of truly out-of-equilibrium systems. To this end, we consider the high-temperature limit $T \gg M$ of the time-dependent Φ width.

In this high-temperature limit $T \gg M$, the Bose–Einstein distribution may be approximated as follows:

$$f_{\rm B}(E) \;\approx\; \frac{T}{E}\,, \tag{9.31}$$

Returning to the Φ width $\Gamma_\Phi^{(1)}$ in (9.8) and substituting for (9.31), we may perform the angular integration by making the following change of variables:

$$\cos\theta \;=\; \frac{|\mathbf{k}|^2 \,+\, |\mathbf{q}|^2 \,+\, m_2^2}{2|\mathbf{k}||\mathbf{q}|}\,(1\,-\,x)^2\,. \tag{9.32}$$

Then, the high-temperature limit $T \gg M$ of the time-dependent width of the heavy scalar Φ becomes

$$\begin{aligned}
\Gamma_\Phi^{(1),\,T\,\gg\,M}(q,t) \;=\; & \frac{g^2}{32\pi^2 M}\sum_{\{\alpha\}}\alpha_1\alpha_\theta\int_0^\infty \mathrm{d}|\mathbf{k}|\,\frac{|\mathbf{k}|}{|\mathbf{q}|}\,\frac{1}{E_1} \\
& \times\,\Bigg[\bigg(1 + \frac{T}{\alpha_1 E_1} + \frac{T}{q_0 - \alpha_1 E_1}\bigg)\mathrm{Si}\big[\big(q_0 - \alpha_1 E_1 - \alpha_2 E_2\big)t\big] \\
& -\,\frac{T}{q_0 - \alpha_1 E_1}\bigg(\sin\big[\big(q_0 - \alpha_1 E_1\big)t\big]\mathrm{Ci}\big(\alpha_2 E_2 t\big) - \cos\big[\big(q_0 - \alpha_1 E_1\big)t\big]\mathrm{Si}\big(\alpha_2 E_2 t\big)\bigg)\Bigg]\,,
\end{aligned} \tag{9.33}$$

where $\mathrm{Si}(x)$ and $\mathrm{Ci}(x)$ are respectively the sine integral and cosine integral functions. We have introduced the short-hand notations: $\{\alpha\}$ for the summation over $\alpha_1,\ \alpha_2,\ \alpha_\theta = \pm 1$; and

$$E_1 \;=\; \sqrt{|\mathbf{k}|^2 + m_1^2}\,, \tag{9.34}$$

$$E_2 \;=\; \sqrt{|\mathbf{k}|^2 - 2\alpha_\theta|\mathbf{k}||\mathbf{q}| + |\mathbf{q}|^2 + m_2^2}\,. \tag{9.35}$$

In terms of the evanescent action u given in (9.13), the high-temperature limit of the one-loop Φ width reads:

$$\begin{aligned}
\Gamma_\Phi^{(1),\,T\,\gg\,M}(q,t) \;=\; & \frac{g^2}{32\pi^2 M t}\sum_{\{\alpha\},\,b\,=\,\pm1}\alpha_1\alpha_2\int_{u_-(t)}^{u_+(t)}\mathrm{d}u \\
& \times\,\Bigg\{\frac{q_u(t)}{\lambda^{1/2}\big(q_u^2(t) - |\mathbf{q}|^2, m_1^2, m_2^2\big)}\Bigg[\frac{m_1^2 + m_2^2}{q_u^2(t) - |\mathbf{q}|^2} - \bigg(\frac{m_1^2 - m_2^2}{q_u^2(t) - |\mathbf{q}|^2}\bigg)^2\Bigg] \\
& +\,\frac{b\alpha_\theta}{2|\mathbf{q}|}\bigg(1 - \frac{m_1^2 - m_2^2}{q_u^2(t) - |\mathbf{q}|^2}\bigg)\Bigg\} \\
& \times\,\Bigg[\bigg(1 + \frac{T}{\omega_1^{(b)}(q,u,t)} + \frac{T}{q_u(t) - \omega_2^{(b)}(q,u,t)}\bigg)\mathrm{Si}(u) \\
& -\,\frac{T}{\omega_1^{(b)}(q,u,t)}\bigg(\sin\big(\omega_1^{(b)}(q,u,t)t\big)\,\mathrm{Ci}\big(\omega_2^{(b)}(q,u,t)t\big) \\
& -\,\cos\big(\omega_1^{(b)}(q,u,t)t\big)\,\mathrm{Si}\big(\omega_2^{(b)}(q,u,t)t\big)\bigg)\Bigg]\,,
\end{aligned} \tag{9.36}$$

where $\omega_1^{(b)}(q,u,t)$ and $\omega_2^{(b)}(q,u,t)$ are time-dependent non-Markovian frequencies defined as

$$\omega_1^{(b)}(q,u,t) \;=\; q_0 - \frac{\big(q_u^2(t) - |\mathbf{q}|^2 + m_1^2 - m_2^2\big)q_u(t) + b\alpha_\theta|\mathbf{q}|\lambda^{1/2}\big(q_u^2(t) - |\mathbf{q}|^2, m_1^2, m_2^2\big)}{2\big(q_u^2(t) - |\mathbf{q}|^2\big)}\,, \tag{9.37a}$$

$$\omega_2^{(b)}(q,u,t) \;=\; \frac{\big(q_u^2(t) - |\mathbf{q}|^2 - m_1^2 + m_2^2\big)q_u(t) - b\alpha_\theta|\mathbf{q}|\lambda^{1/2}\big(q_u^2(t) - |\mathbf{q}|^2, m_1^2, m_2^2\big)}{2\big(q_u^2(t) - |\mathbf{q}|^2\big)}\,. \tag{9.37b}$$



Figure 15: $\omega_1^{(b)}(q,u,t)/M$ versus Mt for the first three extrema of sinc(u): $u_1 = 4.49$ (black), $u_2 = 7.73$ (blue) and $u_3 = 10.90$ (red), obtained by (9.29), for $q^2 = M^2$, $|\mathbf{q}| = 10$ GeV and $\alpha_\theta = +1$. Solid lines correspond to $b = 1$ and dashed lines, $b = -1$ in (9.37a). The dotted lines mark the upper limit of the kinematically disallowed region.

The time dependence of the frequency $\omega_1^{(b)}(q,u,t)$ is shown in Figure 15 for $\alpha_\theta = +1$, with the on-shell constraint $q^2 = M^2$. The $b = 1$ contribution persists for late times, at which point the frequency of the modulations have decayed to zero. The $b = -1$ contribution is disallowed for late times, such that the amplitude of this constant-frequency modulation is damped to zero. For $\alpha_\theta = -1$, the $b = +1$ and $b = -1$ contributions are interchanged, such that the frequency of the $b = -1$ contribution reduces to zero for late times. Thus, we obtain the expected kinematics and equilibrium behaviour in the late time limit. We have not plotted $\omega_2^{(b)}$, as its behaviour is indistinguishable from $\omega_1^{(b)}$, for $m_1 = m_2 = m$.

Looking again at (9.36), we observe that these non-Markovian oscillations occur only for $T \neq 0$. We conclude therefore that this behaviour signifies a genuine non-equilibrium statistical effect.

9.4 Perturbative Time Evolution Equations

In this section, we carry out a loopwise expansion of the time evolution equations derived in Section 8 to leading order in the coupling g, evaluating their one-loop structure for our simple scalar model. For perturbatively small couplings, such a loopwise expansion is expected to accurately capture the early-time dynamics of non-equilibrium systems. This is a regime, in which the applicability of truncated gradient expansions becomes questionable, according to our discussion in Section 4.2. Given the closed analytic form of the free CTP propagators, both the amplitudes of contributing processes and the resulting phase-space integrals describing the kinematics can be analytically determined. The systematic treatment of these kinematic effects is essential to understand the consequences on the early-time dynamics. In particular, we illustrate the significance of contributions from the energy-non-conserving evanescent regime to the dynamics of the system.

Let us first return to the Boltzmann–like equation in (8.17), in which we artificially imposed energy conservation. The collision terms for the real scalar Φ are

$$\Pi_{\Phi,>}(p,t)\Delta_{\Phi,<}(p,t) \;-\; \Pi_{\Phi,<}(p,t)\Delta_{\Phi,>}(p,t)\ . \tag{9.38}$$

Since we have assumed that the two subsystems $\mathscr{S}_{\Phi}$ and $\mathscr{S}_{\chi}$ are separately in thermodynamic equilibrium at the same temperature T at the initial time $t = 0$, the Φ propagators and self-energies will initially satisfy the KMS relation, i.e.

$$\Delta_{\Phi,>}(p,0) \;=\; e^{\beta p_0}\Delta_{\Phi,<}(p,0)\ , \qquad \Pi_{\Phi,>}(p,0) \;=\; e^{\beta p_0}\Pi_{\Phi,<}(p,0)\ . \tag{9.39}$$

As a consequence of the KMS relation, the collision terms in (9.38) are identical to zero. This is true also in the collision terms of the gradient-expanded Kadanoff–Baym kinetic equation in (E.5b). With no external sources present to perturb the combined system $\mathscr{S}$ from this non-interacting equilibrium, we are faced with the problem that the statistical distribution functions will *not* evolve from their initial forms. In reality, we anticipate that the system should evolve to some new interacting thermodynamic equilibrium for late times. We conclude therefore that the energy-non-conserving evanescent regime described by our approach is entirely necessary to account correctly for the evolution of this system.

We now turn our attention to the master time evolution equation (8.14). Truncating to leading order in the coupling g, the RHS of this time evolution equation contains free propagators and one-loop self-energies. For instance, the one-loop collision terms in (8.14) are

$$\begin{aligned}\mathscr{C}^{(1)}(p+\tfrac{P}{2},p-\tfrac{P}{2},t;0) \;\equiv\; \frac{1}{2}\int\frac{\mathrm{d}^4q}{(2\pi)^4}\,\Big[\, & i\Pi^{(1)}_{>}(p_1,q,t;0)i\Delta^{0}_{<}(q,p_2,t;0)\\ & -\; i\Pi^{(1)}_{<}(p_1,q,t)\Big(i\Delta^{0}_{>}(q,p_2,t;0)\;-\;2i\Delta^{0}_{\mathcal{P}}(q,p_2,t;0)\Big)\Big]\ .\end{aligned} \tag{9.40}$$

We insert the one-loop integrals (9.3a) and (9.3b), containing the homogeneous Φ and χ distribution functions, $f_{\Phi}(|\mathbf{p}|,t)$ and $f_{\chi}^{(C)}(|\mathbf{p}|,t)$, into the full time evolution equation (8.14) for the Φ and χ fields. Note that the f distribution functions are real and the g distribution functions vanish in this spatially homogeneous limit. We tacitly assume that a system initially prepared in a spatially homogeneous state remains spatially homogeneous throughout its evolution. This assumption is reasonable if the system has infinite spatial extent.

We may perform the loop and convolution integrals using the techniques and results of Appendix F. After carrying out the p_0 and P_0 integrals in (8.14), we obtain the following one-loop time evolution equation for the spatially homogeneous statistical distribution function $f_\Phi(|\mathbf{p}|,t)$ of the real scalar field:

$$\begin{aligned}\partial_t f_\Phi(|\mathbf{p}|,t) \ &= \ -\frac{g^2}{2}\sum_{\{\alpha\}}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{1}{2E_\Phi(\mathbf{p})}\,\frac{1}{2E_\chi(\mathbf{k})}\,\frac{1}{2E_\chi(\mathbf{p}-\mathbf{k})} \\ &\times\ \frac{t}{2\pi}\,\mathrm{sinc}\big[\big(\alpha E_\Phi(\mathbf{p})\ -\ \alpha_1E_\chi(\mathbf{k})\ -\ \alpha_2E_\chi(\mathbf{p}-\mathbf{k})\big)t/2\big] \\ &\times\ \Big\{\pi\ +\ 2\mathrm{Si}\big[\big(\alpha E_\Phi(\mathbf{p})\ +\ \alpha_1E_\chi(\mathbf{k})\ +\ \alpha_2E_\chi(\mathbf{p}-\mathbf{k})\big)t/2\big]\Big\} \\ &\times\ \Big\{\Big(\theta(-\alpha)\ +\ f_\Phi(|\mathbf{p}|,t)\Big)\Big[\theta(\alpha_1)\Big(1\ +\ f_\chi(|\mathbf{k}|,t)\Big)\ +\ \theta(-\alpha_1)f_\chi^C(|\mathbf{k}|,t)\Big] \\ &\qquad\quad\times\ \Big[\theta(\alpha_2)\Big(1\ +\ f_\chi^C(|\mathbf{p}-\mathbf{k}|,t)\Big)\ +\ \theta(-\alpha_2)f_\chi(|\mathbf{p}-\mathbf{k}|,t)\Big] \\ &-\ \Big(\theta(\alpha)\ +\ f_\Phi(|\mathbf{p}|,t)\Big)\Big[\theta(\alpha_1)f_\chi(|\mathbf{k}|,t)\ +\ \theta(-\alpha_1)\Big(1\ +\ f_\chi^C(|\mathbf{k}|,t)\Big)\Big] \\ &\qquad\quad\times\ \Big[\theta(\alpha_2)f_\chi^C(|\mathbf{p}-\mathbf{k}|,t)\ +\ \theta(-\alpha_2)\Big(1\ +\ f_\chi(|\mathbf{p}-\mathbf{k}|,t)\Big)\Big]\Big\}\,. \end{aligned} \tag{9.41}$$

where $\mathrm{Si}(x)$ is the sine integral function and $\{\alpha\}$ is the short-hand notation for the summation over $\alpha,\ \alpha_1,\ \alpha_2\ =\ \pm1$. With the summation over $\{\alpha\}$, the statistical factors in the braces of the last four lines of (9.41) contain the difference of contributions from the four processes shown in Figure 8 and their inverse processes. For early times, all of these evanescent processes and inverse processes contribute, as can be seen from Figure 12. The presence of $\mathrm{Si}(x)$ ensures that the correct late-time limit and kinematics are obtained on restoration of energy conservation. This factor is missing in the existing descriptions, see e.g. [45, 77]. The dispersive force term and off-shell collision term vanish in the spatially homogeneous case, thanks to the symmetry of the self-energy under $P\ \to\ -P$.

In the large-time limit $t\ \to\ \infty$, we have

$$\begin{aligned}&\frac{t}{2\pi}\,\mathrm{sinc}\big[\big(\alpha E_\Phi(\mathbf{p})\ -\ \alpha_1E_\chi(\mathbf{k})\ -\ \alpha_2E_\chi(\mathbf{p}-\mathbf{k})\big)t/2\big] \\ &\qquad\times\ \Big\{\pi\ +\ 2\mathrm{Si}\big[\big(\alpha E_\Phi(\mathbf{p})\ +\ \alpha_1E_\chi(\mathbf{k})\ +\ \alpha_2E_\chi(\mathbf{p}-\mathbf{k})\big)t/2\big]\Big\} \\ &\qquad\qquad\xrightarrow[t\,\to\,\infty]{}\ 2\pi\theta(\alpha)\delta\big(E_\Phi(\mathbf{p})\ -\ \alpha_1E_\chi(\mathbf{k})\ -\ \alpha_2E_\chi(\mathbf{p}-\mathbf{k})\big)\,.\end{aligned} \tag{9.42}$$

The kinematic constraints then force $\alpha\ =\ \alpha_1\ =\ \alpha_2\ =\ +1$ and we obtain

$$\begin{aligned}\partial_t f_\Phi(|\mathbf{p}|,t)\ &=\ -\frac{g^2}{2}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{1}{2E_\Phi(\mathbf{p})}\,\frac{1}{2E_\chi(\mathbf{k})}\,\frac{1}{2E_\chi(\mathbf{p}-\mathbf{k})} \\ &\times\ 2\pi\delta\big(E_\Phi(\mathbf{p})\ -\ E_\chi(\mathbf{k})\ -\ E_\chi(\mathbf{p}-\mathbf{k})\big) \\ &\times\ \Big[f_\Phi(|\mathbf{p}|,t)\Big(1\ +\ f_\chi(|\mathbf{k}|,t)\Big)\Big(1\ +\ f_\chi^C(|\mathbf{p}-\mathbf{k}|,t)\Big) \\ &-\ \Big(1\ +\ f_\Phi(|\mathbf{p}|,t)\Big)f_\chi(|\mathbf{k}|,t)f_\chi^C(|\mathbf{p}-\mathbf{k}|,t)\Big]\,.\end{aligned} \tag{9.43}$$

Equation (9.43) corresponds to the semi-classical Boltzmann equation [cf. (8.17)].

By analogy, the one-loop time evolution equation for the spatially homogeneous statistical distribution function $f_\chi(|\mathbf{p}|,t)$ of the complex scalar field χ is given by

$$\begin{aligned}
\partial_t f_\chi(|\mathbf{p}|,t) \;=\; & -\frac{g^2}{2}\sum_{\{\alpha\}}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{1}{2E_\Phi(\mathbf{p})}\,\frac{1}{2E_\chi(\mathbf{k})}\,\frac{1}{2E_\chi(\mathbf{p}-\mathbf{k})} \\
& \times\,\frac{t}{2\pi}\,\mathrm{sinc}\big[\big(\alpha E_\chi(\mathbf{p})\,-\,\alpha_1E_\Phi(\mathbf{k})\,-\,\alpha_2E_\chi(\mathbf{p}-\mathbf{k})\big)t/2\big] \\
& \times\,\Big\{\pi\,+\,2\mathrm{Si}\big[\big(\alpha E_\chi(\mathbf{p})\,+\,\alpha_1E_\Phi(\mathbf{k})\,+\,\alpha_2E_\chi(\mathbf{p}-\mathbf{k})\big)t/2\big]\Big\} \\
& \times\,\bigg\{\Big[\theta(\alpha)f_\chi(|\mathbf{p}|,t)\,+\,\theta(-\alpha)\Big(1\,+\,f^C_\chi(|\mathbf{p}|,t)\Big)\Big]\Big(\theta(\alpha_1)\,+\,f_\Phi(|\mathbf{k}|,t)\Big) \\
& \qquad\quad\times\,\Big[\theta(\alpha_2)\Big(1\,+\,f_\chi(|\mathbf{p}-\mathbf{k}|,t)\Big)\,+\,\theta(-\alpha_2)f^C_\chi(|\mathbf{p}-\mathbf{k}|,t)\Big] \\
& -\,\Big[\theta(\alpha)\Big(1\,+\,f_\chi(|\mathbf{p}|,t)\Big)\,+\,\theta(-\alpha)f^C_\chi(|\mathbf{p}|,t)\Big]\Big(\theta(-\alpha_1)\,+\,f_\Phi(|\mathbf{p}|,t)\Big) \\
& \qquad\quad\times\,\Big[\theta(\alpha_2)f_\chi(|\mathbf{p}-\mathbf{k}|,t)\,+\,\theta(-\alpha_2)\Big(1\,+\,f^C_\chi(|\mathbf{p}-\mathbf{k}|,t)\Big)\Big]\bigg\}\,.
\end{aligned}\tag{9.44}$$

In the large-time limit $t\,\to\,\infty$, the kinematics restrict $\alpha\,=\,\alpha_1\,=\,-\alpha_2\,=\,+1$, giving

$$\begin{aligned}
\partial_t f_\chi(|\mathbf{p}|,t) \;=\; & -\frac{g^2}{2}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{1}{2E_\Phi(\mathbf{p})}\,\frac{1}{2E_\chi(\mathbf{k})}\,\frac{1}{2E_\chi(\mathbf{p}-\mathbf{k})} \\
& \times\,2\pi\delta\big(E_\Phi(\mathbf{k})\,-\,E_\chi(\mathbf{p})\,-\,E_\chi(\mathbf{p}-\mathbf{k})\big) \\
& \times\,\Big[\Big(1\,+\,f_\Phi(|\mathbf{k}|,t)\Big)f_\chi(|\mathbf{p}|,t)f^C_\chi(|\mathbf{p}-\mathbf{k}|,t) \\
& -\,f_\Phi(|\mathbf{k}|,t)\Big(1\,+\,f_\chi(|\mathbf{p}|,t)\Big)\Big(1\,+\,f^C_\chi(|\mathbf{p}-\mathbf{k}|,t)\Big)\Big]\,.
\end{aligned}\tag{9.45}$$

This result is consistent with (9.43), differing by an overall sign, as one should expect.

At $t\,=\,0$, the semi-classical Boltzmann transport equation in (9.43) becomes

$$\begin{aligned}
\partial_t f_\Phi(|\mathbf{p}|,t)\big|_{t\,=\,0} \;=\; & -\frac{g^2}{2}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{1}{2E_\Phi(\mathbf{p})}\,\frac{1}{2E_\chi(\mathbf{k})}\,\frac{1}{2E_\chi(\mathbf{p}-\mathbf{k})} \\
& \times\,2\pi\delta\big(E_\Phi(\mathbf{p})\,-\,E_\chi(\mathbf{k})\,-\,E_\chi(\mathbf{p}-\mathbf{k})\big) \\
& \times\,\Big[f_{\mathrm{B}}\big(E_\Phi(\mathbf{p})\big)\Big(1\,+\,f_{\mathrm{B}}\big(E_\chi(\mathbf{k})\big)\,+\,f^C_{\mathrm{B}}\big(E_\chi(\mathbf{p}-\mathbf{k})\big)\Big) \\
& -\,f_{\mathrm{B}}\big(E_\chi(\mathbf{k})\big)f^C_{\mathrm{B}}\big(E(\mathbf{p}-\mathbf{k})\big)\Big]\,,
\end{aligned}\tag{9.46}$$

where we remind the reader that energy conservation is artificially imposed. By virtue of this energy conservation, the first product of statistical factors in (9.46) satisfies the identity

$$f_{\mathrm{B}}\big(E_\Phi(\mathbf{p})\big)\Big(1\,+\,f_{\mathrm{B}}\big(E_\chi(\mathbf{k})\big)\,+\,f^C_{\mathrm{B}}\big(E_\chi(\mathbf{p}-\mathbf{k})\big)\Big)\;=\;f_{\mathrm{B}}\big(E_\chi(\mathbf{k})\big)f^C_{\mathrm{B}}\big(E(\mathbf{p}-\mathbf{k})\big)\,. \tag{9.47}$$

As a result, the RHS of (9.43) is exactly zero at $t\,=\,0$. Consequently, the energy-non-conserving evanescent regime plays a fundamental role in the description of the evolution of the system $\mathscr{S}$.

We now wish to show that the master time evolution equations (9.41) and (9.44) of our perturbative approach describe a non-trivial evolution of the system $\mathscr{S}$. For this



Figure 16: Numerical estimates of $\partial_t f_\Phi(|\mathbf{p}|, t)$, as a function of Mt, assuming that the χ statistical distribution functions $f_\chi^{(C)}(|\mathbf{p}|, t)$ on the RHS of the time evolution equation (9.41) maintain their equilibrium Bose–Einstein form for all times.

purpose, we consider the time evolution equation of the heavy scalar Φ in (9.41) and assume that the χ statistical distribution functions $f_\chi^{(C)}(|\mathbf{p}|, t)$ of the RHS remain in their initial equilibrium forms for all times, i.e. $f_\chi^{(C)}(|\mathbf{p}|, t) = f_{\mathrm{B}}\big(E_\chi(\mathbf{p})\big)$. With this assumption, we see from Figure 16 that the RHS of (9.41) is non-zero for early times. Thus, the system $\mathscr{S}$ is indeed perturbed from its non-interacting equilibrium by the evanescent processes described by our perturbative time evolution equations (9.41) and (9.44).

Let us finally have a closer look at the early-time behaviour that immediately follows after the switching on of the interactions. We expect that this prompt behaviour is dominated by the ultra-violet contribution to the phase-space integral on the RHS of (9.41) due to the Heisenberg uncertainty principle. From Figure 13, we see that this shock regime contributes dominantly to the prompt evolution of the infra-red modes of $f_\Phi(|\mathbf{p}|, t)$. In this regime, we take $|\mathbf{p}| = 0$ and $m = 0$ in (9.41) and introduce the ultra-violet cut-off Λ in the limits of the phase-space integrals of the RHS. Assuming that the χ distribution functions are tempered, vanishing faster than a power law for large momenta, we may ignore their contribution in the ultra-violet limit. Hence, $\partial_t f_\Phi(|\mathbf{p}| = 0, t)$ can be approximated by

$$\partial_t f_\Phi(|\mathbf{p}| = 0, t) \;\sim\; -\frac{g^2}{32\pi^3 M}\Big(1 \,+\, 2f_\Phi(|\mathbf{p}| = 0, t)\Big) \lim_{\Lambda\to\infty} \mathrm{Si}^2(\Lambda t)\ . \tag{9.48}$$

Clearly, $\partial_t f_\Phi(|\mathbf{p}| = 0, t)$ vanishes in the limit $t \ll 1/\Lambda \to 0$, as we expect. Expanding $\mathrm{Si}\,(\Lambda t)$ about $\Lambda t = 0$, i.e. for times infinitesimally close to zero, we obtain

$$\partial_t f_\Phi(|\mathbf{p}| = 0, t) \;\sim\; -\frac{g^2}{32\pi^3 M}\Big(1 \,+\, 2f_\Phi(|\mathbf{p}| = 0, 0)\Big) \lim_{\Lambda\to\infty} (\Lambda t)^2\ . \tag{9.49}$$

For small but finite times, ultra-violet contributions are rapidly varying. It is apparent from this discussion and Figures 13 and 16 that the effect of the transient behaviour of the system upon its subsequent dynamics is significant, since the quantum memories persist on long time scales $Mt \gg 1$. We may therefore conclude that as opposed to other methods relying on a truncated gradient expansion, our approach consistently describes this highly-oscillatory and rapidly-evolving early-time behaviour of the system.

9.5 Inclusion of Thermal Masses

In this section, we describe how local thermal-mass corrections may be incorporated consistently into our approach.

The local part of the one-loop χ self-energy shown in Figure 4 has the explicit form

$$\begin{aligned}\Pi_\chi^{\mathrm{loc}(1)}(p, p', \tilde{t}_f; \tilde{t}_i) \;&=\; -\frac{\lambda}{2}(2\pi)^4\delta_t^{(4)}(p \,-\, p')(2\pi\mu)^{2\epsilon}e^{i(p_0 \,-\, p'_0)\tilde{t}_f} \\ &\times \iint \frac{\mathrm{d}^d k}{(2\pi)^d}\,\frac{\mathrm{d}^4 k'}{(2\pi)^4}\,\bigg(\frac{i}{k^2 \,-\, m^2 \,+\, i\epsilon}(2\pi)^4\delta^{(4)}(k \,-\, k') \\ &+\, 2\pi\delta(k^2 \,-\, m^2)|2k_0|^{1/2}\tilde{f}_\chi(k, k', t)e^{i(k_0 \,-\, k'_0)\tilde{t}_f}|2k'_0|^{1/2}2\pi\delta(k'^2 \,-\, m^2)\bigg)\ ,\end{aligned} \tag{9.50}$$

with $d = 4 - 2\epsilon$ (cf. Appendix F). The first term yields the standard zero-temperature UV divergence, which is usually removed by mass renormalization. The second term yields

$$\Pi_\chi^{\mathrm{loc}(1)}(p, p', \tilde{t}_f; \tilde{t}_i) \;=\; -(2\pi)^4\delta_t^{(4)}(p \,-\, p')e^{i(p_0 \,-\, p'_0)\tilde{t}_f}m_{\mathrm{th}}^2(\tilde{t}_f; \tilde{t}_i)\ , \tag{9.51}$$

where the time-dependent thermal mass $m_{\mathrm{th}}(\tilde{t}_f; \tilde{t}_i)$, given by

$$\begin{aligned}m_{\mathrm{th}}^2(\tilde{t}_f; \tilde{t}_i) \;&=\; \frac{\lambda}{2}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{1}{\sqrt{2E_\chi(\mathbf{k})}}\int\frac{\mathrm{d}^3\mathbf{k}'}{(2\pi)^3}\,\frac{1}{\sqrt{2E_\chi(\mathbf{k}')}} \\ &\times\,\Big(f_\chi(\mathbf{k}, \mathbf{k}', t)e^{i[E(\mathbf{k}) \,-\, E(\mathbf{k}')]\tilde{t}_f} \,+\, f_\chi^{C*}(-\mathbf{k}, -\mathbf{k}', t)e^{-i[E(\mathbf{k}) \,-\, E(\mathbf{k}')]\tilde{t}_f}\Big)\ ,\end{aligned} \tag{9.52}$$

is UV finite. Notice that a unique thermal mass may not be defined in the spatially inhomogeneous case due to the explicit dependence on $\tilde{t}_f$.

From the Schwinger–Dyson equation in (4.32), the inverse quasi-particle CTP propagator $\Delta^{-1}_{\chi,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i)$ of the complex scalar field χ takes the form

$$\Delta^{-1}_{\chi,\,ab}(p,p',\tilde{t}_f;\tilde{t}_i) \;=\; (2\pi)^4\delta_t^{(4)}(p-p')e^{i(p_0\,-\,p_0')\tilde{t}_f}\Big[\Big(p'^2 - m^2_{\rm th}(\tilde{t}_f;\tilde{t}_i)\Big)\eta_{ab} + i\epsilon\mathbb{I}_{ab}\Big]\;, \tag{9.53}$$

in which we have assumed $m_{\rm th}(t) \gg m$. If the quartic χ self-interaction is switched on sufficiently long before $t = 0$, then for $t \geq 0$, we may replace the δ_t function in the inverse quasi-particle propagator (9.53) by an exact energy-conserving delta function. This imposition of energy conservation constitutes a quasi-particle approximation, allowing us to invert (9.53) exactly, using the arguments of Sections 3.2 and 4.3, to obtain the following quasi-particle χ propagators:

$$\begin{aligned}
&\Delta^{0,\,ab}_{\chi}(p,p',\tilde{t}_f;\tilde{t}_i) \qquad\qquad (9.54)\\
&= \begin{bmatrix} \big(p^2 - m^2_{\rm th}(\tilde{t}_f;\tilde{t}_i) + i\epsilon\big)^{-1} & -i2\pi\theta(-p_0)\delta\big(p^2 - m^2_{\rm th}(\tilde{t}_f;\tilde{t}_i)\big) \\ -i2\pi\theta(p_0)\delta\big(p^2 - m^2_{\rm th}(\tilde{t}_f;\tilde{t}_i)\big) & -\big(p^2 - m^2_{\rm th}(\tilde{t}_f;\tilde{t}_i) - i\epsilon\big)^{-1}\end{bmatrix}(2\pi)^4\delta^{(4)}(p-p')\\
&\quad - i2\pi|2p_0|^{1/2}\delta\big(p^2 - m^2_{\rm th}(\tilde{t}_f;\tilde{t}_i)\big)\tilde{f}_\chi(p,p',t)e^{i(p_0\,-\,p_0')\tilde{t}_f}2\pi|2p_0'|^{1/2}\delta\big(p'^2 - m^2_{\rm th}(\tilde{t}_f;\tilde{t}_i)\big)\begin{bmatrix}1 & 1\\ 1 & 1\end{bmatrix}\,.
\end{aligned}$$

If the subsystem $\mathscr{S}_\chi$ is in a state of thermodynamic equilibrium at the initial time $t = 0$, the thermal mass reduces to the known result

$$m^2_{\rm th}(t = 0) \;=\; \frac{\lambda T^2}{24}\;, \tag{9.55}$$

for $m = 0$ and $\mu \ll T$. In order to describe completely the dynamics of the combined system $\mathscr{S} = \mathscr{S}_\Phi \cup \mathscr{S}_\chi$, we couple the evolution of the thermal mass $m_{\rm th}(t)$ to the perturbative time evolution equations (9.41) and (9.44) for the Φ and χ statistical distribution functions. This is achieved by differentiating (9.52) with respect to t, such that

$$\partial_t m_{\rm th}(t) \;=\; \frac{\lambda}{2m_{\rm th}(t)}\int\frac{{\rm d}^3\mathbf{k}}{(2\pi)^3}\,\frac{1}{2E_\chi(\mathbf{k})}\,\frac{1}{2}\Big(\partial_t f_\chi(|\mathbf{k}|,t) + \partial_t f^C_\chi(|\mathbf{k}|,t)\Big)\;. \tag{9.56}$$

Here, $\partial_t f_\chi(|\mathbf{k}|,t)$ and $\partial_t f^C_\chi(|\mathbf{k}|,t)$ depend implicitly upon $m_{\rm th}(t)$ via the coupled first-order differential equations (9.41) and (9.44), derived using the quasi-particle χ propagators in (9.54). On dimensional grounds, we may estimate that the leading contribution to $\partial_t m_{\rm th}(t)$ is of order $\lambda^{1/2}gT$. The latter estimate provides firm support that our time evolution equations may consistently incorporate thermal masses and so describe the main non-perturbative dynamics of the system.

10 Conclusions

We have developed a new perturbative approach to non-equilibrium thermal quantum field theory. Our perturbative approach is based upon non-homogeneous free propagators and time-dependent vertices, which explicitly break space-time translational invariance and properly encode the dependence of the system on the time of observation. The forms of these propagators are constrained by fundamental field-theoretic requirements, such as CPT invariance of the action, Hermiticity properties of correlation functions, causality and unitarity. We have shown that our perturbative approach gives rise to time-dependent diagrammatic perturbation series, which are free of pinch singularities. The absence of these pinch singularities results from the systematic inclusion of finite-time effects and the proper consideration of the time of observation. We emphasize that this is achieved without invoking *ad hoc* prescriptions or effective resummations of finite widths. In our formalism, we have derived the *new* master time evolution equations (8.13) and (8.14) for particle number densities and statistical distribution functions, which are valid *to all orders* in perturbation theory and *to all orders* in gradient expansion. Furthermore, the master time evolution equations (8.13) and (8.14) respect CPT invariance and are also invariant under time translations of the CTP contour. As opposed to other methods, in our perturbative approach, we do not need to employ quasi-particle approximation and no assumption is necessary about the separation of time scales.

We have shown how the effect of a finite time interval since the start of evolution of the system leads to violation of energy conservation, as dictated by the Heisenberg uncertainty principle. Our approach permits the systematic treatment of the pertinent generalized kinematics in this evanescent regime. We have found that the available phase space of $1 \to 2$ decays increases and would-be kinematically disallowed processes can still take place at early times, contributing significantly to our time evolution equations. Within a simple scalar model, we have illustrated that these kinematically forbidden evanescent processes are the $2 \to 1$ processes of Landau damping, as well as $3 \to 0$ processes of *total annihilation* into the thermal bath. The processes of Landau damping and total annihilation have been shown to contribute promptly to the particle width, to a level as high as 20%. The switching on of the interactions leads to a sudden deposition of energy in the system, which manifests itself as a rapid change in both the particle width and the collision terms of the time evolution equations. These early-time effects give rise to an oscillating pattern, which persists even at later times. We have demonstrated that these late-time memory effects exhibit a non-Markovian evolution characterized by oscillations with time-dependent frequencies. The latter constitutes a distinctive feature of proper non-equilibrium dynamics, which is consistently predicted by our perturbative approach.

We note that the rapid transient behaviour of the system makes the method of gradient expansion unsuitable for early times. We emphasize that in our approach no assumption was made to the relative rate of thermalization of either the statistical or spectral behaviour of the system. For the considered initial conditions of thermal equilibrium, we have found that the spectral evolution resulting from evanescent contributions is critical to the early-time statistical dynamics of the system. Consequently, it would have been inappropriate to assume a separation of time scales at early times. A more accurate numerical solution to our time evolution equations turns out to be computationally intensive and may be presented elsewhere.

Finite-time effects, which are systematically incorporated in our perturbative approach, are also relevant to many-body systems whenever a natural characteristic time scale for perturbations arises in such systems. Thus, in addition to possible applications to reheating and preheating, of particular interest are first-order phase transitions. For instance, such characteristic time scales result from the bubble wall velocity of nucleation in the electroweak phase transition, e.g. see [78, 79]. In the vicinity of the bubble wall, evanescent contributions may weaken the constraints of decay and inverse decay thresholds and so affect the washout phenomena and the generation of relic densities. This evanescent regime is particularly relevant to prethermalization [80] and isotropization [81] time scales, which are known to be shorter than the time scale of thermalization.

It is straightforward to generalize our perturbative approach to theories that include fermions and gauge fields. Thus, it would be very interesting to extend the classical approaches [82, 83] to kinetic equations of particle mixing to non-homogeneous backgrounds, by including finite-time evanescent effects in line with the non-equilibrium formalism presented in this paper.

Acknowledgments

The authors thank Stephan Huber and Daniele Teresi for useful comments and discussions. The work of PM and AP is supported in part by the Lancaster–Manchester–Sheffield Consortium for Fundamental Physics under STFC grant ST/J000418/1. AP also acknowledges partial support by an IPPP associateship from Durham University.

A Propagator Properties and Identities

In this appendix, we give a detailed summary of the transformation properties and identities that relate the various propagators defined in Section 2. In detail, the pertinent two-point correlation functions for the complex scalar field χ are given by

$$\begin{aligned}
i\Delta(x,y) &\equiv \langle[\,\chi(x),\ \chi^\dagger(y)\,]\rangle\ , && \text{(A.1a)}\\
i\Delta_1(x,y) &\equiv \langle\{\,\chi(x),\ \chi^\dagger(y)\,\}\rangle\ , && \text{(A.1b)}\\
i\Delta_{\mathrm{R}}(x,y) &\equiv \theta(x_0\,-\,y_0)i\Delta(x,y)\ , && \text{(A.1c)}\\
i\Delta_{\mathrm{A}}(x,y) &\equiv -\theta(y_0\,-\,x_0)i\Delta(x,y)\ , && \text{(A.1d)}\\
i\Delta_{\mathcal{P}}(x,y) &\equiv \tfrac{1}{2}\varepsilon(x_0\,-\,y_0)\,\langle[\,\chi(x),\ \chi^\dagger(y)\,]\rangle\ , && \text{(A.1e)}\\
i\Delta_{>}(x,y) &\equiv \langle\chi(x)\chi^\dagger(y)\rangle\ , && \text{(A.1f)}\\
i\Delta_{<}(x,y) &\equiv \langle\chi^\dagger(y)\chi(x)\rangle\ , && \text{(A.1g)}\\
i\Delta_{\mathrm{F}}(x,y) &\equiv \langle\,\mathrm{T}\,[\,\chi(x)\chi^\dagger(y)\,]\rangle\ , && \text{(A.1h)}\\
i\Delta_{\mathrm{D}}(x,y) &\equiv \langle\,\bar{\mathrm{T}}\,[\,\chi(x)\chi^\dagger(y)\,]\rangle\ . && \text{(A.1i)}
\end{aligned}$$

The definitions of the charge-conjugate propagators follow from the unitary transformation

$$\begin{aligned}
U_C^\dagger\chi(x)U_C &= \chi^C(x) = \eta\chi^\dagger(x)\ , && \text{(A.2a)}\\
U_C^\dagger\chi^\dagger(x)U_C &= \chi^{C\dagger}(x) = \eta^*\chi(x)\ , && \text{(A.2b)}
\end{aligned}$$

where the complex phase η satisfies $|\eta|^2\,=\,1$.

It follows from the definitions that the propagators satisfy the transformations listed below under charge- and Hermitian-conjugation:

$$\begin{aligned}
\Delta(x,y) &= -\Delta^*(y,x) = -\Delta^C(y,x)\ , && \text{(A.3a)}\\
\Delta_1(x,y) &= -\Delta_1^*(y,x) = \Delta_1^C(y,x)\ , && \text{(A.3b)}\\
\Delta_{\mathcal{P}}(x,y) &= \Delta_{\mathcal{P}}^*(y,x) = \Delta_{\mathcal{P}}^C(y,x)\ , && \text{(A.3c)}\\
\Delta_{\mathrm{R}}(x,y) &= \Delta_{\mathrm{R}}^{C*}(x,y) = \Delta_{\mathrm{A}}^*(y,x) = \Delta_{\mathrm{A}}^C(y,x)\ , && \text{(A.3d)}\\
\Delta_{>}(x,y) &= -\Delta_{>}^*(y,x) = \Delta_{<}^C(y,x) = -\Delta_{<}^{C*}(x,y)\ , && \text{(A.3e)}\\
\Delta_{\mathrm{F}}(x,y) &= \Delta_{\mathrm{F}}^C(y,x) = -\Delta_{\mathrm{D}}^*(y,x) = -\Delta_{\mathrm{D}}^{C*}(x,y)\ , && \text{(A.3f)}
\end{aligned}$$

where the action of charge-conjugation is trivial in the case of a real scalar field. In the double momentum representation, these identities take the form:

$$\begin{aligned}
\Delta(p,p') &= -\Delta^*(p',p) = -\Delta^C(-p',-p)\ , && \text{(A.4a)}\\
\Delta_1(p,p') &= -\Delta_1^*(p',p) = \Delta_1^C(-p',-p)\ , && \text{(A.4b)}\\
\Delta_{\mathcal{P}}(p,p') &= \Delta_{\mathcal{P}}^*(p',p) = \Delta_{\mathcal{P}}^C(-p',-p)\ , && \text{(A.4c)}\\
\Delta_{\mathrm{R}}(p,p') &= \Delta_{\mathrm{R}}^{C*}(-p,-p') = \Delta_{\mathrm{A}}^*(p',p) = \Delta_{\mathrm{A}}^C(-p',-p)\ , && \text{(A.4d)}\\
\Delta_{>}(p,p') &= -\Delta_{>}^*(p',p) = \Delta_{<}^C(-p',-p) = -\Delta_{<}^{C*}(-p,-p')\ , && \text{(A.4e)}\\
\Delta_{\mathrm{F}}(p,p') &= \Delta_{\mathrm{F}}^C(-p',-p) = -\Delta_{\mathrm{D}}^*(p',p) = -\Delta_{\mathrm{D}}^{C*}(-p,-p')\ . && \text{(A.4f)}
\end{aligned}$$

Finally, in the Wigner representation, the propagators satisfy the following properties:

$$\begin{aligned}
\Delta(q,X) &= -\Delta^{*}(q,X) = -\Delta^{C}(-q,X)\ , && \text{(A.5a)}\\
\Delta_{1}(q,X) &= -\Delta_{1}^{*}(q,X) = \Delta_{1}^{C}(-q,X)\ , && \text{(A.5b)}\\
\Delta_{\mathrm{R}}(q,X) &= \Delta_{\mathrm{R}}^{C*}(-q,X) = \Delta_{\mathrm{A}}^{*}(q,X) = \Delta_{\mathrm{A}}^{C}(-q,X)\ , && \text{(A.5c)}\\
\Delta_{>}(q,X) &= -\Delta_{>}^{*}(q,X) = \Delta_{<}^{C}(-q,X) = -\Delta_{<}^{C*}(-q,X)\ , && \text{(A.5d)}\\
\Delta_{\mathrm{F}}(q,X) &= \Delta_{\mathrm{F}}^{C}(-q,X) = -\Delta_{\mathrm{D}}^{*}(q,X) = -\Delta_{\mathrm{D}}^{C*}(-q,X)\ . && \text{(A.5e)}
\end{aligned}$$

We also list the following set of useful identities:

$$\begin{aligned}
\Delta(x,y) &= \Delta_{>}(x,y) - \Delta_{<}(x,y) = \Delta_{\mathrm{R}}(x,y) - \Delta_{\mathrm{A}}(x,y)\ , && \text{(A.6a)}\\
&= \varepsilon(x_0 - y_0)\Big(\Delta_{\mathrm{F}}(x,y) - \Delta_{\mathrm{D}}(x,y)\Big)\ , && \text{(A.6b)}\\
\Delta_{1}(x,y) &= \Delta_{>}(x,y) + \Delta_{<}(x,y) = \Delta_{\mathrm{F}}(x,y) + \Delta_{\mathrm{D}}(x,y)\ , && \text{(A.6c)}\\
\Delta_{\mathrm{R(A)}}(x,y) &= \Delta_{\mathrm{F}}(x,y) - \Delta_{<(>)}(x,y) = -\Delta_{\mathrm{D}}(x,y) + \Delta_{>(<)}(x,y)\ , && \text{(A.6d)}\\
\Delta_{\mathrm{F(D)}}(x,y) &= \frac{1}{2}\Big((-)2\Delta_{\mathcal{P}}(x,y) + \Delta_{>}(x,y) + \Delta_{<}(x,y)\Big)\ , && \text{(A.6e)}\\
\Delta_{\mathcal{P}}(x,y) &= \frac{1}{2}\Big(\Delta_{\mathrm{R}}(x,y) + \Delta_{\mathrm{A}}(x,y)\Big) = \frac{1}{2}\Big(\Delta_{\mathrm{F}}(x,y) - \Delta_{\mathrm{D}}(x,y)\Big). && \text{(A.6f)}
\end{aligned}$$

We note that analogous relations hold for the corresponding self-energies and that these identities and relations are true for free and resummed propagators.

B Correspondence between Imaginary and Real Time Formalisms

In this appendix, we briefly outline a number of relevant details of the Imaginary Time Formalism (ITF) of thermal field theory. These details are discussed in the context of the real scalar field theory introduced in Section 2. Subsequently, we identify the correspondence of the ITF Φ-scalar propagator and its one-loop non-local self-energy with results calculated explicitly in real time, using the CTP formalism of Section 3 in the equilibrium limit discussed in Section 5.

The equilibrium density operator ρ_{eq} in (5.6) permits a path-integral representation in negative imaginary time. The ITF generating functional is

$$\mathcal{Z}[J] = \int[\mathrm{d}\Phi]\ \exp\Big(-\bar{S}[\Phi] + \int_0^{\beta}\mathrm{d}\tau_x\int\mathrm{d}^3\mathbf{x}\ J(\bar{x})\Phi(\bar{x})\Big)\ , \tag{B.1}$$

with action

$$\bar{S}[\Phi] = \int_0^{\beta}\mathrm{d}\tau_x\int\mathrm{d}^3\mathbf{x}\ \Big(\tfrac{1}{2}\partial_\mu\Phi(\bar{x})\partial_\mu\Phi(\bar{x}) + \tfrac{1}{2}M^2\Phi^2(\bar{x}) + \tfrac{1}{3!}g\Phi^3(\bar{x}) + \tfrac{1}{4!}\lambda\Phi^4(\bar{x})\Big)\ . \tag{B.2}$$

We emphasize the restricted domain of integration over the imaginary time $\tau_x \in [0,\ \beta]$. Four-dimensional Euclidean space-time coordinates are denoted by a horizontal bar, i.e. $\bar{x}_\mu \equiv (\tau_x, \mathbf{x})$. In the limit $\beta \to \infty$, (B.1) is precisely the Wick rotation to Euclidean space-time of the Minkowski-space generating functional via the analytic continuation $x_0 \to -i\tau_x$.

The free imaginary-time propagator $\bar{\Delta}^0$ may be written as follows:

$$\bar{\Delta}^0(\bar{x}-\bar{y}) \;=\; \frac{1}{\beta}\sum_{\ell\,=\,-\infty}^{+\infty}\int\frac{\mathrm{d}^3\mathbf{p}}{(2\pi)^3}\,e^{i[\omega_\ell(\tau_x\,-\,\tau_y)\,+\,\mathbf{p}\cdot(\mathbf{x}\,-\,\mathbf{y})]}\bar{\Delta}_0(i\omega_\ell,\mathbf{p})\;, \tag{B.3}$$

where

$$\bar{\Delta}^0(i\omega_\ell,\mathbf{p}) \;=\; \frac{1}{\omega_\ell^2\;+\;\mathbf{p}^2\;+\;M^2} \tag{B.4}$$

is the so-called Matsubara propagator. The discrete Matsubara frequencies $\omega_\ell \;=\; 2\pi\ell/\beta$, $\ell \;\in\; \mathbb{Z}$, arise from the periodicity of the imaginary time direction in order to satisfy the KMS relation [cf. (5.15)]

$$\bar{\Delta}^0(\bar{x}-\bar{y}) \;=\; \bar{\Delta}^0(\bar{x}-\bar{y}+\beta)\;. \tag{B.5}$$

The resummed Matsubara propagator is given by the imaginary-time Schwinger–Dyson equation

$$\bar{\Delta}^{-1}(i\omega_\ell,\mathbf{p}) \;=\; \bar{\Delta}^{0,\,-1}(i\omega_\ell,\mathbf{p})\;+\;\bar{\Pi}(i\omega_\ell,\mathbf{p})\;, \tag{B.6}$$

where $\bar{\Pi}(\omega_\ell,\mathbf{p})$ is the imaginary-time self-energy. Equation (B.6) may be inverted directly to obtain

$$\bar{\Delta}(i\omega_\ell,\mathbf{p}) \;=\; \frac{1}{\omega_\ell^2\;+\;\mathbf{p}^2\;+\;M^2\;+\;\bar{\Pi}(i\omega_\ell,\mathbf{p})}\;. \tag{B.7}$$

The free Matsubara propagator in (B.4) may also be written in the following spectral representation:

$$\bar{\Delta}^0(i\omega_\ell,\mathbf{p}) \;=\; -i\int\frac{\mathrm{d}k_0}{2\pi}\,\frac{\Delta^0(k_0,\mathbf{p})}{i\omega_\ell\;-\;k_0}\;, \tag{B.8}$$

where

$$i\Delta^0(k_0,\mathbf{p}) \;=\; 2\pi\varepsilon(k_0)\delta(k_0^2\;-\;\mathbf{p}^2\;-\;M^2) \tag{B.9}$$

is the single-momentum representation of the free Pauli–Jordan propagator, consistent with (2.16a). Making the analytic continuation $i\omega_\ell \to p_0 + i\epsilon$ to real frequencies and comparing with the spectral representation of the retarded propagator Δ_{R} in (2.22), we may establish the correspondence

$$\bar{\Delta}^0(i\omega_\ell \to p_0+i\epsilon,\mathbf{p}) \;=\; -\Delta_{\mathrm{R}}^0(p)\;. \tag{B.10}$$

This correspondence must also hold for the resummed Matsubara propagator $\bar{\Delta}$ via (B.6). As such, the analytic continuation of the ITF self-energy $\bar{\Pi}$,

$$\bar{\Pi}(i\omega_\ell \to p_0+i\epsilon,\mathbf{p}) \;=\; -\Pi_{\mathrm{R}}(p)\;, \tag{B.11}$$

yields the equilibrium retarded self-energy Π_{R}.

Thus, in thermodynamic equilibrium, an exact correspondence can be established between the ITF and the real-time approach of the CTP formalism, by means of *retarded* propagators and self-energies. The full complement of propagators exhibited in Table 2 and the corresponding self-energies may be obtained using the constraints of causality (2.24) and unitarity (2.28) in combination with the KMS relation (5.16) and the condition of detailed balance (5.25). The relationships between the retarded and CTP propagators and self-energies are listed explicitly in (5.19) and (5.26).

To illustrate the correspondence in (B.11), we consider the one-loop bubble diagram of the real scalar theory in (2.1). The real and imaginary parts of the retarded self-energy $\Pi_{\rm R}$ may be calculated in the CTP formalism from the time-ordered Π and positive-frequency Wightman $\Pi_>$ self-energies, respectively, using the relations in (5.26).

The real part of the one-loop retarded self-energy $\Pi^{(1)}_{\rm R}$ is given by

$$\mathrm{Re}\,\Pi^{(1)}_{\rm R}(p) \;=\; \mathrm{Re}\bigg[-\,i\frac{(-ig)^2}{2}\int\frac{\mathrm{d}^4k}{(2\pi)^4}\;i\Delta^0_{\rm F}(k)i\Delta^0_{\rm F}(p-k)\bigg]\;, \tag{B.12}$$

where $i\Delta^0_{\rm F}(k)$ is the equilibrium CTP Feynman propagator in (5.10a). Explicitly, we have

$$\begin{aligned}\mathrm{Re}\,\Pi^{(1)}_{\rm R}(p) \;=\; & -\frac{g^2}{2}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{1}{4E_1E_2} \\ & \times\bigg[\sum_{\alpha_1\,=\,\pm1}\;\int\limits_{(p-k)^2\neq M^2}\mathrm{d}k_0\,\frac{2E_2}{(p-k)^2-M^2}\,\delta\big(k_0\,-\,\alpha_1E_1\big)\Big(\tfrac{1}{2}\,+\,f_{\rm B}(k_0)\Big) \\ & +\sum_{\alpha_2\,=\,\pm1}\;\int\limits_{k^2\neq M^2}\mathrm{d}k_0\,\frac{2E_1}{k^2-M^2}\,\delta\big(p_0\,-\,k_0\,-\,\alpha_2E_2\big)\Big(\tfrac{1}{2}\,+\,f_{\rm B}(p_0-k_0)\Big)\bigg]\;,\end{aligned} \tag{B.13}$$

where $E_1\;\equiv\;E(\mathbf{k})$ and $E_2\;\equiv\;E(\mathbf{p}-\mathbf{k})$. The integral subscripts remind us that the integration around the on-shell poles are understood in the Cauchy principle value sense. Integration over k_0 yields the result

$$\mathrm{Re}\,\Pi^{(1)}_{\rm R}(p) \;=\; -\frac{g^2}{2}\sum_{\{\alpha\}}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{\alpha_1\alpha_2}{4E_1E_2}\,\frac{1\,+\,f_{\rm B}(\alpha_1E_1)\,+\,f_{\rm B}(\alpha_2E_2)}{p_0\,-\,\alpha_1E_1\,-\,\alpha_2E_2}\;, \tag{B.14}$$

where we have used the short-hand notation $\{\alpha\}$ to denote summation over $\alpha_1,\;\alpha_2\;=\;\pm1$.

The one-loop positive-frequency Wightman self-energy is given by

$$i\Pi^{(1)}_>(p) \;=\; \frac{(-ig)^2}{2}\int\frac{\mathrm{d}^4k}{(2\pi)^4}\;i\Delta^0_>(k)i\Delta^0_>(p-k)\;. \tag{B.15}$$

Using the signum form of the equilibrium positive-frequency Wightman propagator $i\Delta^0_>(k)$ in (5.10b), we obtain

$$\begin{aligned}\Pi^{(1)}_>(p) \;=\; & i\pi g^2\sum_{\{\alpha\}}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\int\mathrm{d}k_0\,\frac{\alpha_1\alpha_2}{4E_1E_2}\,\delta(k_0\,-\,\alpha_1E_1)\delta(p_0\,-\,k_0\,-\,\alpha_2E_2) \\ & \times\Big(1\,+\,f_{\rm B}(k_0)\Big)\Big(1\,+\,f_{\rm B}(p_0-k_0)\Big)\;.\end{aligned} \tag{B.16}$$

The product of Bose–Einstein distributions in (B.16) satisfies the following relation:

$$f_{\rm B}(k_0)f_{\rm B}(p_0-k_0) \;=\; f_{\rm B}(p_0)\Big(1\,+\,f_{\rm B}(k_0)\,+\,f_{\rm B}(p_0-k_0)\Big)\;. \tag{B.17}$$

Thus, upon integration over k_0, we find

$$\begin{aligned}\Pi^{(1)}_>(p) \;=\; & i\pi g^2\Big(1\,+\,f_{\rm B}(p_0)\Big)\sum_{\{\alpha\}}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{\alpha_1\alpha_2}{4E_1E_2}\,\delta(p_0\,-\,\alpha_1E_1\,-\,\alpha_2E_2) \\ & \times\Big(1\,+\,f_{\rm B}(\alpha_1E_1)\,+\,f_{\rm B}(\alpha_2E_2)\Big)\;.\end{aligned} \tag{B.18}$$

Using the relation in (5.26c), the imaginary part of the one-loop retarded self-energy may then be written down as

$$\begin{aligned}\mathrm{Im}\,\Pi^{(1)}_{\mathrm{R}}(p) \;&=\; \frac{\pi g^2}{2}\sum_{\{\alpha\}}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{\alpha_1\alpha_2}{4E_1E_2}\,\delta(p_0\,-\,\alpha_1E_1\,-\,\alpha_2E_2)\\ &\quad\times\,\Big(1\,+\,f_{\mathrm{B}}(\alpha_1E_1)\,+\,f_{\mathrm{B}}(\alpha_2E_2)\Big)\;. \qquad\qquad (\mathrm{B}.19)\end{aligned}$$

In the ITF, the one-loop self-energy is given by

$$-\,\bar{\Pi}^{(1)}(i\omega_\ell,\mathbf{p})\;=\;\frac{(-g)^2}{2\beta}\sum_{n\,=\,-\infty}^{+\infty}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\bar{\Delta}(i\omega_n,\mathbf{k})\bar{\Delta}(i(\omega_\ell-\omega_n),\mathbf{p}-\mathbf{k})\;. \qquad (\mathrm{B}.20)$$

After performing the summation over n (see for instance [68]), we obtain

$$\bar{\Pi}^{(1)}(i\omega_\ell,\mathbf{p})\;=\;\frac{g^2}{2}\sum_{\{\alpha\}}\int\frac{\mathrm{d}^3\mathbf{k}}{(2\pi)^3}\,\frac{\alpha_1\alpha_2}{4E_1E_2}\,\frac{1\,+\,f_{\mathrm{B}}(\alpha_1E_1)\,+\,f_{\mathrm{B}}(\alpha_2E_2)}{i\omega_\ell\,-\,\alpha_1E_1\,-\,\alpha_2E_2}\;. \qquad (\mathrm{B}.21)$$

Making the analytic continuation $i\omega_\ell \to p_0 + i\epsilon$ and subsequently extracting the real and imaginary parts, the result in (B.21) agrees with (B.14) and (B.19) via the correspondence in (B.11).

C The Complex Scalar Field

This appendix describes the generalization of the results in Sections 4.3 and 5 to the case of a complex scalar field χ. In particular, we derive the full complement of non-homogeneous free propagators and expand upon the ITF correspondence identified in Appendix B.

Our starting point is the complex-scalar Lagrangian

$$\mathcal{L}(x)\;=\;\partial_\mu\chi^\dagger(x)\partial^\mu\chi(x)\;-\;m^2\chi^\dagger(x)\chi(x)\;-\;\tfrac{1}{4}\lambda\big[\chi^\dagger(x)\chi(x)\big]^2\;. \qquad (\mathrm{C}.1)$$

In analogy to (2.11), the complex scalar field $\chi(x)$ may be written in the interaction picture as

$$\chi(x;\tilde{t}_i)\;=\;\int\mathrm{d}\Pi_{\mathbf{p}}\,\Big(a(\mathbf{p},0;\tilde{t}_i)e^{-iE(\mathbf{p})x_0}e^{i\mathbf{p}\cdot\mathbf{x}}\;+\;b^\dagger(\mathbf{p},0;\tilde{t}_i)e^{iE(\mathbf{p})x}e^{-i\mathbf{p}\cdot\mathbf{x}}\Big)\;, \qquad (\mathrm{C}.2)$$

where $a^\dagger(\mathbf{p},0;\tilde{t}_i)$ and $a(\mathbf{p},0;\tilde{t}_i)$ ($b^\dagger(\mathbf{p},0;\tilde{t}_i)$ and $b(\mathbf{p},0;\tilde{t}_i)$) are the interaction-picture particle (anti-particle) creation and annihilation operators, respectively. Under C-conjugation [cf. (A.2a)] these creation and annihilation operators satisfy the transformations

$$U_C^\dagger\,a(\mathbf{p},\tilde{t};\tilde{t}_i)U_C\;=\;\eta b(\mathbf{p},\tilde{t};\tilde{t}_i)\;,\qquad U_C^\dagger\,b^\dagger(\mathbf{p},\tilde{t};\tilde{t}_i)U_C\;=\;\eta a^\dagger(\mathbf{p},\tilde{t};\tilde{t}_i)\;. \qquad (\mathrm{C}.3)$$

Introducing the four-dimensional LIPS measure from (2.3), (C.2) and its Hermitian conjugate may be recast in the form

$$\chi(x;\tilde{t}_i)\;=\;\int\frac{\mathrm{d}^4p}{(2\pi)^4}\,e^{-ip\cdot x}\,\chi(p;\tilde{t}_i)\;,\qquad \chi^\dagger(x;\tilde{t}_i)\;=\;\int\frac{\mathrm{d}^4p}{(2\pi)^4}\,e^{-ip\cdot x}\,\chi^\dagger(-p;\tilde{t}_i)\;, \qquad (\mathrm{C}.4)$$

where the Fourier amplitudes are given by

$$\chi(p;\tilde{t}_i) \;=\; 2\pi\delta(p^2 \,-\, m^2)\Big(\,\theta(p_0)a(\mathbf{p},0;\tilde{t}_i) \,+\, \theta(-p_0)b^\dagger(-\mathbf{p},0;\tilde{t}_i)\,\Big)\;. \tag{C.5}$$

For the real scalar field, the quantization scheme was only depending on the restriction placed upon the form of the field commutator. In the case of the complex scalar field, we have two degrees of freedom to fix. Thus, we begin with the following two commutators of interaction-picture fields:

$$\big[\,\chi(x),\;\chi(y)\,\big] \;=\; 0\;, \qquad \big[\,\chi(x),\;\chi^\dagger(y)\,\big] \;=\; i\Delta^0(x,y;m^2)\;, \tag{C.6}$$

where the Pauli–Jordan propagator has precisely the form in (2.15). In analogy to the real scalar field, we may derive from (C.6) the equal-time commutation relations

$$i\Delta^0(x,y;m^2)\big|_{x^0\,=\,y^0\,=\,\tilde{t}} \;=\; \big[\,\chi(\tilde{t},\mathbf{x}),\;\chi^\dagger(\tilde{t},\mathbf{y})\,\big] \;=\; 0\;, \tag{C.7a}$$

$$\partial_{x_0} i\Delta^0(x,y;m^2)\big|_{x^0\,=\,y^0\,=\,\tilde{t}} \;=\; \big[\,\pi^\dagger(\tilde{t},\mathbf{x}),\;\chi^\dagger(\tilde{t},\mathbf{y})\,\big] \;=\; -i\delta^{(3)}(\mathbf{x}\,-\,\mathbf{y})\;, \tag{C.7b}$$

$$\partial_{y_0} i\Delta^0(x,y;m^2)\big|_{x^0\,=\,y^0\,=\,\tilde{t}} \;=\; \big[\,\chi(\tilde{t},\mathbf{x}),\;\pi(\tilde{t},\mathbf{y})\,\big] \;=\; i\delta^{(3)}(\mathbf{x}\,-\,\mathbf{y})\;, \tag{C.7c}$$

$$\partial_{x_0}\partial_{y_0} i\Delta^0(x,y;m^2)\big|_{x^0\,=\,y^0\,=\,\tilde{t}} \;=\; \big[\,\pi^\dagger(\tilde{t},\mathbf{x}),\;\pi(t,\mathbf{y})\,\big] \;=\; 0\;, \tag{C.7d}$$

where $\pi(x) = \partial_{x_0}\chi^\dagger(x)$ is the conjugate momentum operator. The particle and anti-particle creation and annihilation operators necessarily satisfy the algebra

$$\big[\,a(\mathbf{p},\tilde{t}),\;a^\dagger(\mathbf{p}',\tilde{t}')\,\big] \;=\; \big[\,b(\mathbf{p},\tilde{t}),\;b^\dagger(\mathbf{p}',\tilde{t}')\,\big] \;=\; (2\pi)^3 2E(\mathbf{p})\delta^{(3)}(\mathbf{p}-\mathbf{p}')e^{-iE(\mathbf{p})(\tilde{t}\,-\,\tilde{t}')}\;, \tag{C.8}$$

with all other commutators vanishing. We stress again that the phase factor $e^{-iE(\mathbf{p})(\tilde{t}\,-\,\tilde{t}')}$ on the RHS of (C.8) has appeared due to the difference in microscopic times of the interaction-picture creation and annihilation operators.

In addition to the C-conserving propagators listed in Appendix A, we may also define C-violating propagators. As an example, the C-violating Hadamard propagator $\Delta_{1\not{C}}(x,y)$ would read:

$$i\Delta_{1\not{C}}(x,y) \;=\; \langle\,\{\,\chi(x),\;\chi(y)\,\}\,\rangle\;, \tag{C.9}$$

which satisfies

$$\Delta_{1\not{C}}(x,y) \;=\; \Delta_{1\not{C}}(y,x) \;=\; -\eta^2\Delta^{C*}_{1\not{C}}(x,y)\;. \tag{C.10}$$

This Hadamard correlation function may, in general, be non-zero for early times, thus permitting extra C-violating evanescent processes in addition to those described in Section 9.

In analogy to (4.53), we write the following set of EEVs of two-point products of particle and anti-particle creation and annihilation operators:

$$\langle a^\dagger(\mathbf{p}',\tilde{t}_f;\tilde{t}_i)a(\mathbf{p},\tilde{t}_f;\tilde{t}_i)\rangle_t \;=\; 2\mathscr{E}(\mathbf{p},\mathbf{p}')f(\mathbf{p},\mathbf{p}',t)\;, \tag{C.11a}$$

$$\langle b(\mathbf{p}',\tilde{t}_f;\tilde{t}_i)a(\mathbf{p},\tilde{t}_f;\tilde{t}_i)\rangle_t \;=\; 2\mathscr{E}(\mathbf{p},\mathbf{p}')g(\mathbf{p},\mathbf{p}',t)\;, \tag{C.11b}$$

$$\langle a(\mathbf{p}',\tilde{t}_f;\tilde{t}_i)a(\mathbf{p},\tilde{t}_f;\tilde{t}_i)\rangle_t \;=\; 2\mathscr{E}(\mathbf{p},\mathbf{p}')h(\mathbf{p},\mathbf{p}',t)\;, \tag{C.11c}$$

$$\langle b^\dagger(\mathbf{p}',\tilde{t}_f;\tilde{t}_i)a(\mathbf{p},\tilde{t}_f;\tilde{t}_i)\rangle_t \;=\; 2\mathscr{E}(\mathbf{p},\mathbf{p}')d(\mathbf{p},\mathbf{p}',t)\;, \tag{C.11d}$$

where the remaining EEVs are obtained by Hermitian and charge conjugation. The four statistical distribution functions $f,\ g,\ h,\ d$ satisfy the following identities:

$$\begin{aligned} f(\mathbf{p},\mathbf{p}',t) &= f^{*}(\mathbf{p}',\mathbf{p},t)\ , && \text{(C.12a)}\\ g(\mathbf{p},\mathbf{p}',t) &= g^{C}(\mathbf{p}',\mathbf{p},t)\ , && \text{(C.12b)}\\ h(\mathbf{p},\mathbf{p}',t) &= h(\mathbf{p}',\mathbf{p},t)\ , && \text{(C.12c)}\\ d(\mathbf{p},\mathbf{p}',t) &= \eta^{2}d^{C*}(\mathbf{p}',\mathbf{p},t)\ . && \text{(C.12d)} \end{aligned}$$

The non-homogeneous free propagators of the complex scalar field χ may be written as listed in Table 2 with the substitution of the following ensemble function:

$$\begin{aligned} \tilde{f}(p,p',t) \ &= \ \theta(p_0)\theta(p'_0)f(\mathbf{p},\mathbf{p}',t) \ + \ \theta(-p_0)\theta(-p'_0)f^{C*}(-\mathbf{p},-\mathbf{p}',t)\\ &\quad + \ \theta(p_0)\theta(-p'_0)g(\mathbf{p},-\mathbf{p}',t) \ + \ \theta(-p_0)\theta(p'_0)g^{C*}(-\mathbf{p},\mathbf{p}',t)\ , \end{aligned} \tag{C.13}$$

satisfying the relations: $\tilde{f}(p,p',t) \ = \ \tilde{f}^{C}(-p',-p,t) \ = \ \tilde{f}^{C*}(-p,-p',t)$ in accordance with (A.4). The free C-violating Hadamard propagator $\Delta^{0}_{1\not{C}}(p,p',\tilde{t}_f;\tilde{t}_i)$, defined in (C.9), may be written down as

$$\Delta_{1\not{C}}(p,p',\tilde{t}_f;\tilde{t}_i) \ = \ -i2\pi\delta(p^2 \ - \ m^2)|2p_0|^{1/2}2\tilde{d}(p,p',t)e^{i(p_0\,-\,p'_0)\tilde{t}_f}|2p'_0|^{1/2}2\pi\delta(p'^2 \ - \ m^2)\ , \tag{C.14}$$

where we have defined the C-violating ensemble function $\tilde{d}(p,p',t)$. Evaluating the EEVs directly, we find

$$\begin{aligned} \tilde{d}(p,p',t) \ &= \ \theta(p_0)\theta(p'_0)d(\mathbf{p},\mathbf{p}',t) \ + \ \theta(-p_0)\theta(-p'_0)\eta^2 d^{C*}(-\mathbf{p},-\mathbf{p}',t)\\ &\quad + \ \theta(p_0)\theta(-p'_0)h(\mathbf{p},-\mathbf{p}',t) \ + \ \theta(-p_0)\theta(p'_0)\eta^2 h^{C*}(-\mathbf{p},\mathbf{p}',t)\ . \end{aligned} \tag{C.15}$$

satisfying the relations: $\tilde{d}(p,p',t) \ = \ \tilde{d}^{C}(-p',-p,t) \ = \ \eta^2\tilde{d}^{C*}(-p,-p',t)$.

The inclusion of the C-violating distribution functions requires the addition of the C-violating source l_{ab} to the expansion of the kernel of the density operator in the CTP generating functional, according to our discussion in Section 4.1. Omitting the t-dependence of the sources for notational convenience, the CTP generating functional for the complex scalar field χ takes the form

$$\begin{aligned} \mathcal{W}[j_a,k_{ab},l_{ab}] \ &= \ -i\hbar\ln\iint[\mathrm{d}(\chi^{a\dagger},\chi^{a})]\\ &\quad\times\ \exp\bigg\{\frac{i}{\hbar}\bigg[S\big[\chi^{a\dagger},\chi^{a}\big] \ + \ \int_{\Omega_t}\mathrm{d}^4x\ \Big(j^{\dagger}_a(x)\chi^a(x) \ + \ \chi^{\dagger}_a(x)j^a(x)\Big)\\ &\quad + \ \frac{1}{2}\iint_{\Omega_t}\mathrm{d}^4x\,\mathrm{d}^4x'\ \Big(\chi^{a\dagger}(x)k_{ab}(x,x')\chi^b(x') \ + \ \chi^a(x)k^{C}_{ab}(x,x')\chi^{b\dagger}(x')\\ &\quad + \ \chi^a(x)l^{\dagger}_{ab}(x,x')\chi^b(x') \ + \ \chi^{a\dagger}(x)l_{ab}(x,x')\chi^{b\dagger}(x')\Big) \ + \ \cdots\bigg]\bigg\}\ . \end{aligned} \tag{C.16}$$

The bi-local sources k_{ab} and l_{ab} necessarily satisfy the identities

$$k_{ab}(x,x') \ = \ k^{\dagger}_{ba}(x',x)\ , \qquad l_{ab}(x,x') \ = \ l_{ba}(x',x) \ = \ \eta^2 l^{C\dagger}_{ab}(x,x')\ , \tag{C.17}$$

to ensure that the exponent of the generating functional is CPT-invariant. The subsequent derivation of the effective action then follows analogously to Section 4.

The Lagrangian in (C.1) is invariant under the global $U(1)$ transformation

$$\chi(x) \;\to\; \chi'(x) \;=\; e^{-i\alpha}\,\chi(x)\;, \qquad \chi^\dagger(x) \;\to\; \chi'^\dagger(x) \;=\; e^{i\alpha}\,\chi^\dagger(x)\;, \tag{C.18}$$

entailing the conserved Noether current

$$j_\mu(x) \;=\; i\Big(\,\chi^\dagger(x)\partial_\mu\chi(x) \;-\; \big(\partial_\mu\chi^\dagger(x)\big)\chi(x)\,\Big)\;, \tag{C.19}$$

with corresponding conserved charge

$$:\mathcal{Q}(x_0): \;=\; \int \mathrm{d}^3\mathbf{x}\;\; :j_0(x): \;=\; \int \mathrm{d}\Pi_{\mathbf{p}}\;\Big(\,a^\dagger(\mathbf{p},0)a(\mathbf{p},0) \;-\; b^\dagger(\mathbf{p},0)b(\mathbf{p},0)\,\Big)\;, \tag{C.20}$$

where : : denotes normal ordering. The existence of this conserved charge necessitates the introduction of a chemical potential μ and, as such, the equilibrium density operator is of the grand-canonical form

$$\rho(\beta,\mu) \;=\; e^{-\beta(H-\mu Q)}\;. \tag{C.21}$$

In the presence of this chemical potential, the KMS relation in (5.15) generalizes to

$$\Delta_>(x^0-y^0,\mathbf{x}-\mathbf{y}) \;=\; e^{-\beta\mu}\Delta_<(x^0-y^0+i\beta,\mathbf{x}-\mathbf{y}) \tag{C.22}$$

or, in the momentum representation,

$$\Delta_>(p) \;=\; e^{\beta(p_0-\mu)}\Delta_<(p)\;. \tag{C.23}$$

Proceeding as in Section 3, we find that the final constraint on $\tilde{f}(p)$, generalising (5.17), is

$$\tilde{f}(p) \;=\; \theta(p_0)f_{\mathrm{B}}(p_0) \;+\; \theta(-p_0)f_{\mathrm{B}}^{C}(-p_0)\;, \tag{C.24}$$

where $f_{\mathrm{B}}^{(C)}(p_0) = (e^{\beta[p_0-(+)\mu]}-1)^{-1}$ is the particle (anti-particle) Bose–Einstein distribution function. In equilibrium, translational invariance is restored and the elements of the free CTP propagator may be written in the single momentum representations

$$\begin{aligned} i\Delta_{\mathrm{F}}^0(p) \;&=\; i\Big(\,p^2 \;-\; m^2 \;+\; i\varepsilon\,\Big)^{-1} \\ &\qquad +\; 2\pi\,\Big(\,\theta(p_0)f_{\mathrm{B}}(p_0) \;+\; \theta(-p_0)f_{\mathrm{B}}^{C}(-p_0)\,\Big)\delta(p^2 \;-\; m^2)\;, \end{aligned} \tag{C.25a}$$

$$i\Delta_>^0(p) \;=\; 2\pi\Big[\,\theta(p_0)\Big(\,1 \;+\; f_{\mathrm{B}}(p_0)\,\Big) \;+\; \theta(-p_0)f_{\mathrm{B}}^{C}(-p_0)\,\Big]\delta(p^2 \;-\; m^2)\;, \tag{C.25b}$$

$$i\Delta_<^0(p) \;=\; 2\pi\Big[\,\theta(p_0)f_{\mathrm{B}}(p_0) \;+\; \theta(-p_0)\Big(\,1 \;+\; f_{\mathrm{B}}^{C}(-p_0)\,\Big)\Big]\delta(p^2 \;-\; m^2)\;. \tag{C.25c}$$

In order to define the ITF generating functional for the grand canonical partition function in (C.21), one may consider the Hamiltonian form of the path integral directly (see for instance [68]). We may write

$$\begin{aligned} \mathcal{Z}[j] \;=\; \iint [\mathrm{d}(\pi^\dagger,\pi)]\,[\mathrm{d}(\chi^\dagger,\chi)]\;\exp\bigg\{ &-\int_0^\beta \mathrm{d}\tau_x \int \mathrm{d}^3\mathbf{x}\;\Big[\,\mathcal{H}(\pi^{(\dagger)},\chi^{(\dagger)}) \\ &-\; i\,\Big(\,\pi(\bar{x})\partial_{\tau_x}\chi(\bar{x}) \;+\; \pi^\dagger(\bar{x})\partial_{\tau_x}\chi^\dagger(\bar{x})\Big) \;-\; i\mu\,\Big(\,\pi^\dagger(\bar{x})\chi^\dagger(\bar{x}) \;-\; \pi(\bar{x})\chi(\bar{x})\,\Big) \\ &-\; j^\dagger(\bar{x})\chi(\bar{x}) - j(\bar{x})\chi(\bar{x})\,\Big]\bigg\}\;. \end{aligned} \tag{C.26}$$

Expanding the fields and conjugate momenta in terms of two real degrees of freedom $\chi_{1,2}(\bar{x})$ and $\pi_{1,2}(\bar{x})$ as

$$\chi(\bar{x}) \;=\; \frac{1}{\sqrt{2}}\Big(\chi_1(\bar{x}) \,+\, i\chi_2(\bar{x})\Big)\,, \qquad \pi(\bar{x}) \;=\; \frac{1}{\sqrt{2}}\Big(\pi_1(\bar{x}) \,-\, i\pi_2(\bar{x})\Big)\,, \tag{C.27}$$

we may analytically calculate the Gaussian integrals over $\pi_{1,2}(\bar{x})$, yielding

$$\begin{aligned}\mathcal{Z}[j] \;=\; \int[\mathrm{d}(\chi^\dagger,\chi)]\,\exp\bigg[-\int_0^\beta \mathrm{d}\tau_x\int \mathrm{d}^3\mathbf{x}\,\Big((\partial_{\tau_x}+\mu)\chi^\dagger(\bar{x})(\partial_{\tau_x}-\mu)\chi(\bar{x})\\ +\,\nabla\chi^\dagger(\bar{x})\cdot\nabla\chi(\bar{x}) \,+\, m^2\chi^\dagger(\bar{x})\chi(\bar{x}) \,-\, j^\dagger(\bar{x})\chi(\bar{x}) \,-\, j(\bar{x})\chi^\dagger(\bar{x})\Big)\bigg]\,.\end{aligned} \tag{C.28}$$

In order to derive the form of the ITF χ-propagator, we insert into (C.28) the Fourier transform

$$\chi(\bar{x}) \;=\; \frac{1}{\beta}\sum_{\ell\,=\,-\infty}^{+\infty}\int\frac{\mathrm{d}^3\mathbf{p}}{(2\pi)^3}\,e^{i[\omega_\ell\tau_x\,+\,\mathbf{p}\cdot\mathbf{x}]}\chi(i\omega_\ell,\mathbf{p})\,, \tag{C.29}$$

where ω_ℓ are the discrete Matsubara frequencies described in Appendix B. The effect of the chemical potential is to shift the poles of the Matsubara propagator, which becomes

$$\bar{\Delta}^0(i\omega_\ell-\mu,\mathbf{p}) \;=\; \frac{1}{(\omega_\ell\,+\,i\mu)^2\,+\,\mathbf{p}^2\,+\,m^2}\,. \tag{C.30}$$

In generalization of the correspondence in (B.10), the equilibrium retarded propagator Δ_{R} is obtained by the analytic continuation $i\omega_\ell\,-\,\mu\,\to\,p_0\,+\,i\epsilon$ via

$$\bar{\Delta}^0(i\omega_\ell-\mu\to p_0+i\epsilon,\mathbf{p}) \;=\; -\Delta^0_{\mathrm{R}}(p)\,. \tag{C.31}$$

Correspondingly, the equilibrium retarded self-energy Π_{R} is given by

$$\bar{\Pi}(i\omega_\ell-\mu\to p_0+i\epsilon) \;=\; -\Pi_{\mathrm{R}}(p)\,, \tag{C.32}$$

thereby generalising (B.11).

D Non-Homogeneous Density Operator

In (D.1), we highlight the full form of the series expansion of the general Gaussian-like density operator described in Section 4.3. Symmetric and asymmetric multi-particle states are built up by summing over all possible convolutions of the W amplitudes. To facilitate our presentation task, the time dependence and phase-space integrals have been omitted. In fact, all momenta are integrated with the LIPS measure given in (2.3).

$$
\begin{aligned}
\rho \;=\; & \Big(1 + \frac{1}{2}W_{10}(\mathbf{p}_1:0)W_{01}(0:\mathbf{p}_1) + \frac{1}{4}W_{20}(\mathbf{p}_1,\mathbf{p}_2:0)W_{02}(0:\mathbf{p}_2,\mathbf{p}_1) + \cdots\Big)\Big\{\, |0\rangle\langle 0| \\
& + \Big(- W_{10}(\mathbf{k}_1:0) + \frac{1}{2}W_{11}(\mathbf{k}_1:\mathbf{q}_1)W_{10}(\mathbf{q}_1:0) \\
& \qquad + \frac{1}{2}W_{20}(\mathbf{k}_1,\mathbf{q}_1:0)W_{01}(0:\mathbf{q}_1) + \cdots\Big)|\mathbf{k}_1\rangle\langle 0| \\
& + \Big(- W_{01}(0:\mathbf{k}'_1) + \frac{1}{2}W_{01}(0:\mathbf{q}_1)W_{11}(\mathbf{q}_1:\mathbf{k}'_1) \\
& \qquad + \frac{1}{2}W_{02}(0:\mathbf{k}'_1,\mathbf{q}_1)W_{10}(\mathbf{q}_1:0) + \cdots\Big)|0\rangle\langle \mathbf{k}'_1| \\
& + \Big((2\pi)^3 2E(\mathbf{k}_1)\delta^{(3)}(\mathbf{k}_1 - \mathbf{k}'_1) - W_{11}(\mathbf{k}_1:\mathbf{k}'_1) + W_{10}(\mathbf{k}_1:0)W_{01}(0:\mathbf{k}'_1) \\
& \qquad + \frac{1}{2}W_{11}(\mathbf{k}_1:\mathbf{q}_1)W_{11}(\mathbf{q}_1:\mathbf{k}'_1) \\
& \qquad + \frac{1}{4}W_{20}(\mathbf{k}_1,\mathbf{q}_1:0)W_{02}(0:\mathbf{q}_1,\mathbf{k}'_1) + \cdots\Big)|\mathbf{k}_1\rangle\langle \mathbf{k}'_1| \\
& + \frac{1}{2}\Big(- W_{20}(\mathbf{k}_1,\mathbf{k}_2:0) + W_{10}(\mathbf{k}_1:0)W_{10}(\mathbf{k}_2:0) + \frac{1}{2}W_{11}(\mathbf{k}_1:\mathbf{q}_1)W_{20}(\mathbf{q}_1,\mathbf{k}_2:0) \\
& \qquad + \frac{1}{2}W_{11}(\mathbf{k}_2:\mathbf{q}_1)W_{20}(\mathbf{q}_1,\mathbf{k}_1:0) + \cdots\Big)|\mathbf{k}_1,\mathbf{k}_2\rangle\langle 0| \\
& + \frac{1}{2}\Big(- W_{02}(0:\mathbf{k}'_1,\mathbf{k}'_2) + W_{01}(0:\mathbf{k}'_1)W_{01}(0:\mathbf{k}'_2) + \frac{1}{2}W_{02}(0:\mathbf{k}'_1,\mathbf{q}_1)W_{11}(\mathbf{q}_1:\mathbf{k}'_2) \\
& \qquad + \frac{1}{2}W_{02}(0:\mathbf{k}'_2,\mathbf{q}_1)W_{11}(\mathbf{q}_1:\mathbf{k}'_1) + \cdots\Big)|0\rangle\langle \mathbf{k}'_1,\mathbf{k}'_2| \\
& + \frac{1}{2}\Big(W_{11}(\mathbf{k}_1:\mathbf{k}'_1)W_{10}(\mathbf{k}_2:0) + W_{11}(\mathbf{k}_2:\mathbf{k}'_1)W_{10}(\mathbf{k}_1:0) \\
& \qquad + W_{20}(\mathbf{k}_1,\mathbf{k}_2:0)W_{01}(0:\mathbf{k}'_1) + \cdots\Big)|\mathbf{k}_1,\mathbf{k}_2\rangle\langle \mathbf{k}'_1| \\
& + \frac{1}{2}\Big(W_{11}(\mathbf{k}_1:\mathbf{k}'_1)W_{01}(0:\mathbf{k}'_2) + W_{11}(\mathbf{k}_1:\mathbf{k}'_2)W_{01}(0:\mathbf{k}'_1) \\
& \qquad + W_{02}(0:\mathbf{k}'_1,\mathbf{k}'_2)W_{10}(\mathbf{k}_1:0) + \cdots\Big)|\mathbf{k}_1\rangle\langle \mathbf{k}'_1,\mathbf{k}'_2| \\
& + \frac{1}{2}\Big(W_{20}(\mathbf{k}_1,\mathbf{k}_2:0)W_{10}(\mathbf{k}_3:0) + \cdots\Big)|\mathbf{k}_1,\mathbf{k}_2,\mathbf{k}_3\rangle\langle 0| \\
& + \frac{1}{2}\Big(W_{02}(0:\mathbf{k}'_1,\mathbf{k}'_2)W_{01}(0:\mathbf{k}'_3) + \cdots\Big)|0\rangle\langle \mathbf{k}'_1,\mathbf{k}'_2,\mathbf{k}'_3| \\
& + \Big[\frac{1}{2}\Big((2\pi)^3 2E(\mathbf{k}_1)\delta^{(3)}(\mathbf{k}_1 - \mathbf{k}'_1) - W_{11}(\mathbf{k}_1:\mathbf{k}'_1) + W_{10}(\mathbf{k}_1:0)W_{01}(0:\mathbf{k}'_1) \\
& \qquad\qquad + \frac{1}{2}W_{11}(\mathbf{k}_1:\mathbf{q}_1)W_{11}(\mathbf{q}_1:\mathbf{k}'_1) + \frac{1}{4}W_{20}(\mathbf{k}_1,\mathbf{q}_1:0)W_{02}(0:\mathbf{q}_1,\mathbf{k}'_1) + \cdots\Big) \\
& \qquad \times \Big((2\pi)^3 2E(\mathbf{k}_2)\delta^{(3)}(\mathbf{k}_2 - \mathbf{k}'_2) - W_{11}(\mathbf{k}_2:\mathbf{k}'_2) + W_{10}(\mathbf{k}_2:0)W_{01}(0:\mathbf{k}'_2) \\
& \qquad\qquad + \frac{1}{2}W_{11}(\mathbf{k}_2:\mathbf{q}_2)W_{11}(\mathbf{q}_2:\mathbf{k}'_2) + \frac{1}{4}W_{20}(\mathbf{k}_2,\mathbf{q}_2:0)W_{02}(0:\mathbf{q}_2,\mathbf{k}'_2) + \cdots\Big) \\
& \quad + \frac{1}{4}\Big(- W_{20}(\mathbf{k}_1,\mathbf{k}_2:0) + W_{10}(\mathbf{k}_1:0)W_{10}(\mathbf{k}_2:0) + \frac{1}{2}W_{11}(\mathbf{k}_1:\mathbf{q}_1)W_{20}(\mathbf{q}_1,\mathbf{k}_2:0) \\
& \qquad\qquad + \frac{1}{2}W_{11}(\mathbf{k}_2:\mathbf{q}_1)W_{20}(\mathbf{q}_1,\mathbf{k}_1:0) + \cdots\Big) \\
& \qquad \times \Big(- W_{02}(0:\mathbf{k}'_1,\mathbf{k}'_2) + W_{01}(0:\mathbf{k}'_1)W_{01}(0:\mathbf{k}'_2) + \frac{1}{2}W_{02}(0:\mathbf{k}'_1,\mathbf{q}_1)W_{11}(\mathbf{q}_1:\mathbf{k}'_2) \\
& \qquad\qquad + \frac{1}{2}W_{02}(0:\mathbf{k}'_2,\mathbf{q}_1)W_{11}(\mathbf{q}_1:\mathbf{k}'_1) + \cdots\Big)\Big]\,|\mathbf{k}_1,\mathbf{k}_2\rangle\langle \mathbf{k}'_1,\mathbf{k}'_2| + \cdots\Big\}\,. \qquad \text{(D.1)}
\end{aligned}
$$

E Kadanoff–Baym Equations

In this appendix, we outline the derivation of the Kadanoff–Baym equations [52, 53]. The resulting time evolution equations are included here for comparison with the new results described in Sections 8 and 9. The Kadanoff–Baym equations are obtained by the partial inversion and subsequent gradient expansion of the Schwinger–Dyson equation derived in Section 4.1; see also our discussion in Section 4.2. We omit the $\tilde{t}_f$- and $\tilde{t}_i$-dependence of all propagators and self-energies for notational convenience.

Inverse Fourier transforming (8.5), we obtain the coordinate-space representation of the partially inverted Schwinger–Dyson equation

$$\begin{aligned}-\Big(\Box_x^2\,+\,M^2\Big)\Delta_{\gtrless}(x,y)\,+\int_{\Omega_t}\mathrm{d}^4z\,\Big(\Pi_{\gtrless}(x,z)\Delta_{\mathcal{P}}(z,y)\,+\,\Pi_{\mathcal{P}}(x,z)\Delta_{\gtrless}(z,y)\Big)\\ =\,\frac{1}{2}\int_{\Omega_t}\mathrm{d}^4z\,\Big(\Pi_{<}(x,z)\Delta_{>}(z,y)\,-\,\Pi_{>}(x,z)\Delta_{<}(z,y)\Big)\,,\end{aligned}\tag{E.1}$$

where we have also included the positive-frequency contribution for completeness. Introducing the relative and central coordinates R^{μ} and X^{μ} via (4.38), we may show that the d'Alembertian operator may be rewritten as

$$\Box_x^2\;=\;\Box_R^2\,+\,\partial_{R,\mu}\partial_X^{\mu}\,+\,\frac{1}{4}\Box_X^2\,.\tag{E.2}$$

Performing a gradient expansion of the Wigner transform of (E.1), as described in Section 4.2, we obtain

$$\begin{aligned}&\Big(q^2\,-\,M^2\,+\,iq\cdot\partial_X\,-\,\frac{1}{4}\Box_X^2\Big)\Delta_{\gtrless}(q,X)\\&\quad+\int\frac{\mathrm{d}^4Q}{(2\pi)^4}\,(2\pi)^4\delta_t^{(4)}(Q)\,\exp\Big[-i\Big(Q\cdot X\,+\,\Diamond_{q,X}^{-}\,+\,2\Diamond_{Q,X}^{+}\Big)\Big]\\&\quad\Big(\{\Pi_{\gtrless}(q+Q/2,X)\}\{\Delta_{\mathcal{P}}(q-Q/2,X)\}\,+\,\{\Pi_{\mathcal{P}}(q+Q/2,X)\}\{\Delta_{\gtrless}(q-Q/2,X)\}\Big)\\&=\,\frac{1}{2}\int\frac{\mathrm{d}^4Q}{(2\pi)^4}\,(2\pi)^4\delta_t^{(4)}(Q)\,\exp\Big[-i\Big(Q\cdot X\,+\,\Diamond_{q,X}^{-}\,+\,2\Diamond_{Q,X}^{+}\Big)\Big]\\&\quad\Big(\{\Pi_{<}(q+Q/2,X)\}\{\Delta_{>}(q-Q/2,X)\}\,-\,\{\Pi_{>}(q+Q/2,X)\}\{\Delta_{<}(q-Q/2,X)\}\Big)\,.\end{aligned}\tag{E.3}$$

The diamond operators $\Diamond^{\pm}\{\bullet\}\{\bullet\}$ are defined in (4.46) and (4.47) of Section 4.2. Subsequently separating the Hermitian and anti-Hermitian parts of (E.3), we find the so-called constraint and kinetic equations

$$\begin{aligned}&\Big(q^2\,-\,M^2\,-\,\frac{1}{4}\Box_X^2\Big)\Delta_{\gtrless}(q,X)\,+\int\frac{\mathrm{d}^4Q}{(2\pi)^4}\,(2\pi)^4\delta_t^{(4)}(Q)\,\cos\Big(Q\cdot X\,+\,\Diamond_{q,X}^{-}\,+\,2\Diamond_{Q,X}^{+}\Big)\\&\quad\Big(\{\Pi_{\gtrless}(q+\tfrac{Q}{2},X)\}\{\Delta_{\mathcal{P}}(q-\tfrac{Q}{2},X)\}\,+\,\{\Pi_{\mathcal{P}}(q+\tfrac{Q}{2},X)\}\{\Delta_{\gtrless}(q-\tfrac{Q}{2},X)\}\Big)\\&=\,\frac{i}{2}\int\frac{\mathrm{d}^4Q}{(2\pi)^4}\,(2\pi)^4\delta_t^{(4)}(Q)\,\sin\Big(Q\cdot X\,+\,\Diamond_{q,X}^{-}\,+\,2\Diamond_{Q,X}^{+}\Big)\\&\quad\Big(\{\Pi_{>}(q+\tfrac{Q}{2},X)\}\{\Delta_{<}(q-\tfrac{Q}{2},X)\}\,+\,\{\Pi_{<}(q+\tfrac{Q}{2},X)\}\{\Delta_{>}(q-\tfrac{Q}{2},X)\}\Big)\,,\end{aligned}\tag{E.4a}$$

$$\begin{aligned} & q \cdot \partial_X \Delta_{\gtrless}(q, X) \;-\; \int \frac{\mathrm{d}^4 Q}{(2\pi)^4}\,(2\pi)^4 \delta_t^{(4)}(Q)\,\sin\Big(Q \cdot X \;+\; \diamondsuit_{q,X}^{-} \;+\; 2\diamondsuit_{Q,X}^{+}\Big) \\ & \qquad \Big(\{\Pi_{\gtrless}(q + \tfrac{Q}{2}, X)\}\{\Delta_{\mathcal{P}}(q - \tfrac{Q}{2}, X)\} \;+\; \{\Pi_{\mathcal{P}}(q + \tfrac{Q}{2}, X)\}\{\Delta_{\gtrless}(q - \tfrac{Q}{2}, X)\}\Big) \\ & \quad = \frac{i}{2} \int \frac{\mathrm{d}^4 Q}{(2\pi)^4}\,(2\pi)^4 \delta_t^{(4)}(Q)\,\cos\Big(Q \cdot X \;+\; \diamondsuit_{q,X}^{-} \;+\; 2\diamondsuit_{Q,X}^{+}\Big) \\ & \qquad \Big(\{\Pi_{>}(q + \tfrac{Q}{2}, X)\}\{\Delta_{<}(q - \tfrac{Q}{2}, X)\} \;-\; \{\Pi_{<}(q + \tfrac{Q}{2}, X)\}\{\Delta_{>}(q - \tfrac{Q}{2}, X)\}\Big)\,, \end{aligned} \tag{E.4b}$$

respectively.

In the late-time limit $t \to \infty$, the Q dependence is removed and the microscopic violation of energy conservation resulting from the uncertainty principle is neglected. As was established in Section 9, this microscopic violation of energy conservation is significant to the early-time dynamics of the system. Subsequently, keeping terms to zeroth order in the gradient expansion, the above expressions reduce to the more familiar form of the Kadanoff–Baym equations [1, 42, 43, 52, 53]:

$$\Big(q^2 \;-\; M^2\Big)\Delta_{\gtrless}(q, X) \;=\; -\Big(\Pi_{\gtrless}(q, X)\Delta_{\mathcal{P}}(q, X) \;+\; \Pi_{\mathcal{P}}(q, X)\Delta_{\gtrless}(q, X)\Big)\,, \tag{E.5a}$$

$$q \cdot \partial_X \Delta_{\gtrless}(q, X) \;=\; \frac{i}{2}\Big(\Pi_{>}(q, X)\Delta_{<}(q, X) \;-\; \Pi_{<}(q, X)\Delta_{>}(q, X)\Big)\,. \tag{E.5b}$$

The kinetic equation (E.5b) is to be compared with the semi-classical Boltzmann transport equation (6.18) and the energy-conserving limit of our time evolution equations in (8.17).

F Non-Homogeneous Loop Integrals

In this final appendix, we outline the techniques we have been using to perform the loop integrals that occur with the modified time-dependent Feynman rules of this new perturbative approach. In particular, we develop a method for dealing with the energy-non-conserving vertices that utilizes the Laplace transform. Lastly, we summarize the time-*independent* equilibrium and time-*dependent* spatially homogeneous limits of these integrals.

We begin by defining a generalization of the zero-temperature B_0 function [84]

$$B_0^{T\,=\,0}(q, m_1, m_2) \;=\; (2\pi\mu)^{4-d} \int \frac{\mathrm{d}^d k}{i\pi^2}\,\frac{1}{k^2 \;-\; m_1^2 \;+\; i\epsilon}\,\frac{1}{(k - q)^2 \;-\; m_1^2 \;+\; i\epsilon}\,. \tag{F.1}$$

To this end, we define the non-homogeneous CTP B_0 function, which may be written in the 2×2 matrix form as

$$B_0^{ab} \;=\; \begin{bmatrix} B_0 & B_0^{<} \\ B_0^{>} & -B_0^{*} \end{bmatrix}\,, \tag{F.2}$$



Figure 17: The non-homogeneous B_0^{ab} function.

consistent with the propagators and self-energies of the CTP formalism in Sections 3 and 4. The elements of B_0^{ab} are given by the following integral:

$$\begin{aligned} B_0^{ab}(q_1, q_2, m_1, m_2, \tilde{t}_f; \tilde{t}_i) \ \equiv \ & (2\pi\mu)^{4-d} e^{i(q_1^0 \, - \, q_2^0)\tilde{t}_f} \int \cdots \int \frac{\mathrm{d}^d k_1}{i\pi^2} \, \frac{\mathrm{d}^4 k_1'}{(2\pi)^4} \, \frac{\mathrm{d}^4 k_2}{(2\pi)^4} \, \frac{\mathrm{d}^4 k_2'}{(2\pi)^4} \\ & \times \ (2\pi)^4 \delta_t^{(4)}(q_1 \, - \, k_1 \, - \, k_2)(2\pi)^4 \delta_t^{(4)}(q_2 \, - \, k_1' \, - \, k_2') \\ & \times \ \eta^{acd} \Delta_{ce}^0(k_1, k_1', m_1, \tilde{t}_f; \tilde{t}_i) \Delta_{df}^{C,\,0}(k_2, k_2', m_2, \tilde{t}_f; \tilde{t}_i) \eta^{efb}, \end{aligned} \tag{F.3}$$

where $\delta_t^{(4)}$ is defined in (3.49) and (3.50), $\Delta^{0,\,ab}(p, p', t)$ is the free CTP propagator derived in Section 4.3 and the upper-case Roman C denotes charge-conjugate free CTP propagator. We reiterate that lower-case Roman CTP indices are raised and lowered by contraction with the $\mathbb{SO}\,(1,\ 1)$ 'metric' $\eta_{ab} \ = \ \mathrm{diag}\,(1,\ -1)$ and that $\eta_{abc} \ = \ +1$ is for $a \ = \ b \ = \ c \ = \ 1$, $\eta_{abc} \ = \ -1$ for $a \ = \ b \ = \ c \ = \ 2$ and $\eta_{abc} \ = \ 0$ otherwise.

In the definition (F.3), we have assumed that the statistical distribution functions are cut-off in the ultra-violet. Thus, any ultra-violet divergences anticipated from the superficial degree of divergence of the integral will be those that result from the homogeneous zero-temperature contribution. As a result, the dimensional regularization of the integral has been restricted to the k_1 dependence only.

In order to deal with the product of time-dependent energy-non-conserving vertices, we make the following replacement:

$$\delta_t(x)\delta_t(y) \ = \ \int_{\sigma - i\infty}^{\sigma + i\infty} \frac{\mathrm{d}s}{2\pi i} \, e^{st} \, \frac{2}{\pi^2} \, \frac{4s}{(x \, - \, y)^2 \, + \, 4s^2} \, \frac{1}{(x \, + \, y)^2 \, + \, 4s^2} \ , \tag{F.4}$$

where the RHS is the inverse Laplace transform and $s \ \in \ \mathbb{C}$ is a complex variable. The Bromwich contour is chosen so that $\sigma \ \in \ \mathbb{R}$ is larger than the real part of the right-most pole in the integrand, in order to ensure convergence. We then introduce the representation $B_0^{ab}(q_1, q_2, m_1, m_2, s)$ of the non-homogeneous CTP B_0 function through

$$B^{ab}(q_1, q_2, m_1, m_2, \tilde{t}_f; \tilde{t}_i) \ = \ \int_{\sigma - i\infty}^{\sigma + i\infty} \frac{\mathrm{d}s}{2\pi i} \, e^{st} \, B_0^{ab}(q_1, q_2, m_1, m_2, s) \ . \tag{F.5}$$

Note that $B_0^{ab}(q_1, q_2, m_1, m_2, s)$ is *not* the exact Laplace transform of $B_0^{ab}(q_1, q_2, m_1, m_2, t)$, since we have not transformed the t-dependence of the distribution functions. We suppress the dependence of $B_0^{ab}(q_1, q_2, m_1, m_2, s)$ on $\tilde{t}_f$ and $\tilde{t}_i$ for notational convenience.

After making the replacement (F.4) in (F.3), we obtain the integral

$$\begin{aligned} B_0^{ab}(q_1, q_2, m_1, m_2, s) \ &= \ 8(2\pi\mu)^{4-d} e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f} \int \cdots \int \frac{\mathrm{d}^d k_1}{i\pi^2}\,\frac{\mathrm{d}^4 k_1'}{(2\pi)^4}\,\frac{\mathrm{d}^4 k_2}{(2\pi)^4}\,\frac{\mathrm{d}^4 k_2'}{(2\pi)^4} \\ &\times\ (2\pi)^3\delta^{(3)}(\mathbf{q}_1 - \mathbf{k}_1 - \mathbf{k}_2)(2\pi)^3\delta^{(3)}(\mathbf{q}_2 - \mathbf{k}_1' - \mathbf{k}_2') \\ &\times\ 4s\Big[\Big(q_1^0 - q_2^0 - k_1^0 + {k_1'}^0 - k_2^0 + {k_2'}^0\Big)^2 + 4s^2\Big]^{-1} \\ &\times\ \Big[\Big(q_1^0 + q_2^0 - k_1^0 - {k_1'}^0 - k_2^0 - {k_2'}^0\Big)^2 + 4s^2\Big]^{-1} \\ &\times\ \eta^{acd}\Delta^0_{ce}(k_1, k_1', m_1, \tilde{t}_f; \tilde{t}_i)\Delta^{0,C}_{df}(k_2, k_2', m_2, \tilde{t}_f; \tilde{t}_i)\eta^{efb}\ , \end{aligned} \tag{F.6}$$

in which the analytic structure of the product of formerly t-dependent vertex functions is now manifest.

For conciseness, throughout this appendix we use the short-hand notation:

$$\sum_{\{\alpha\}} \tag{F.7}$$

for summation over only the $\alpha_i = \pm 1$ and $\alpha_i' = \pm 1$ that appear explicitly in an expression. On-shell energies are denoted by

$$E_i \ \equiv\ E_i(\mathbf{k}_i) \ =\ \sqrt{\mathbf{k}_i^2 + m_i^2}\ , \qquad E_i' \ \equiv\ E_i'(\mathbf{k}_i') \ =\ \sqrt{{\mathbf{k}_i'}^2 + m_i^2}\ , \tag{F.8}$$

and on-shell four-momenta by

$$\hat{k}_i \ \equiv\ \hat{k}_i^{\mu} \ =\ (\alpha_i E_i(\mathbf{k}_i), \mathbf{k}_i)\ , \qquad \hat{k}_i' \ \equiv\ \hat{k}_i'^{\mu} \ =\ (\alpha_i' E_i'(\mathbf{k}_i'), \mathbf{k}_i')\ . \tag{F.9}$$

For the energy factor in (4.54), we use the notation

$$\mathscr{E}_i \ \equiv\ \mathscr{E}_i(\mathbf{k}_i, \mathbf{k}_i') \ =\ \sqrt{E_i(\mathbf{k}_i)E_i(\mathbf{k}_i')}\ . \tag{F.10}$$

Products of unit-step functions are denoted by

$$\theta(x, y, \cdots, z) \ \equiv\ \theta(x)\theta(y)\cdots\theta(z)\ . \tag{F.11}$$

Time-Ordered Functions

The $(a, b) = (1, 1)$ element of (F.3) coincides with the time-ordered function B_0, which we may separate into four contributions (suppressing all arguments):

$$B_0 \ =\ I^{(\mathrm{i})} + I^{(\mathrm{ii})} + I^{(\mathrm{iiia})} + I^{(\mathrm{iiib})}\ . \tag{F.12}$$

These four contributions are the zero temperature limit ($I^{(i)}$), the purely thermal term ($I^{(ii)}$) and the cross terms ($I^{(iiia)}$ and $I^{(iiib)}$).

(i) **The zero-temperature part** may be extracted from the product of terms that remain in the limit of vanishing statistical distribution functions. This part is

$$\begin{aligned}
I^{(\mathrm{i})}(q_1, q_2, m_1, m_2, s) \;=\; & 8(2\pi\mu)^{4-d} e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f} \int \cdots \int \frac{\mathrm{d}^d k_1}{i\pi^2} \frac{\mathrm{d}^4 k_1'}{(2\pi)^4} \frac{\mathrm{d}^4 k_2}{(2\pi)^4} \frac{\mathrm{d}^4 k_2'}{(2\pi)^4} \\
\times\; & (2\pi)^3 \delta^{(3)}(\mathbf{q}_1 - \mathbf{k}_1 - \mathbf{k}_2)(2\pi)^3 \delta^{(3)}(\mathbf{q}_2 - \mathbf{k}_1' - \mathbf{k}_2') \\
\times\; & 4s\Big[\Big(q_1^0 - q_2^0 - k_1^0 + k_1'^{\,0} - k_2^0 + k_2'^{\,0}\Big)^2 + 4s^2\Big]^{-1} \\
\times\; & \Big[\Big(q_1^0 + q_2^0 - k_1^0 - k_1'^{\,0} - k_2^0 - k_2'^{\,0}\Big)^2 + 4s^2\Big]^{-1} \\
\times\; & \frac{1}{k_1^2 - m_1^2 + i\epsilon}(2\pi)^4 \delta^{(4)}(k_1 - k_1') \frac{1}{k_2^2 - m_2^2 + i\epsilon}(2\pi)^4 \delta^{(4)}(k_2 - k_2')\,. \qquad \text{(F.13)}
\end{aligned}$$

The $k_1'^{\,0}$ and $k_2'^{\,0}$ integrations are performed by means of the delta functions, giving

$$\begin{aligned}
I^{(\mathrm{i})}(q_1, q_2, m_1, m_2, s) \;=\; & (2\pi\mu)^{4-d} e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f} \int \cdots \int \frac{\mathrm{d}^{d-1} k_1}{i\pi^2} \frac{\mathrm{d}^3 \mathbf{k}_1'}{(2\pi)^3} \frac{\mathrm{d}^3 \mathbf{k}_2}{(2\pi)^3} \frac{\mathrm{d}^3 \mathbf{k}_2'}{(2\pi)^3}\, \mathrm{d}k_1^0\, \mathrm{d}k_2^0 \\
\times\; & (2\pi)^3 \delta^{(3)}(\mathbf{q}_1 - \mathbf{k}_1 - \mathbf{k}_2)(2\pi)^3 \delta^{(3)}(\mathbf{q}_2 - \mathbf{k}_1' - \mathbf{k}_2') \\
\times\; & \frac{4s}{\pi}\Big[\Big(q_1^0 - q_2^0\Big)^2 + 4s^2\Big]^{-1}\Big[\Big(\tfrac{q_1^0\,+\,q_2^0}{2} - k_1^0 - k_2^0\Big)^2 + s^2\Big]^{-1} \\
\times\; & \frac{1}{(k_1^0)^2 - E_1^2 + i\epsilon}\, \frac{1}{(k_2^0)^2 - E_2^2 + i\epsilon}(2\pi)^3 \delta^{(3)}(\mathbf{k}_1 - \mathbf{k}_1')(2\pi)^3 \delta^{(3)}(\mathbf{k}_2 - \mathbf{k}_2')\,.
\end{aligned}$$

(F.14)

By virtue of the residue theorem, we may perform the k_1^0 and k_2^0 integrations by closing contours in the lower-halves of the k_1^0 and k_2^0 complex planes. After collecting together the resulting terms, we find

$$\begin{aligned}
I^{(\mathrm{i})}(q_1, q_2, m_1, m_2, s) \;=\; & -2(2\pi)^3 \mu^{4-d} e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f} \sum_{\{\alpha\}} \int \cdots \int \frac{\mathrm{d}^{d-1} \mathbf{k}_1}{(2\pi)^{d-1}} \frac{\mathrm{d}^3 \mathbf{k}_1'}{(2\pi)^3} \frac{\mathrm{d}^3 \mathbf{k}_2}{(2\pi)^3} \frac{\mathrm{d}^3 \mathbf{k}_2'}{(2\pi)^3} \\
\times\; & \frac{1}{E_1 E_2}\,(2\pi)^3 \delta^{(3)}(\mathbf{q}_1 - \mathbf{k}_1 - \mathbf{k}_2)(2\pi)^3 \delta^{(3)}(\mathbf{q}_2 - \mathbf{k}_1' - \mathbf{k}_2') \\
\times\; & \frac{1}{\pi}\Big[\Big(q_1^0 - q_2^0\Big)^2 + 4s^2\Big]^{-1} \alpha_1 \Big[\tfrac{q_1^0\,+\,q_2^0}{2} - \alpha_1\Big(E_1 + E_2 - is\Big)\Big]^{-1} \\
\times\; & (2\pi)^3 \delta^{(3)}(\mathbf{k}_1 - \mathbf{k}_1')(2\pi)^3 \delta^{(3)}(\mathbf{k}_2 - \mathbf{k}_2')\,, \qquad \text{(F.15)}
\end{aligned}$$

or, in terms of the Lorentz-invariant phase-space measures in (2.3),

$$\begin{aligned}
I^{(\mathrm{i})}(q_1, q_2, m_1, m_2, s) \;=\; & -8(2\pi)^3 \mu^{4-d} e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f} \sum_{\{\alpha\}} \int \cdots \int \mathrm{d}\Pi_{\mathbf{k}_1}^{d-1}\, \mathrm{d}\Pi_{\mathbf{k}_1'}\, \mathrm{d}\Pi_{\mathbf{k}_2}\, \mathrm{d}\Pi_{\mathbf{k}_2'} \\
\times\; & (2\pi)^3 \delta^{(3)}(\mathbf{q}_1 - \mathbf{k}_1 - \mathbf{k}_2)(2\pi)^3 \delta^{(3)}(\mathbf{q}_2 - \mathbf{k}_1' - \mathbf{k}_2') \\
\times\; & \frac{1}{\pi}\Big[\Big(q_1^0 - q_2^0\Big)^2 + 4s^2\Big]^{-1} \alpha_1 \Big[\tfrac{q_1^0\,+\,q_2^0}{2} - \alpha_1\Big(E_1 + E_2 - is\Big)\Big]^{-1} \\
\times\; & (2\pi)^3 2E_1' \delta^{(3)}(\mathbf{k}_1 - \mathbf{k}_1')(2\pi)^3 2E_2' \delta^{(3)}(\mathbf{k}_2 - \mathbf{k}_2')\,, \qquad \text{(F.16)}
\end{aligned}$$

where the superscript $d-1$ indicates that the $\mathbf{k}_1$ integration is to be taken over a $(d-1)$-dimensional phase-space.

The Laplace transform $\mathcal{F}(s)$ of a complex-valued function $F(t)$ has the following properties:

$$\mathcal{L}_t[\mathrm{Re}\,F](s) \;=\; \frac{1}{2}\Big[\,\mathcal{F}(s)\,+\,\Big(\mathcal{F}(s^*)\Big)^*\Big]\,, \tag{F.17a}$$

$$\mathcal{L}_t[\mathrm{Im}\,F](s) \;=\; \frac{1}{2i}\Big[\,\mathcal{F}(s)\,-\,\Big(\mathcal{F}(s^*)\Big)^*\Big]\,. \tag{F.17b}$$

We may then identify the dispersive and absorptive parts of the t-dependent B_0 function with the parts symmetric and anti-symmetric in s, respectively. With this separation, (F.16) becomes

$$\begin{aligned} I^{(\mathrm{i})}(q_1,q_2,m_1,m_2,s) \;&=\; 8(2\pi)^3\mu^{4-d}e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f}\sum_{\{\alpha\}}\int\cdots\int \mathrm{d}\Pi_{k_1}^{d-1}\,\mathrm{d}\Pi_{\mathbf{k}_1'}\,\mathrm{d}\Pi_{\mathbf{k}_2}\,\mathrm{d}\Pi_{\mathbf{k}_2'} \\ &\times\; (2\pi)^3\delta^{(3)}(\mathbf{q}_1\,-\,\mathbf{k}_1\,-\,\mathbf{k}_2)(2\pi)^3\delta^{(3)}(\mathbf{q}_2\,-\,\mathbf{k}_1'\,-\,\mathbf{k}_2') \\ &\times\; \frac{1}{\pi}\Big[\Big(\,q_1^0\,-\,q_2^0\Big)^2\,+\,4s^2\Big]^{-1}\Big\{\Big[\,\tfrac{q_1^0\,+\,q_2^0}{2}\,-\,\alpha_1\Big(\,E_1\,+\,E_2\Big)\Big]^2\,+\,s^2\Big\}^{-1} \\ &\times\; \Big\{\,-\,\alpha_1\Big[\,\tfrac{q_1^0\,+\,q_2^0}{2}\,-\,\alpha_1\Big(\,E_1\,+\,E_2\Big)\Big]\,+\,is\Big\} \\ &\times\; (2\pi)^3 2E_1'\delta^{(3)}(\mathbf{k}_1\,-\,\mathbf{k}_1')(2\pi)^3 2E_2'\delta^{(3)}(\mathbf{k}_2\,-\,\mathbf{k}_2')\,, \end{aligned} \tag{F.18}$$

where the delineation between dispersive and absorptive parts is identified within the curly brackets of the fourth line.

Isolating the dispersive part of this result and performing all but one of the remaining phase-space integrals, we find

$$\begin{aligned} \mathrm{Disp}\,B_0(q_1,q_2,m_1,m_2,s) \;&=\; (2\pi)^4\frac{1}{\pi}\,\frac{1}{\Big(\,q_1^0\,-\,q_2^0\,\Big)^2\,+\,4s^2}\,\delta^{(3)}(\mathbf{q}_1\,-\,\mathbf{q}_2)e^{i(q_1^0\,-\,q_2^0)t/2} \\ \times\sum_{\{\alpha\}}-(2\pi\mu)^{4-d}&\int \mathrm{d}^{d-1}\mathbf{k}_1\,\frac{E_1\,+\,E_2}{E_1E_2}\,\frac{1}{\pi}\Big[\Big(\,\tfrac{q_1^0+q_2^0}{2}\,-\,i\alpha s\Big)^2\,-\,\Big(\,E_1\,+\,E_2\Big)^2\Big]^{-1}\,, \end{aligned} \tag{F.19}$$

where Disp stands for the dispersive part. We compare this result to the form of the zero-temperature B_0 function (F.1) after the k_0 integration has been performed:

$$B_0^{T\,=\,0}(q,m_1,m_2) \;=\; -(2\pi\mu)^{4-d}\int \mathrm{d}^{d-1}\mathbf{k}_1\;\frac{E_1\,+\,E_2}{E_1E_2}\;\frac{1}{\pi}\;\Big[\;q_0^2\;-\;\Big(\;E_1\;+\;E_2\Big)^2\Big]^{-1}\,. \tag{F.20}$$

Hence, we may write

$$\begin{aligned} \mathrm{Disp}\,I^{(\mathrm{i})}(q_1,q_2,m_1,m_2,s) \;&=\; (2\pi)^4\frac{1}{\pi}\,\frac{1}{\Big(\,q_1^0\,-\,q_2^0\,\Big)^2\,+\,4s^2}\,\delta^{(3)}(\mathbf{q}_1\,-\,\mathbf{q}_2)e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f} \\ &\times\;\sum_{\{\alpha\}}B_0^{T\,=\,0}\Big(\tfrac{q_1^0+q_2^0}{2}\,-\,i\alpha s,\,\tfrac{\mathbf{q}_1+\mathbf{q}_2}{2},m_1,m_2\Big)\,, \end{aligned} \tag{F.21}$$

where, for $d\,=\,4\,-\,2\epsilon$ dimensions,

$$\begin{aligned} B_0^{T=0}(q^0-i\alpha s,\mathbf{q},m_1,m_2) \;&=\; \frac{1}{\epsilon}-\gamma_{\mathrm{E}}+\ln\frac{4\pi\mu^2}{m_1m_2}\,+\,\frac{1}{\Big(q^0\,-\,i\alpha s\Big)^2\,-\,\mathbf{q}^2}\Big[\,\Big(m_2^2-m_1^2\Big)\ln\frac{m_1^2}{m_2^2} \\ &+\;\lambda^{1/2}\Big((q^0-i\alpha s)^2-\mathbf{q}^2,m_1^2,m_2^2\Big)\cosh^{-1}\Big(\frac{m_1^2\,+\,m_2^2\,-\,(q^0-i\alpha s)^2\,+\,\mathbf{q}^2}{2m_1m_2}\Big)\Big]\,, \end{aligned} \tag{F.22}$$

containing the zero-temperature UV divergence [84]. In (F.22), γ_{E} is the Euler constant and μ is the 't Hooft mass scale.
(ii) **The purely thermal part** is more straightforward to analyze:

$$\begin{aligned}
I^{(\mathrm{ii})}(q_1, q_2, m_1, m_2, s) \ &= \ -8(2\pi\mu)^{4-d} e^{i(q_1^0 \,-\, q_2^0)\tilde{t}_f} \int \cdots \int \frac{\mathrm{d}^d k_1}{i\pi^2} \, \frac{\mathrm{d}^4 k_1'}{(2\pi)^4} \, \frac{\mathrm{d}^4 k_2}{(2\pi)^4} \, \frac{\mathrm{d}^4 k_2'}{(2\pi)^4} \\
&\times \ (2\pi)^3 \delta^{(3)}(\mathbf{q}_1 - \mathbf{k}_1 - \mathbf{k}_2)(2\pi)^3 \delta^{(3)}(\mathbf{q}_2 - \mathbf{k}_1' - \mathbf{k}_2') \\
&\times \ 4s\Big[\Big(q_1^0 - q_2^0 - k_1^0 + k_1'^{\,0} - k_2^0 + k_2'^{\,0}\Big)^2 + 4s^2\Big]^{-1} \\
&\times \ \Big[\Big(q_1^0 + q_2^0 - k_1^0 - k_1'^{\,0} - k_2^0 - k_2'^{\,0}\Big)^2 + 4s^2\Big]^{-1} \\
&\times \ 2\pi\delta(k_1^2 - m_1^2)|2k_1^0|^{1/2} \tilde{f}_1(k_1, k_1', t) e^{i(k_1^0 \,-\, k_1'^{\,0})\tilde{t}_f} |2k_1'^{\,0}|^{1/2} 2\pi\delta(k_1'^{\,2} - m_1^2) \\
&\times \ 2\pi\delta(k_2^2 - m_2^2)|2k_2^0|^{1/2} \tilde{f}_2^C(k_2, k_2', t) e^{i(k_2^0 \,-\, k_2'^{\,0})\tilde{t}_f} |2k_2'^{\,0}|^{1/2} 2\pi\delta(k_2'^{\,2} - m_2^2) \, .
\end{aligned} \tag{F.23}$$

Performing the four zeroth-component momentum integrations by virtue of the on-shell delta functions $\delta(k_i^2 - m_i^2)$ and $\delta(k_i'^2 - m_i^2)$ in (F.23), we find

$$\begin{aligned}
I^{(\mathrm{ii})}(q_1, q_2, m_1, m_2, s) \ &= \ 8(2\pi)^3 \mu^{4-d} e^{i(q_1^0 \,-\, q_2^0)\tilde{t}_f} \sum_{\{\alpha\}} \int \cdots \int \mathrm{d}\Pi_{k_1}^{d-1} \, \mathrm{d}\Pi_{\mathbf{k}_1'} \, \mathrm{d}\Pi_{\mathbf{k}_2} \, \mathrm{d}\Pi_{\mathbf{k}_2'} \\
&\times \ (2\pi)^3 \delta^{(3)}(\mathbf{q}_1 - \mathbf{k}_1 - \mathbf{k}_2)(2\pi)^3 \delta^{(3)}(\mathbf{q}_2 - \mathbf{k}_1' - \mathbf{k}_2') \\
&\times \ \Big[\Big(q_1^0 - q_2^0 - \alpha_1 E_1 + \alpha_1' E_1' - \alpha_2 E_2 + \alpha_2' E_2'\Big)^2 + 4s^2\Big]^{-1} \\
&\times \ \Big[\Big(\tfrac{q_1^0 + q_2^0}{2} - \tfrac{\alpha_1 E_1 + \alpha_1' E_1'}{2} - \tfrac{\alpha_2 E_2 + \alpha_2' E_2'}{2}\Big)^2 + s^2\Big]^{-1} 2is \\
&\times \ 2\mathscr{E}_1 \tilde{f}_1(\hat{k}_1, \hat{k}_1', t) e^{i(\alpha_1 E_1 \,-\, \alpha_1' E_1')\tilde{t}_f} 2\mathscr{E}_2 \tilde{f}_2^C(\hat{k}_2, \hat{k}_2', t) e^{i(\alpha_2 E_2 \,-\, \alpha_2' E_2')\tilde{t}_f} \, .
\end{aligned} \tag{F.24}$$

Note that the purely thermal part in (F.24) contains only absorptive contributions.
(iii) **The cross-terms** yield both dispersive and absorptive contributions. The first of the cross terms yields

$$\begin{aligned}
I^{(\mathrm{iiia})}(q_1, q_2, m_1, m_2, s) \ &= \ -8i(2\pi\mu)^{4-d} e^{i(q_1^0 \,-\, q_2^0)\tilde{t}_f} \int \cdots \int \frac{\mathrm{d}^d k_1}{i\pi^2} \, \frac{\mathrm{d}^4 k_1'}{(2\pi)^4} \, \frac{\mathrm{d}^4 k_2}{(2\pi)^4} \, \frac{\mathrm{d}^4 k_2'}{(2\pi)^4} \\
&\times \ (2\pi)^3 \delta^{(3)}(\mathbf{q}_1 - \mathbf{k}_1 - \mathbf{k}_2)(2\pi)^3 \delta^{(3)}(\mathbf{q}_2 - \mathbf{k}_1' - \mathbf{k}_2') \\
&\times \ 4s\Big[\Big(q_1^0 - q_2^0 - k_1^0 + k_1'^{\,0} - k_2^0 + k_2'^{\,0}\Big)^2 + 4s^2\Big]^{-1} \\
&\times \ \Big[\Big(q_1^0 + q_2^0 - k_1^0 - k_1'^{\,0} - k_2^0 - k_2'^{\,0}\Big)^2 + 4s^2\Big]^{-1} \\
&\times \ \frac{1}{k_1^2 - m_1^2 + i\epsilon} (2\pi)^4 \delta^{(4)}(k_1 - k_1') \\
&\times \ 2\pi\delta(k_2^2 - m_2^2)|2k_2^0|^{1/2} \tilde{f}_2^C(k_2, k_2', t) e^{i(k_2^0 \,-\, k_2'^{\,0})\tilde{t}_f} |2k_2'^{\,0}|^{1/2} 2\pi\delta(k_2'^{\,2} - m_2^2) \, .
\end{aligned} \tag{F.25}$$

After evaluating the $k_1'^{\,0}$, k_2^0 and $k_2'^{\,0}$ integrals by virtue of the delta functions, we perform the k_1^0 integral by closing a contour in the lower half of the k_1^0 complex plane. Equation (F.25) may then be written in the more compact form

$$\begin{aligned} I^{(\mathrm{iiia})}(q_1, q_2, m_1, m_2, s) \;=\; & -8(2\pi)^3 \mu^{4-d} e^{i(q_1^0 \,-\, q_2^0)\tilde{t}_f} \sum_{\{\alpha\}} \int \cdots \int \mathrm{d}\Pi_{\mathbf{k}_1}^{d-1}\, \mathrm{d}\Pi_{\mathbf{k}_1'}\, \mathrm{d}\Pi_{\mathbf{k}_2}\, \mathrm{d}\Pi_{\mathbf{k}_2'} \\ & \times\; (2\pi)^3 \delta^{(3)}(\mathbf{q}_1 \,-\, \mathbf{k}_1 \,-\, \mathbf{k}_2)(2\pi)^3 \delta^{(3)}(\mathbf{q}_2 \,-\, \mathbf{k}_1' \,-\, \mathbf{k}_2') \\ & \times\; \frac{1}{\pi}\Big[\Big(q_1^0 \,-\, q_2^0 \,-\, \alpha_2 E_2 \,+\, \alpha_2' E_2'\Big)^2 \,+\, 4s^2\Big]^{-1} \\ & \times\; \alpha_1 \Big[\tfrac{q_1^0 + q_2^0}{2} \,-\, \tfrac{\alpha_2 E_2 + \alpha_2' E_2'}{2} \,-\, \alpha_1\Big(E_1 \,-\, is\Big)\Big]^{-1} \\ & \times (2\pi)^3 2E_1' \delta^{(3)}(\mathbf{k}_1 \,-\, \mathbf{k}_1') 2\mathscr{E}_2 \tilde{f}_2^C(\hat{k}_2, \hat{k}_2', t) e^{i(\alpha_2 E_2 \,-\, \alpha_2' E_2')\tilde{t}_f}\ . \end{aligned} \tag{F.26}$$

Separating again into dispersive and absorptive parts as in (F.18), $I^{(\mathrm{iiia})}$ may be written as

$$\begin{aligned} I^{(\mathrm{iiia})}(q_1, q_2, m_1, m_2, s) \;=\; & 8(2\pi)^3 \mu^{4-d} e^{i(q_1^0 \,-\, q_2^0)\tilde{t}_f} \sum_{\{\alpha\}} \int \cdots \int \mathrm{d}\Pi_{\mathbf{k}_1}^{d-1}\, \mathrm{d}\Pi_{\mathbf{k}_1'}\, \mathrm{d}\Pi_{\mathbf{k}_2}\, \mathrm{d}\Pi_{\mathbf{k}_2'} \\ & \times\; (2\pi)^3 \delta^{(3)}(\mathbf{q}_1 \,-\, \mathbf{k}_1 \,-\, \mathbf{k}_2)(2\pi)^3 \delta^{(3)}(\mathbf{q}_2 \,-\, \mathbf{k}_1' \,-\, \mathbf{k}_2') \\ & \times\; \frac{1}{\pi}\Big[\Big(q_1^0 \,-\, q_2^0 \,-\, \alpha_2 E_2 \,+\, \alpha_2' E_2'\Big)^2 \,+\, 4s^2\Big]^{-1} \\ & \times\; \Big[\Big(\tfrac{q_1^0 + q_2^0}{2} \,-\, \alpha_1 E_1 \,-\, \tfrac{\alpha_2 E_2 + \alpha_2' E_2'}{2}\Big)^2 \,+\, s^2\Big]^{-1} \\ & \times\; \Big\{ -\alpha_1\Big(\tfrac{q_1^0 + q_2^0}{2} \,-\, \alpha_1 E_1 \,-\, \tfrac{\alpha_2 E_2 + \alpha_2' E_2'}{2}\Big) \,+\, is\Big\} \\ & \times\; (2\pi)^3 2E_1' \delta^{(3)}(\mathbf{k}_1 \,-\, \mathbf{k}_1') 2\mathscr{E}_1 \tilde{f}_2^C(\hat{k}_2, \hat{k}_2', t) e^{i(\alpha_2 E_2 \,-\, \alpha_2' E_2')\tilde{t}_f}\ . \end{aligned} \tag{F.27}$$

Similarly, for the second cross term $I^{(\mathrm{iiib})}$ in (F.12) , we obtain

$$\begin{aligned} I^{(\mathrm{iiib})}(q_1, q_2, m_1, m_2, s) \;=\; & 8(2\pi)^3 \mu^{4-d} e^{i(q_1^0 \,-\, q_2^0)\tilde{t}_f} \sum_{\{\alpha\}} \int \cdots \int \mathrm{d}\Pi_{\mathbf{k}_1}^{d-1}\, \mathrm{d}\Pi_{\mathbf{k}_1'}\, \mathrm{d}\Pi_{\mathbf{k}_2}\, \mathrm{d}\Pi_{\mathbf{k}_2'} \\ & \times\; (2\pi)^3 \delta^{(3)}(\mathbf{q}_1 \,-\, \mathbf{k}_1 \,-\, \mathbf{k}_2)(2\pi)^3 \delta^{(3)}(\mathbf{q}_2 \,-\, \mathbf{k}_1' \,-\, \mathbf{k}_2') \\ & \times\; \frac{1}{\pi}\Big[\Big(q_1^0 \,-\, q_2^0 \,-\, \alpha_1 E_1 \,+\, \alpha_1' E_1'\Big)^2 \,+\, 4s^2\Big]^{-1} \\ & \times\; \Big[\Big(\tfrac{q_1^0 + q_2^0}{2} \,-\, \tfrac{\alpha_1 E_1 + \alpha_1' E_1'}{2} \,-\, \alpha_2 E_2\Big)^2 \,+\, s^2\Big]^{-1} \\ & \times\; \Big\{ -\alpha_2\Big(\tfrac{q_1^0 + q_2^0}{2} \,-\, \tfrac{\alpha_1 E_1 + \alpha_1' E_1'}{2} \,-\, \alpha_2 E_2\Big) \,+\, is\Big\} \\ & \times\; (2\pi)^3 2E_2' \delta^{(3)}(\mathbf{k}_2 \,-\, \mathbf{k}_2') 2\mathscr{E}_1 \tilde{f}_1(\hat{k}_1, \hat{k}_1', t) e^{i(\alpha_1 E_1 \,-\, \alpha_1' E_1')\tilde{t}_f}\ . \end{aligned} \tag{F.28}$$

Collecting the four contributions from (F.18), (F.24), (F.27) and (F.28) together, we find the full time-ordered function

$$\begin{aligned} B_0(q_1, q_2, m_1, m_2, s) \;&=\; 8(2\pi)^3\mu^{4-d}e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f}\sum_{\{\alpha\}}\int\cdots\int \mathrm{d}\Pi^{d-1}_{\mathbf{k}_1}\,\mathrm{d}\Pi_{\mathbf{k}'_1}\,\mathrm{d}\Pi_{\mathbf{k}_2}\,\mathrm{d}\Pi_{\mathbf{k}'_2} \\ &\times\; (2\pi)^3\delta^{(3)}(\mathbf{q}_1 - \mathbf{k}_1 - \mathbf{k}_2)(2\pi)^3\delta^{(3)}(\mathbf{q}_2 - \mathbf{k}'_1 - \mathbf{k}'_2) \\ &\times\; 2\mathscr{E}_1 e^{i(\alpha_1E_1\,-\,\alpha'_1E'_1)\tilde{t}_f}\,2\mathscr{E}_2 e^{i(\alpha_2E_2\,-\,\alpha'_2E'_2)\tilde{t}_f} \\ &\times\; \frac{1}{\pi}\bigg[\Big(q_1^0 - q_2^0 - \alpha_1E_1 + \alpha'_1E'_1 - \alpha_2E_2 + \alpha'_2E'_2\Big)^2 + 4s^2\bigg]^{-1} \\ &\times\; \bigg[\Big(\frac{q_1^0+q_2^0}{2} - \frac{\alpha_1E_1+\alpha'_1E'_1}{2} - \frac{\alpha_2E_2+\alpha'_2E'_2}{2}\Big)^2 + s^2\bigg]^{-1} \\ &\times\; \bigg\{-\Big(\frac{q_1^0+q_2^0}{2} - \frac{\alpha_1E_1+\alpha'_1E'_1}{2} - \frac{\alpha_2E_2+\alpha'_2E'_2}{2}\Big)F^{\mathrm{R}}(\{\hat{k}\},t) + isF^{1}(\{\hat{k}\},t)\bigg\}\,, \end{aligned} \tag{F.29}$$

where $\{\hat{k}\} \equiv \{\hat{k}_1, \hat{k}'_1, \hat{k}_2, \hat{k}'_2\}$ is the set of on-shell four momenta. The statistical structure is contained within the distributions $F^{\mathrm{R}}(\{\hat{k}\},t)$ and $F^{1}(\{\hat{k}\},t)$ defined as follows:

$$\begin{aligned} F^{\mathrm{R}}(\{\hat{k}\},t) \;&=\; (2\pi)^3\delta^{(3)}(\mathbf{k}_1 - \mathbf{k}'_1)(2\pi)^3\delta^{(3)}(\mathbf{k}_2 - \mathbf{k}'_2) \\ &\qquad\times\; \Big(\theta(\alpha_1,\alpha'_1,\alpha_2,\alpha'_2) - \theta(-\alpha_1,-\alpha'_1,-\alpha_2,-\alpha'_2)\Big) \\ &\quad+\; (2\pi)^3\delta^{(3)}(\mathbf{k}_1 - \mathbf{k}'_1)\Big(\theta(\alpha_1,\alpha'_1) - \theta(-\alpha_1,-\alpha'_1)\Big)\tilde{f}_2^{C}(\hat{k}_2,\hat{k}'_2,t) \\ &\quad+\; \tilde{f}_1(\hat{k}_1,\hat{k}'_1,t)(2\pi)^3\delta^{(3)}(\mathbf{k}_2 - \mathbf{k}'_2)\Big(\theta(\alpha_2,\alpha'_2) - \theta(-\alpha_2,-\alpha'_2)\Big)\,, \end{aligned} \tag{F.30a}$$

$$\begin{aligned} F^{1}(\{\hat{k}\},t) \;&=\; (2\pi)^3\delta^{(3)}(\mathbf{k}_1 - \mathbf{k}'_1)(2\pi)^3\delta^{(3)}(\mathbf{k}_2 - \mathbf{k}'_2) \\ &\qquad\times\; \Big(\theta(\alpha_1,\alpha'_1,\alpha_2,\alpha'_2) + \theta(-\alpha_1,-\alpha'_1,-\alpha_2,-\alpha'_2)\Big) \\ &\quad+\; (2\pi)^3\delta^{(3)}(\mathbf{k}_1 - \mathbf{k}'_1)\Big(\theta(\alpha_1,\alpha'_1) + \theta(-\alpha_1,-\alpha'_1)\Big)\tilde{f}_2^{C}(\hat{k}_2,\hat{k}'_2,t) \\ &\quad+\; \tilde{f}_1(\hat{k}_1,\hat{k}'_1,t)(2\pi)^3\delta^{(3)}(\mathbf{k}_2 - \mathbf{k}'_2)\Big(\theta(\alpha_2,\alpha'_2) + \theta(-\alpha_2,-\alpha'_2)\Big) \\ &\quad+\; 2\tilde{f}_1(\hat{k}_1,\hat{k}'_1,t)\tilde{f}_2^{C}(\hat{k}_2,\hat{k}'_2,t)\,. \end{aligned} \tag{F.30b}$$

For completeness, the anti-time-ordered function, obtained from the RHS (F.29) by taking $s \to -s$ and multiplying by an overall minus sign, is given by

$$\begin{aligned} -\,B_0^*(q_1, q_2, m_1, m_2, s) \;&=\; 8(2\pi)^3\mu^{4-d}e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f}\sum_{\{\alpha\}}\int\cdots\int \mathrm{d}\Pi^{d-1}_{\mathbf{k}_1}\,\mathrm{d}\Pi_{\mathbf{k}'_1}\,\mathrm{d}\Pi_{\mathbf{k}_2}\,\mathrm{d}\Pi_{\mathbf{k}'_2} \\ &\times\; (2\pi)^3\delta^{(3)}(\mathbf{q}_1 - \mathbf{k}_1 - \mathbf{k}_2)(2\pi)^3\delta^{(3)}(\mathbf{q}_2 - \mathbf{k}'_1 - \mathbf{k}'_2) \\ &\times\; 2\mathscr{E}_1 e^{i(\alpha_1E_1\,-\,\alpha'_1E'_1)\tilde{t}_f}\,2\mathscr{E}_2 e^{i(\alpha_2E_2\,-\,\alpha'_2E'_2)\tilde{t}_f} \\ &\times\; \frac{1}{\pi}\bigg[\Big(q_1^0 - q_2^0 - \alpha_1E_1 + \alpha'_1E'_1 - \alpha_2E_2 + \alpha'_2E'_2\Big)^2 + 4s^2\bigg]^{-1} \\ &\times\; \bigg[\Big(\frac{q_1^0+q_2^0}{2} - \frac{\alpha_1E_1+\alpha'_1E'_1}{2} - \frac{\alpha_2E_2+\alpha'_2E'_2}{2}\Big)^2 + s^2\bigg]^{-1} \\ &\times\; \bigg\{\Big(\frac{q_1^0+q_2^0}{2} - \frac{\alpha_1E_1+\alpha'_1E'_1}{2} - \frac{\alpha_2E_2+\alpha'_2E'_2}{2}\Big)F^{\mathrm{R}}(\{\hat{k}\},t) + isF^{1}(\{\hat{k}\},t)\bigg\}\,, \end{aligned} \tag{F.31}$$

Finally, from (4.27a), the 'Hadamard' function B_0^1 is given by

$$\begin{aligned} B_0^1(q_1, q_2, m_1, m_2, s) \ &= \ 16(2\pi)^3\mu^{4-d}e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f}\sum_{\{\alpha\}}\int\cdots\int \mathrm{d}\Pi^{d-1}_{\mathbf{k}_1}\,\mathrm{d}\Pi_{\mathbf{k}'_1}\,\mathrm{d}\Pi_{\mathbf{k}_2}\,\mathrm{d}\Pi_{\mathbf{k}'_2} \\ &\times\ (2\pi)^3\delta^{(3)}(\mathbf{q}_1\,-\,\mathbf{k}_1\,-\,\mathbf{k}_2)(2\pi)^3\delta^{(3)}(\mathbf{q}_2\,-\,\mathbf{k}'_1\,-\,\mathbf{k}'_2) \\ &\times\ 2\mathscr{E}_1e^{i(\alpha_1E_1\,-\,\alpha'_1E'_1)\tilde{t}_f}2\mathscr{E}_2e^{i(\alpha_2E_2\,-\,\alpha'_2E'_2)\tilde{t}_f} \\ &\times\ \frac{1}{\pi}\bigg[\Big(q_1^0\,-\,q_2^0\,-\,\alpha_1E_1\,+\,\alpha'_1E'_1\,-\,\alpha_2E_2\,+\,\alpha'_2E'_2\Big)^2\,+\,4s^2\bigg]^{-1} \\ &\times\ \bigg[\Big(\frac{q_1^0+q_2^0}{2}\,-\,\frac{\alpha_1E_1+\alpha'_1E'_1}{2}\,-\,\frac{\alpha_2E_2+\alpha'_2E'_2}{2}\Big)^2\,+\,s^2\bigg]^{-1} isF^1(\{\hat{k}\},t)\ , \qquad \text{(F.32)} \end{aligned}$$

which depends only on $F^1(\{\hat{k}\},t)$, as one should expect.

Absolutely-Ordered Functions

We turn our attention now to the $(a,b) = (2,1)$ and $(a,b) = (1,2)$ elements of (F.2). These are the positive- and negative-frequency absolutely-ordered functions, respectively:

$$\begin{aligned} B_0^{>(<)}(q_1, q_2, m_1, m_2, s) \ &= \ -8(2\pi\mu)^{4-d}e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f}\int\cdots\int\frac{\mathrm{d}^dk_1}{i\pi^2}\,\frac{\mathrm{d}^4k'_1}{(2\pi)^4}\,\frac{\mathrm{d}^4k_2}{(2\pi)^4}\,\frac{\mathrm{d}^4k'_2}{(2\pi)^4} \\ &\times\ (2\pi)^3\delta^{(3)}(\mathbf{q}_1\,-\,\mathbf{k}_1\,-\,\mathbf{k}_2)(2\pi)^3\delta^{(3)}(\mathbf{q}_2\,-\,\mathbf{k}'_1\,-\,\mathbf{k}'_2) \\ &\times\ 4s\bigg[\Big(q_1^0\,-\,q_2^0\,-\,k_1^0\,+\,{k'_1}^0\,-\,k_2^0\,+\,{k'_2}^0\Big)^2\,+\,4s^2\bigg]^{-1} \\ &\times\ \bigg[\Big(q_1^0\,+\,q_2^0\,-\,k_1^0\,-\,{k'_1}^0\,-\,k_2^0\,-\,{k'_2}^0\Big)^2\,+\,4s^2\bigg)^{-1} \\ &\times\ 2\pi\delta(k_1^2\,-\,m_1^2)|2k_1^0|^{1/2}\Big(\theta((-)k_1^0,(-){k'_1}^0)(2\pi)^3\delta^{(3)}(\mathbf{k}_1\,-\,\mathbf{k}'_1) \\ &\qquad +\ \tilde{f}_1(k_1,k'_1,t)\Big)\,e^{i(k_1^0\,-\,k_1^{0\prime})\tilde{t}_f}|2{k'_1}^0|^{1/2}2\pi\delta({k'_1}^2\,-\,m_1^2) \\ &\times\ 2\pi\delta(k_2^2\,-\,m_2^2)|2k_2^0|^{1/2}\Big(\theta((-)k_2^0,(-){k'_2}^0)(2\pi)^3\delta^{(3)}(\mathbf{k}_2\,-\,\mathbf{k}'_2) \\ &\qquad +\ \tilde{f}_2^C(k_2,k'_2,t)\Big)\,e^{i(k_2^0\,-\,{k'_2}^0)\tilde{t}_f}|2k_2^0|^{1/2}2\pi\delta({k'_2}^2\,-\,m_2^2)\ , \qquad \text{(F.33)} \end{aligned}$$

where $\theta(x,y)$ is defined in (F.11).

After carrying out the zeroth-component momentum integrals, we obtain

$$\begin{aligned} B_0^{>(<)}(q_1, q_2, m_1, m_2, s) \ &= \ 16(2\pi)^3\mu^{4-d}e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f}\sum_{\{\alpha\}}\int\cdots\int \mathrm{d}\Pi^{d-1}_{\mathbf{k}_1}\,\mathrm{d}\Pi_{\mathbf{k}'_1}\,\mathrm{d}\Pi_{\mathbf{k}_2}\,\mathrm{d}\Pi_{\mathbf{k}'_2} \\ &\times\ (2\pi)^3\delta^{(3)}(\mathbf{q}_1\,-\,\mathbf{k}_1\,-\,\mathbf{k}_2)(2\pi)^3\delta^{(3)}(\mathbf{q}_2\,-\,\mathbf{k}'_1\,-\,\mathbf{k}'_2) \\ &\times\ 2\mathscr{E}_1e^{i(\alpha_1E_1\,-\,\alpha'_1E'_1)\tilde{t}_f}2\mathscr{E}_2e^{i(\alpha_2E_2\,-\,\alpha'_2E'_2)\tilde{t}_f} \\ &\times\ \frac{1}{\pi}\bigg[\Big(q_1^0\,-\,q_2^0\,-\,\alpha_1E_1\,+\,\alpha'_1E'_1\,-\,\alpha_2E_2\,+\,\alpha'_2E'_2\Big)^2\,+\,4s^2\bigg]^{-1} \\ &\times\ \bigg[\Big(\frac{q_1^0+q_2^0}{2}\,-\,\frac{\alpha_1E_1+\alpha'_1E'_1}{2}\,-\,\frac{\alpha_2E_2+\alpha'_2E'_2}{2}\Big)^2\,+\,s^2\bigg]^{-1} isF^{>(<)}(\{\hat{k}\},t)\ , \quad \text{(F.34)} \end{aligned}$$

where

$$\begin{aligned} F^{>(<)}(\{\hat{k}\},t) \;&=\; \Big(\theta((-)\alpha_1,(-)\alpha'_1)(2\pi)^3\delta^{(3)}(\mathbf{k}_1 \,-\, \mathbf{k}'_1) \,+\, \tilde{f}_1(\hat{k}_1,\hat{k}'_1,t)\Big) \\ &\quad\times\, \Big(\theta((-)\alpha_2,(-)\alpha'_2)(2\pi)^3\delta^{(3)}(\mathbf{k}_2 \,-\, \mathbf{k}'_2) \,+\, \tilde{f}_2^{C}(\hat{k}_2,\hat{k}'_2,t)\Big)\,. \end{aligned} \tag{F.35}$$

We may confirm the following relations between the F's:

$$F^{\mathrm{R}}(\{\hat{k}\},t) \;=\; F^{>}(\{\hat{k}\},t) \,-\, F^{<}(\{\hat{k}\},t)\,, \tag{F.36a}$$

$$F^{1}(\{\hat{k}\},t) \;=\; F^{>}(\{\hat{k}\},t) \,+\, F^{<}(\{\hat{k}\},t)\,, \tag{F.36b}$$

which are in line with our notation for the propagators and the self-energies.

Causal Functions

We are now in a position to obtain the causal counterparts. Given the relations in (A.6d), the retarded and advanced functions $B_0^{\mathrm{R(A)}}$ are given by

$$\begin{aligned} B_0^{\mathrm{R(A)}}(q_1,q_2,m_1,m_2,s) \;&=\; -8(2\pi)^3\mu^{4-d}e^{i(q_1^0\,-\,q_2^0)\tilde{t}_f}\sum_{\{\alpha\}}\int\cdots\int \mathrm{d}\Pi_{\mathbf{k}_1}^{d-1}\,\mathrm{d}\Pi_{\mathbf{k}'_1}\,\mathrm{d}\Pi_{\mathbf{k}_2}\,\mathrm{d}\Pi_{\mathbf{k}'_2} \\ &\quad\times\, (2\pi)^3\delta^{(3)}(\mathbf{q}_1 \,-\, \mathbf{k}_1 \,-\, \mathbf{k}_2)(2\pi)^3\delta^{(3)}(\mathbf{q}_2 \,-\, \mathbf{k}'_1 \,-\, \mathbf{k}'_2) \\ &\quad\times\, 2\mathscr{E}_1 e^{i(\alpha_1E_1\,-\,\alpha'_1E'_1)\tilde{t}_f}2\mathscr{E}_2 e^{i(\alpha_2E_2\,-\,\alpha'_2E'_2)\tilde{t}_f} \\ &\quad\times\, \frac{1}{\pi}\Big[\Big(q_1^0 \,-\, q_2^0 \,-\, \alpha_1E_1 \,+\, \alpha'_1E'_1 \,-\, \alpha_2E_2 \,+\, \alpha'_2E'_2\Big)^2 \,+\, 4s^2\Big]^{-1} \\ &\quad\times\, \Big(\tfrac{q_1^0\,+\,q_2^0}{2} \,-\, \tfrac{\alpha_1E_1\,+\,\alpha'_1E'_1}{2} \,-\, \tfrac{\alpha_2E_2\,+\,\alpha'_2E'_2}{2} \,+(-)\,is\Big)^{-1}F^{\mathrm{R(A)}}(\{\hat{k}\},t)\,, \end{aligned} \tag{F.37}$$

where $F^{\mathrm{R}}(\{\hat{k}\},t) \;=\; F^{\mathrm{A}}(\{\hat{k}\},t)$.

The Thermodynamic Equilibrium Limit

In the thermodynamic equilibrium limit, we expect to be able to recover the results from the discussions of Section 5, using the correspondence identified in (5.8). It follows that the various distribution functions satisfy the following factorization:

$$\begin{aligned} F(\{\hat{k}\},t) \;\to\; &(2\pi)^3\delta^{(3)}(\mathbf{k}_1 \,-\, \mathbf{k}'_1)\Big(\theta(\alpha_1,\alpha'_1) \,+\, \theta(-\alpha_1,-\alpha'_1)\Big) \\ &\times\, (2\pi)^3\delta^{(3)}(\mathbf{k}_2 \,-\, \mathbf{k}'_2)\Big(\theta(\alpha_2,\alpha'_2) \,+\, \theta(-\alpha_2,-\alpha'_2)\Big)\alpha_1\alpha_2F_{\mathrm{eq}}(\alpha_1E_1,\alpha_2E_2)\,, \end{aligned} \tag{F.38}$$

where

$$F_{\mathrm{eq}}^{>}(\alpha_1E_1,\alpha_2E_2) \;=\; \Big(1 \,+\, f_{\mathrm{B}}(\alpha_1E_1)\Big)\Big(1 \,+\, f_{\mathrm{B}}^{C}(\alpha_2E_2)\Big)\,, \tag{F.39a}$$

$$F_{\mathrm{eq}}^{<}(\alpha_1E_1,\alpha_2E_2) \;=\; f_{\mathrm{B}}(\alpha_1E_1)f_{\mathrm{B}}^{C}(\alpha_2E_2)\,. \tag{F.39b}$$

Using the above expressions, we may perform all but one of the three-dimensional phase-space integrations and both of the 'primed' summations in the various B_0 functions. We then obtain the following set of ‘equilibrium’ results:

$$\begin{aligned} B_0(-B_0^*)(q_1, q_2, m_1, m_2, s) \ &= \ (2\pi)^4 \frac{1}{\pi} \frac{1}{\big(q_1^0 \, - \, q_2^0\big)^2 \, + \, 4s^2} e^{i(q_1^0 \, - \, q_2^0)\tilde{t}_f} \delta^{(3)}(\mathbf{q}_1 \, - \, \mathbf{q}_2) \\ & \times \ (2\pi\mu)^{4-d} \sum_{\{\alpha\}} \int \mathrm{d}^{d-1}\mathbf{k}_1 \, \frac{1}{\pi} \, \frac{\alpha_1 \alpha_2}{E_1 E_2} \bigg[\Big(\tfrac{q_1^0 \, + \, q_2^0}{2} \, - \, \alpha_1 E_1 \, - \, \alpha_2 E_2 \Big)^2 \, + \, s^2 \bigg]^{-1} \\ & \times \ \bigg[-(+) \Big(\tfrac{q_1^0 \, + \, q_2^0}{2} \, - \, \alpha_1 E_1 \, - \, \alpha_2 E_2 \Big) \, F_{\mathrm{eq}}^{\mathrm{R}}(\alpha_1 E_1, \alpha_2 E_2) \, + \, isF_{\mathrm{eq}}^{1}(\alpha_1 E_1, \alpha_2 E_2) \bigg] \, , \end{aligned} \tag{F.40a}$$

$$\begin{aligned} B_0^{>,<,1}(q_1, q_2, m_1, m_2, s) \ &= \ (2\pi)^4 \frac{1}{\pi} \frac{1}{\big(q_1^0 \, - \, q_2^0\big)^2 \, + \, 4s^2} e^{i(q_1^0 \, - \, q_2^0)\tilde{t}_f} \delta^{(3)}(\mathbf{q}_1 \, - \, \mathbf{q}_2) \\ & \times \ (2\pi\mu)^{4-d} \sum_{\{\alpha\}} \int \mathrm{d}^{d-1}\mathbf{k}_1 \, \frac{1}{\pi} \, \frac{\alpha_1 \alpha_2}{E_1 E_2} \bigg[\Big(\tfrac{q_1^0 \, + \, q_2^0}{2} \, - \, \alpha_1 E_1 \, - \, \alpha_2 E_2 \Big)^2 \, + \, s^2 \bigg]^{-1} \\ & \times \ 2isF_{\mathrm{eq}}^{>,<,1}(\alpha_1 E_1, \alpha_2 E_2) \, , \end{aligned} \tag{F.40b}$$

$$\begin{aligned} B_0^{\mathrm{R(A)}}(q_1, q_2, m_1, m_2, s) \ &= \ -(2\pi)^4 \frac{1}{\pi} \frac{1}{\big(q_1^0 \, - \, q_2^0\big)^2 \, + \, 4s^2} e^{i(q_1^0 \, - \, q_2^0)\tilde{t}_f} \delta^{(3)}(\mathbf{q}_1 \, - \, \mathbf{q}_2) \\ & \times \ (2\pi\mu)^{4-d} \sum_{\{\alpha\}} \int \mathrm{d}^{d-1}\mathbf{k}_1 \, \frac{1}{\pi} \, \frac{\alpha_1 \alpha_2}{E_1 E_2} \Big(\tfrac{q_1^0 \, + \, q_2^0}{2} \, - \, \alpha_1 E_1 \, - \, \alpha_2 E_2 \, + (-) \, is \Big)^{-1} \\ & \times \ F_{\mathrm{eq}}^{\mathrm{R\,(A)}}(\alpha_1 E_1, \alpha_2 E_2) \, , \end{aligned} \tag{F.40c}$$

with $F_{\mathrm{eq}}^{\mathrm{R}}(\alpha_1 E_1, \alpha_2 E_2) = F_{\mathrm{eq}}^{\mathrm{A}}(\alpha_1 E_1, \alpha_2 E_2)$.

At this point, we caution the reader that (F.40a)–(F.40c) are not the exact equilibirum B_0 functions. For late times, we may use the final value theorem

$$\lim_{t \, \to \, \infty} \, F(t) \ = \ \lim_{s \, \to \, 0} \, s\mathcal{L}_t\big[F\big](s) \tag{F.41}$$

to obtain the true time-invariant equilibrium functions:

$$\begin{aligned} B_0(-B_0^*)(q_1, q_2, m_1, m_2) \ &= \ (2\pi)^4 \delta^{(4)}(q_1 \, - \, q_2)(2\pi\mu)^{4-d} \sum_{\{\alpha\}} \int \mathrm{d}^{d-1}\mathbf{k}_1 \, \frac{\alpha_1 \alpha_2}{E_1 E_2} \\ & \times \ \bigg(-(+) \frac{1}{2\pi} \, \frac{1}{q_1^0 \, - \, \alpha_1 E_1 \, - \, \alpha_2 E_2} \, F_{\mathrm{eq}}^{\mathrm{R}}(\alpha_1 E_1, \alpha_2 E_2) \\ & + \ \frac{i}{2} \, \delta(q_1^0 \, - \, \alpha_1 E_1 \, - \, \alpha_2 E_2) F_{\mathrm{eq}}^{1}(\alpha_1 E_1, \alpha_2 E_2) \bigg) \, , \end{aligned} \tag{F.42a}$$

$$\begin{aligned} B_0^{>,<,1}(q_1, q_2, m_1, m_2) \ &= \ i(2\pi)^4 \delta^{(4)}(q_1 \, - \, q_2)(2\pi\mu)^{4-d} \sum_{\{\alpha\}} \int \mathrm{d}^{d-1}\mathbf{k}_1 \, \frac{\alpha_1 \alpha_2}{E_1 E_2} \\ & \times \ \delta(q_1^0 \, - \, \alpha_1 E_1 \, - \, \alpha_2 E_2) F_{\mathrm{eq}}^{>,<,1}(\alpha_1 E_1, \alpha_2 E_2) \, , \end{aligned} \tag{F.42b}$$

$$\begin{aligned} B_0^{\mathrm{R(A)}}(q_1, q_2, m_1, m_2) \ &= \ -(2\pi)^4 \delta^{(4)}(q_1 \, - \, q_2)(2\pi\mu)^{4-d} \sum_{\{\alpha\}} \int \mathrm{d}^{d-1}\mathbf{k}_1 \, \frac{\alpha_1 \alpha_2}{E_1 E_2} \\ & \times \ \frac{1}{2\pi} \, \frac{1}{q_1^0 \, - \, \alpha_1 E_1 \, - \, \alpha_2 E_2 \, + (-) \, i\epsilon} \, F_{\mathrm{eq}}^{\mathrm{R}}(\alpha_1 E_1, \alpha_2 E_2) \, . \end{aligned} \tag{F.42c}$$

The above expressions are consistent with known results calculated in the ITF or equilibrium CTP formalism (see Appendix B).

The Homogeneous Limit

Lastly, we summarize the time-dependent homogeneous limit of the B_0 functions. In the spatially homogeneous case, the F's satisfy the following factorization

$$
\begin{aligned}
F(\{\hat{k}\},t) \;&\to\; (2\pi)^3\delta^{(3)}(\mathbf{k}_1 - \mathbf{k}_1')\Big(\,\theta(\alpha_1,\alpha_1') + \theta(-\alpha_1,-\alpha_1')\,\Big) \qquad \text{(F.43)}\\
&\quad\times (2\pi)^3\delta^{(3)}(\mathbf{k}_2 - \mathbf{k}_2')\Big(\,\theta(\alpha_2,\alpha_2') + \theta(-\alpha_2,-\alpha_2')\,\Big)F_{\text{hom}}(\alpha_1,\alpha_2,\mathbf{k}_1,\mathbf{k}_2)\;.
\end{aligned}
$$

The set of homogeneous distributions F_{hom} can be obtained from

$$
\begin{aligned}
F_{\text{hom}}^{>(<)}(\alpha_1,\alpha_2,\mathbf{k}_1,\mathbf{k}_2,t) \;&=\; \Big(\,\theta((-)\alpha_1) + \theta(\alpha_1)f\big(|\mathbf{k}_1|,t\big) + \theta(-\alpha_1)f^C\big(|\mathbf{k}_1|,t\big)\Big) \qquad \text{(F.44)}\\
&\quad\times \Big(\,\theta((-)\alpha_2) + \theta(\alpha_2)f^C\big(|\mathbf{k}_2|,t\big) + \theta(-\alpha_2)f\big(|\mathbf{k}_2|,t\big)\Big)\;,
\end{aligned}
$$

using the relations in (F.36). Substituting these distributions into the non-homogeneous B_0 functions, we perform all but one of the three-dimensional phase-space integrals and both 'primed' summations. Subsequently, with the aid of an inverse Laplace transformation with respect to s, the following set of t-dependent homogeneous B_0 functions are obtained:

$$
\begin{aligned}
B_0(-B_0^*)(q_1,q_2,m_1,m_2,\tilde{t}_f;\tilde{t}_i) \;&=\; (2\pi)^4\delta^{(3)}(\mathbf{q}_1 - \mathbf{q}_2)(2\pi\mu)^{4-d}e^{i(q_1^0 - q_2^0)\tilde{t}_f} \qquad \text{(F.45a)}\\
&\times \sum_{\{\alpha\}}\int \mathrm{d}^{d-1}\mathbf{k}_1\,\frac{1}{E_1E_2}\bigg[-(+)\frac{1}{2\pi}\,\frac{\frac{q_1^0+q_2^0}{2} - \alpha_1E_1 - \alpha_2E_2}{\big(q_1^0 - \alpha_1E_1 - \alpha_2E_2\big)\big(q_2^0 - \alpha_1E_1 - \alpha_2E_2\big)}\\
&\times \Big(\,\delta_t(q_1^0 - q_2^0) - \delta_t\big(q_1^0 + q_2^0 - 2\alpha_1E_1 - 2\alpha_2E_2\big)\,\Big)F_{\text{hom}}^{\text{R}}(\alpha_1,\alpha_2,\mathbf{k}_1,\mathbf{q}_1-\mathbf{k}_1,t)\\
&+ \frac{i}{2}\,\delta_t\big(q_1^0 - \alpha_1E_1 - \alpha_2E_2\big)\delta_t\big(q_2^0 - \alpha_1E_1 - \alpha_2E_2\big)F_{\text{hom}}^{1}(\alpha_1,\alpha_2,\mathbf{k}_1,\mathbf{q}_1-\mathbf{k}_1,t)\bigg]\;,\\
B_0^{>,<,1}(q_1,q_2,m_1,m_2,\tilde{t}_f;\tilde{t}_i) \;&=\; i(2\pi)^4\delta^{(3)}(\mathbf{q}_1 - \mathbf{q}_2)(2\pi\mu)^{4-d}e^{i(q_1^0 - q_2^0)\tilde{t}_f}\\
&\times \sum_{\{\alpha\}}\int \mathrm{d}^{d-1}\mathbf{k}_1\,\frac{1}{E_1E_2}\,\delta_t\big(q_1^0 - \alpha_1E_1 - \alpha_2E_2\big)\delta_t\big(q_2^0 - \alpha_1E_1 - \alpha_2E_2\big)\\
&\times F_{\text{hom}}^{>,<,1}(\alpha_1,\alpha_2,\mathbf{k}_1,\mathbf{q}_1-\mathbf{k}_1,t)\;, \qquad \text{(F.45b)}\\
B_0^{\text{R(A)}}(q_1,q_2,m_1,m_2,\tilde{t}_f;\tilde{t}_i) \;&=\; (2\pi)^4\delta^{(3)}(\mathbf{q}_1 - \mathbf{q}_2)(2\pi\mu)^{4-d}e^{i(q_1^0 - q_2^0)\tilde{t}_f}\\
&\times \sum_{\{\alpha\}}\int \mathrm{d}^{d-1}\mathbf{k}_1\,\frac{1}{E_1E_2}\bigg[-\frac{1}{2\pi}\,\frac{\frac{q_1^0+q_2^0}{2} - \alpha_1E_1 - \alpha_2E_2}{\big(q_1^0 - \alpha_1E_1 - \alpha_2E_2\big)\big(q_2^0 - \alpha_1E_1 - \alpha_2E_2\big)}\\
&\times \Big(\,\delta_t(q_1^0 - q_2^0) - \delta_t\big(q_1^0 + q_2^0 - 2\alpha_1E_1 - 2\alpha_2E_2\big)\,\Big)\\
&+(-)\,\frac{i}{2}\,\delta_t\big(q_1^0 - \alpha_1E_1 - \alpha_2E_2\big)\delta_t\big(q_2^0 - \alpha_1E_1 - \alpha_2E_2\big)\bigg]\\
&\times F_{\text{hom}}^{\text{R(A)}}(\alpha_1,\alpha_2,\mathbf{k}_1,\mathbf{q}_1-\mathbf{k}_1,t)\;. \qquad \text{(F.45c)}
\end{aligned}
$$

In the limit $t \to \infty$, using (3.51) and making the replacement

$$
F_{\text{hom}}(\alpha_1,\alpha_2,\mathbf{k}_1,\mathbf{q}_1-\mathbf{k}_1,t) \;\to\; \alpha_1\alpha_2F_{\text{eq}}(\alpha_1,\alpha_2,\mathbf{k}_1,\mathbf{q}_1-\mathbf{k}_1)\;, \qquad \text{(F.46)}
$$

we recover the time-independent equilibrium results in (F.42).

References

[1] J.-P. Blaizot and E. Iancu, *The quark gluon plasma: Collective dynamics and hard thermal loops*, *Phys.Rep.* **359** (2002) 355–528, [hep-ph/0101103].

[2] **ATLAS** Collaboration, M. Spousta, *Jet reconstruction and jet quenching in heavy ion collisions at ATLAS*, arXiv:1106.1604.

[3] **CMS** Collaboration, M. B. Tonjes, *Study of jet quenching using dijets in Pb Pb collisions with CMS*, *J.Phys.G* **G38** (2011) 124084.

[4] **ALICE** Collaboration, J. F. Grosse-Oetringhaus, *Measurement of jet quenching with I_{CP} and I_{AA}, Pythia in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76\,\mathrm{TeV}$ with ALICE*, arXiv:1105.3901.

[5] **WMAP Collaboration** Collaboration, D. N. Spergel et al., *First year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Determination of cosmological parameters*, *Astrophys.J.Suppl.* **148** (2003) 175–194, [astro-ph/0302209].

[6] **WMAP Collaboration** Collaboration, D. N. Spergel et al., *Wilkinson Microwave Anisotropy Probe (WMAP) three year results: implications for cosmology*, *Astrophys.J.Suppl.* **170** (2007) 377, [astro-ph/0603449].

[7] G. Steigman, *Neutrinos and big bang nucleosynthesis*, *Phys. Scripta* **T121** (2005) 142–146, [hep-ph/0501100].

[8] A. D. Sakharov, *Violation of CP invariance, C asymmetry, and baryon asymmetry of the Universe*, *Pisma Zh.Eksp.Teor.Fiz.* **5** (1967) 32–35.

[9] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, *On the anomalous electroweak baryon number nonconservation in the early Universe*, *Phys.Lett.* **B155** (1985) 36.

[10] M. Fukugita and T. Yanagida, *Baryogenesis without grand unification*, *Phys.Lett.* **B174** (1986) 45–47.

[11] M. Fukugita and T. Yanagida, *Sphaleron-induced baryon-number nonconservation and a constraint on Majorana neutrino masses*, *Phys.Rev.* **D42** (1990) 1285–1286.

[12] L. Kofman, A. D. Linde, and A. A. Starobinsky, *Reheating after inflation*, *Phys.Rev.Lett.* **73** (1994) 3195–3198, [hep-th/9405187].

[13] D. Boyanovsky, H. J. de Vega, R. Holman, D. S. Lee, and A. Singh, *Dissipation via particle production in scalar field theories*, *Phys.Rev.* **D51** (1995) 4419–4444, [hep-ph/9408214].

[14] J. Baacke, K. Heitmann, and C. Pätzold, *Nonequilibrium dynamics: Preheating in the $SU(2)$ Higgs model*, *Phys.Rev.* **D55** (1997) 7815–7825, [hep-ph/9612264].

[15] L. Kofman, A. D. Linde, and A. A. Starobinsky, *Towards the theory of reheating after inflation*, *Phys.Rev.* **D56** (1997) 3258–3295, [hep-ph/9704452].

[16] C. M. Bender and S. Sarkar, *Asymptotic analysis of the Boltzmann equation for dark matter relics*, arXiv:1203.1822.

[17] E. W. Kolb and S. Wolfram, *Baryon number generation in the early universe*, *Nucl.Phys.* **B172** (1980) 224–284.

[18] M. S. Carena, M. Quiros, M. Seco, and C. E. M. Wagner, *Improved results in supersymmetric electroweak baryogenesis*, *Nucl.Phys.* **B650** (2003) 24–42, [hep-ph/0208043].

[19] G. F. Giudice, A. Notari, M. Raidal, A. Riotto, and A. Strumia, *Towards a complete theory of thermal leptogenesis in the SM and MSSM*, *Nucl.Phys.* **B685** (2004) 89–149, [hep-ph/0310123].

[20] A. Pilaftsis and T. E. J. Underwood, *Resonant leptogenesis*, *Nucl.Phys.* **B692** (2004) 303–345, [hep-ph/0309342].

[21] W. Buchmüller, P. Di Bari, and M. Plümacher, *Leptogenesis for pedestrians*, *Annals Phys.* **315** (2005) 305–351, [hep-ph/0401240].

[22] A. Pilaftsis and T. E. J. Underwood, *Electroweak-scale resonant leptogenesis*, *Phys.Rev.* **D72** (2005) 113001, [hep-ph/0506107].

[23] S. Davidson, E. Nardi, and Y. Nir, *Leptogenesis*, *Phys.Rep.* **466** (2008) 105–177, [arXiv:0802.2962].

[24] F. F. Deppisch and A. Pilaftsis, *Lepton flavour violation and θ_{13} in minimal resonant leptogenesis*, *Phys.Rev.* **D83** (2011) 076007, [arXiv:1012.1834].

[25] T. Matsubara, *A new approach to quantum statistical mechanics*, *Prog.Theor.Phys.* **14** (1955) 351–378.

[26] K.-C. Chou, Z.-B. Su, B.-L. Hao, and L. Yu, *Equilibrium and nonequilibrium formalisms made unified*, *Phys.Rep.* **118** (1985) 1–131.

[27] N. P. Landsman and C. G. van Weert, *Real- and imaginary-time field theory at finite temperature and density*, *Phys.Rep.* **145** (1987) 141–249.

[28] J. S. Schwinger, *Brownian motion of a quantum oscillator*, *J.Math.Phys.* **2** (1961) 407–432.

[29] L. Keldysh, *Diagram technique for nonequilibrium processes*, *Zh.Eksp.Teor.Fiz.* **47** (1964) 1515–1527.

[30] R. Kobes, *A correspondence between imaginary-time and real-time finite-temperature field theory*, *Phys.Rev.* **D42** (1990) 562–572.

[31] P. Aurenche and T. Becherrawy, *A comparison of the real-time and the imaginary-time formalisms of finite-temperature field theory for 2, 3, and 4 point green functions*, *Nucl.Phys.* **B379** (1992) 259–303.

[32] H.-H. Xu, *Relation between real-time and imaginary-time formalisms of finite-temperature quantum field theory*, *Phys.Rev.* **D47** (1993) 2622–2624.

[33] R. Baier and A. Niégawa, *Analytic continuation of thermal N-point functions from imaginary to real energies*, *Phys.Rev.* **D49** (1994) 4107–4112, [hep-ph/9307362].

[34] T. S. Evans and A. C. Pearson, *A reexamination of the path-ordered approach to real time thermal field theory*, *Phys.Rev.* **D52** (1995) 4652–4659, [hep-ph/9412217].

[35] M. A. van Eijck, R. Kobes, and C. G. van Weert, *Transformations of real-time finite-temperature Feynman rules*, *Phys.Rev.* **D50** (1994) 4097–4109, [hep-ph/9406214].

[36] B.-R. Zhou, *Explicit proof of equivalence of two-point functions in the two formalisms of thermal field theory*, *Commun.Theor.Phys.* **37** (2002) 49–54, [hep-th/0108146].

[37] J. M. Cornwall, R. Jackiw, and E. Tomboulis, *Effective action for composite operators*, *Phys.Rev.* **D10** (1974) 2428–2445.

[38] M. E. Carrington, *The 4PI effective action for φ^4 theory*, *Eur.Phys.J.* **C35** (2004) 383–392, [hep-ph/0401123].

[39] E. Calzetta and B. L. Hu, *Closed-time-path functional formalism in curved spacetime: application to cosmological back-reaction problems*, *Phys.Rev.* **D35** (1987) 495–509.

[40] E. Calzetta and B. L. Hu, *Nonequilibrium quantum fields: closed-time-path effective action, Wigner function and Boltzmann equation*, *Phys.Rev.* **D37** (1988) 2878–2900.

[41] G. Aarts and J. Berges, *Nonequilibrium time evolution of the spectral function in quantum field theory*, *Phys.Rev.* **D64** (2001) 105010, [hep-ph/0103049].

[42] T. Prokopec, M. G. Schmidt, and S. Weinstock, *Transport equations for chiral fermions to order $\hbar$ and electroweak baryogenesis. Part I*, *Annals Phys.* **314** (2004) 208–265, [hep-ph/0312110].

[43] T. Prokopec, M. G. Schmidt, and S. Weinstock, *Transport equations for chiral fermions to order $\hbar$ and electroweak baryogenesis. Part II*, *Annals Phys.* **314** (2004) 267–320, [hep-ph/0406140].

[44] M. Lindner and M. M. Müller, *Comparison of Boltzmann equations with quantum dynamics for scalar fields*, *Phys.Rev.* **D73** (2006) 125002, [hep-ph/0512147].

[45] A. De Simone and A. Riotto, *Quantum Boltzmann equations and leptogenesis*, *JCAP* **0708** (2007) 002, [hep-ph/0703175].

[46] V. Cirigliano, A. De Simone, G. Isidori, I. Masina, and A. Riotto, *Quantum resonant leptogenesis and minimal lepton flavour violation*, *JCAP* **0801** (2008) 004, [arXiv:0711.0778].

[47] M. Garny, A. Hohenegger, A. Kartavtsev, and M. Lindner, *Systematic approach to leptogenesis in nonequilibrium QFT: self-energy contribution to the CP-violating parameter*, *Phys.Rev.* **D81** (2010) 085027, [arXiv:0911.4122].

[48] V. Cirigliano, C. Lee, M. J. Ramsey-Musolf, and S. Tulin, *Flavored quantum Boltzmann equations*, *Phys.Rev.* **D81** (2010) 103503, [arXiv:0912.3523].

[49] M. Beneke, B. Garbrecht, C. Fidler, M. Herranen, and P. Schwaller, *Flavoured leptogenesis in the CTP formalism*, *Nucl.Phys.* **B843** (2011) 177–212, [arXiv:1007.4783].

[50] A. Anisimov, W. Buchmüller, M. Drewes, and S. Mendizabal, *Quantum leptogenesis I*, *Annals Phys.* **326** (2011) 1998–2038, [arXiv:1012.5821].

[51] J. Winter, *Wigner transformation in curved space-time and the curvature correction of the Vlasov equation for semiclassical gravitating systems*, *Phys.Rev.* **D32** (1985) 1871–1888.

[52] G. Baym and L. P. Kadanoff, *Conservation laws and correlation functions*, *Phys.Rev.* **124** (1961) 287–299.

[53] L. P. Kadanoff and G. Baym, *Quantum statistical mechanics; Green's function methods in equilibrium and nonequilibrium problems.* Addison-Wesley Publishing Company, 1989.

[54] P. Danielewicz, *Quantum theory of nonequilibrium processes, I*, *Annals Phys.* **152** (1984) 239–304.

[55] T. Bornath, D. Kremp, W. D. Kraeft, and M. Schlanges, *Kinetic equations for a nonideal quantum system*, *Phys.Rev.* **E54** (1996) 3274–3284.

[56] J. Berges and S. Schlichting, *The non-linear glasma*, arXiv:1209.0817.

[57] T. Gasenzer, S. Kessler, and J. M. Pawlowski, *Far-from-equilibrium quantum many-body dynamics*, *Eur.Phys.J.* **C70** (2010) 423–443, [arXiv:1003.4163].

[58] G. Källén, *On the definition of the renormalization constants in quantum electrodynamics*, *Helv.Phys.Acta* **25** (1952) 417–434.

[59] H. Lehmann, *On the properties of propagation functions and renormalization contants of quantized fields*, *Nuovo Cim.* **11** (1954) 342–357.

[60] W. Greiner and J. Reinhardt, *Field quantization*, pp. 138–140. Springer, 1996.

[61] H. Lehmann, K. Symanzik, and W. Zimmermann, *On the formulation of quantized field theories*, *Nuovo Cim.* **1** (1955) 205–225.

[62] G. 't Hooft and M. J. G. Veltman, *Diagrammar*, *NATO Adv. Study Inst. Ser. B Phys.* **4** (1974) 177–322.

[63] R. L. Kobes and G. W. Semenoff, *Discontinuities of Green functions in field theory at finite temperature and density*, *Nucl.Phys.* **B260** (1985) 714–746.

[64] R. L. Kobes and G. W. Semenoff, *Discontinuities of Green functions in field theory at finite temperature and density (II)*, *Nucl.Phys.* **B272** (1986) 329–364.

[65] M. J. G. Veltman, *Diagrammatica; the path to Feynman diagrams*, pp. 183–206. Cambridge University Press, 1994.

[66] R. Jackiw, *Functional evaluation of the effective potential*, *Phys.Rev.* **D9** (1974) 1686.

[67] J. Berges and S. Borsányi, *Range of validity of transport equations*, *Phys.Rev.* **D74** (2006) 045022, [hep-ph/0512155].

[68] M. Le Bellac, *Thermal field theory*, pp. 54–58. Cambridge University Press, 2000.

[69] R. Kobes, *Retarded functions, dispersion relations, and Cutkosky rules at zero and finite temperature*, *Phys.Rev.* **D43** (1991) 1269–1282.

[70] M. A. van Eijck and C. G. van Weert, *Finite-temperature retarded and advanced Green functions*, *Phys.Lett.* **B278** (1992) 305–310.

[71] T. Altherr and D. Seibert, *Problems of perturbation series in nonequilibrium quantum field theories*, *Phys.Lett.* **B333** (1994) 149–152, [hep-ph/9405396].

[72] C. Greiner and S. Leupold, *Interpretation and resolution of pinch singularities in non-equilibrium quantum field theory*, *Eur.Phys.J.* **C8** (1999) 517–522, [hep-ph/9804239].

[73] P. F. Bedaque, *Thermalization and pinch singularities in non-equilibrium quantum field theory*, *Phys.Lett.* **B344** (1995) 23–28, [hep-ph/9410415].

[74] I. Dadić, *Out-of-equilibrium thermal field theories: finite time after switching on the interaction and Wigner transforms of the projected functions*, *Phys.Rev.* **D63** (2001) 025011, [hep-ph/9910337].

[75] E. Kolb and M. Turner, *The Early Universe*, p. 116. Westview Press, 1994.

[76] E. E. Burniston and C. E. Siewert, *The use of Riemann problems in solving a class of transcendental equations*, *Proc.Camb.Phil.Soc.* **73** (1973) 111–118.

[77] D. Boyanovsky, H. J. de Vega, R. Holman, S. P. Kumar, and R. D. Pisarski, *Real-time relaxation and kinetics in hot scalar QED: Landau damping*, *Phys.Rev.* **D58** (1998) 125009, [hep-ph/9802370].

[78] S. J. Huber and M. G. Schmidt, *Electroweak baryogenesis: concrete in a SUSY model with a gauge singlet*, *Nucl.Phys.* **B606** (2001) 183–230, [hep-ph/0003122].

[79] S. J. Huber and M. Sopena, *The bubble wall velocity in the minimal supersymmetric light stop scenario*, *Phys.Rev.* **D85** (2012) 103507, [arXiv:1112.1888].

[80] J. Berges, S. Borsányi, and C. Wetterich, *Prethermalization*, *Phys.Rev.Lett.* **93** (2004) 142002, [hep-ph/0403234].

[81] J. Berges, S. Borsányi, and C. Wetterich, *Isotropization far from equilibrium*, *Nucl.Phys.* **B727** (2005) 244–263, [hep-ph/0505182].

[82] G. Sigl and G. Raffelt, *General kinetic description of relativistic mixed neutrinos*, *Nucl.Phys.* **B406** (1993) 423–451.

[83] D. Boyanovsky and C. M. Ho, *Nonequilibrium dynamics of mixing, oscillations and equilibration: a model study*, *Phys.Rev.* **D75** (2007) 085004, [hep-ph/0610036].

[84] G. 't Hooft and M. J. G. Veltman, *Scalar one loop integrals*, *Nucl.Phys.* **B153** (1979) 365–401.